

**American College of Radiology
ACR Appropriateness Criteria®**

Clinical Condition: Dysphagia

Variant 1: Oropharyngeal dysphagia with an attributable cause.

Radiologic Procedure	Rating	Comments	RRL*
X-ray barium swallow modified	8		Med
X-ray pharynx dynamic and static imaging	6		Med
X-ray biphasic esophagram	4	Double contrast and single contrast.	Med
X-ray barium swallow	4		Med
NUC Tc-99m transit scintigraphy esophagus	2		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 2: Unexplained oropharyngeal dysphagia.

Radiologic Procedure	Rating	Comments	RRL*
X-ray pharynx dynamic and static imaging	8	Both pharyngeal and esophageal exams needed, since patient may have referred dysphagia.	Med
X-ray biphasic esophagram	8	Both pharyngeal and esophageal exams needed, since patient may have referred dysphagia. Double contrast and single contrast.	Med
X-ray barium swallow modified	6		Med
X-ray barium swallow	6		Med
NUC Tc-99m transit scintigraphy esophagus	4		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Dysphagia****Variant 3:****Substernal dysphagia in immunocompetent patients.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray biphasic esophagram	8	Endoscopy and biphasic esophagram are both excellent diagnostic tests in this setting. Double contrast and single contrast.	Med
X-ray barium swallow	6	Probably indicated if that is all the patient can do.	Med
X-ray barium swallow modified	4		Med
NUC Tc-99m transit scintigraphy esophagus	4		Med
X-ray pharynx dynamic and static imaging	4		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 4:**Substernal dysphagia in immunocompromised patients.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray biphasic esophagram	8	Endoscopy and biphasic esophagram are both excellent diagnostic tests in this setting. Double contrast and single contrast.	Med
X-ray barium swallow	5		Med
X-ray barium swallow modified	4		Med
X-ray pharynx dynamic and static imaging	3		Med
NUC Tc-99m transit scintigraphy esophagus	2		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

DYSPHAGIA

Expert Panel on Gastrointestinal Imaging: Marc S. Levine, MD¹; Robert L. Bree, MD, MHSA²; Max Paul Rosen, MD, MPH³; W. Dennis Foley, MD⁴; Spencer B. Gay, MD⁵; Thomas H. Grant DO⁶; Jay P. Heiken, MD⁷; James E. Huprich, MD⁸; Tasneem Lalani, MD⁹; Frank H. Miller, MD¹⁰; Pablo R. Ros, MD, MPH¹¹; Gary S. Sudakoff, MD¹²; Frederick L. Greene, MD¹³; Don C. Rockey, MD.¹⁴

Summary of Literature Review

Dysphagia is defined as the subjective awareness of swallowing difficulty during passage of a solid or liquid bolus from the mouth to the stomach. This symptom can be caused by functional or structural abnormalities of the oral cavity, pharynx, esophagus, or even the gastric cardia. A barium study may be performed with videofluoroscopy to assess pharyngeal function and esophageal motility, as well as a series of double-contrast and single-contrast static images to assess structural abnormalities such as rings, strictures, and tumors. Other possible diagnostic tests include a modified barium swallow, endoscopy, manometry, and nuclear scintigraphy esophageal transit studies. The choice of test may depend on the clinical setting as well as the nature and location of the patient's dysphagia.

Clinical Perspective

Many patients with dysphagia can subjectively localize a sensation of blockage or discomfort to the throat or substernal region. Patients with pharyngeal dysphagia typically complain of food sticking in the throat or of a globus sensation with a lump in the throat. Other symptoms of oropharyngeal dysfunction include coughing or choking during swallowing due to laryngeal penetration or aspiration, a nasal-quality voice or nasal regurgitation due to soft-palate insufficiency, and food dribbling from the mouth or difficulty chewing due to an abnormal oral phase of swallowing. When oropharyngeal dysphagia has an attributable cause (eg, recent stroke), a modified barium swallow may be the appropriate test to assess the patient's swallowing status and initiate treatment by a speech therapist. In patients with unexplained oropharyngeal dysphagia, however, a more

detailed barium study may be needed to determine the cause. It also is important to recognize that abnormalities of the mid or distal esophagus or even the gastric cardia may cause referred dysphagia to the upper chest or pharynx, whereas abnormalities of the pharynx rarely cause referred dysphagia to the lower chest [1]. The esophagus and cardia should therefore be evaluated in patients with pharyngeal symptoms, particularly if no abnormalities are found in the pharynx to explain these symptoms. Thus, a combined radiologic examination of the pharynx, esophagus, and gastric cardia is appropriate for patients with unexplained pharyngeal dysphagia.

Other patients may have substernal dysphagia with a sensation of blockage or discomfort anywhere from the thoracic inlet to the xiphoid process. This symptom may be caused by esophageal motility disorders or by structural abnormalities of the esophagus or cardia such as esophagitis, rings, strictures, and tumors. When barium studies are performed on these patients, the esophagram usually consists of a biphasic examination that includes upright double-contrast views with a high-density barium suspension to assess mucosal disease and prone single-contrast views with a low-density barium suspension to assess distensibility and motility.

Optimal evaluation of patients with dysphagia depends on the nature and location of the dysphagia and the clinical setting. The following four scenarios are considered separately:

1. Oropharyngeal dysphagia with an attributable cause;
2. Unexplained oropharyngeal dysphagia;
3. Substernal dysphagia in immunocompetent patients; and
4. Substernal dysphagia in immunocompromised patients.

Oropharyngeal Dysphagia with an Attributable Cause

When oropharyngeal dysphagia has an attributable cause (eg, recent stroke, worsening dementia, myasthenia gravis, amyotrophic lateral sclerosis), a modified barium swallow may be performed with the assistance of a speech therapist [2]. The study is facilitated by examining the patient in a speech therapy chair. The modified barium swallow focuses on the oral cavity, pharynx, and cervical esophagus with videofluoroscopy or cine recording to assess abnormalities of both the oral phase of swallowing (eg, difficulty propelling the bolus) and the pharyngeal phase (eg, laryngeal penetration, tracheal aspiration, cricopharyngeal dysfunction). It has been shown that the risk of developing aspiration pneumonia is directly related to the degree of swallowing dysfunction on video fluoroscopic studies [3]. The patient may be given high-density and low-density barium suspensions as well as

¹Principal Author, Hospital of the University of Pennsylvania, Philadelphia, Pa; ²Panel Chair, University of Washington, Seattle, Wash; ³Panel Vice-Chair, University of Pennsylvania, Philadelphia, Pa; ⁴Froedtert Hospital East, Milwaukee, Wis; ⁵University of Virginia Health Science Center, Charlottesville, Va; ⁶Northwestern University Feinberg School of Medicine/NMH, Chicago, Ill; ⁷Mallinckrodt Institute of Radiology, Saint Louis, Mo; ⁸Mayo Clinic, Rochester, Minn; ⁹Inland Imaging Associates, Seattle, Wash; ¹⁰Northwestern University Feinberg School of Medicine/NMH, Chicago, Ill; ¹¹Brigham & Women's Hospital/Harvard, Boston, Mass; ¹²Froedtert Hospital, Milwaukee, Wis; ¹³Carolinas Medical Center, Charlotte, NC, American College of Surgeons; ¹⁴University of Texas, Southwest Medical Center, Dallas, Texas, American Gastroenterological Association.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

other substances of varying consistency (eg, barium paste or barium-impregnated crackers) to assess the patient's ability to swallow solid or semisolid substances. In conjunction with a speech therapist, various compensatory maneuvers (eg, a chin-tuck position) may be tried to prevent aspiration or other types of swallowing dysfunction [2].

Unexplained Oropharyngeal Dysphagia

In patients with unexplained oropharyngeal dysphagia, a more detailed barium study may be performed to assess both functional and structural abnormalities of the pharynx [4,5]. As in the modified barium swallow, a dynamic examination of the pharynx with videofluoroscopy or cine recording permits assessment of both the oral and pharyngeal phases of swallowing. However, static images of the pharynx (eg, double-contrast spot films of the pharynx in frontal and lateral projections with high-density barium) should also be obtained to detect structural abnormalities (eg, pharyngeal tumors, Zenker's diverticulum). Because some patients with lesions in the esophagus or at the gastric cardia can have referred dysphagia, the esophagus and cardia should also be carefully evaluated as part of the barium study in these patients, particularly if no abnormalities are found in the pharynx to account for their symptoms (see below) [4]. In patients with unexplained pharyngeal dysphagia, it has been shown that the combination of videofluoroscopy and static images of the pharynx and esophagus has a higher diagnostic value than either videofluoroscopy or static images alone [6].

Substernal Dysphagia in Immunocompetent Patients

The biphasic esophagram is a valuable technique for evaluating substernal dysphagia in immunocompetent patients [4]. This technique permits detection of both structural and functional abnormalities of the esophagus. Perhaps the most important structural lesion is carcinoma of the esophagus or esophagogastric junction. In one study, double-contrast esophagography was found to have a sensitivity of 96% in diagnosing cancer of the esophagus or esophagogastric junction [7], which is comparable to the reported sensitivity of endoscopy for diagnosing these lesions. In two other large series of patients, endoscopy failed to reveal any cases of esophageal carcinoma that had been missed on the barium studies [8,9]. The findings in these series suggest that endoscopy is not routinely warranted to rule out missed tumors in patients who have normal findings on radiologic examinations.

While double-contrast views are best for detecting mucosal lesions (eg, tumors, esophagitis), prone single-contrast views with continuous drinking of a low-density barium suspension are best for detecting lower esophageal rings or strictures. It has been shown that lower esophageal rings are two to three times more likely to be

diagnosed on prone single-contrast views than on upright double-contrast views because of inadequate distention of the distal esophagus when the patient is upright [10,11]. In one study, the biphasic esophagram was found to detect about 95% of all lower esophageal rings, whereas endoscopy detected only 76% of these rings [11]. Similarly, biphasic esophagrams have been found to have a sensitivity of about 95% in detecting peptic strictures, sometimes revealing strictures that are missed with endoscopy [12,13].

Alternatively, endoscopy may be performed to evaluate the esophagus for structural abnormalities in patients with dysphagia. It is a highly accurate test for esophageal cancer when multiple endoscopic biopsy specimens and brushings are obtained. It also is more sensitive than double-contrast esophagography for detecting mild reflux esophagitis or other subtle forms of esophagitis. However, endoscopy is a more expensive and invasive test than the barium study. It also is less sensitive than the barium study for detecting lower esophageal rings or strictures (see above) [10-13] and does not permit evaluation of esophageal motility disorders. For these reasons, the barium study is often recommended, even by gastroenterologists, as the initial diagnostic test for patients with dysphagia [4,14-17].

The biphasic esophagram is also a useful test in patients with esophageal motility disorders causing dysphagia. Videofluoroscopy of discrete swallows of a low-density barium suspension in the prone right anterior oblique position permits detailed assessment of esophageal motility. In various studies, videofluoroscopy has been found to have an overall sensitivity of 80%-89% and specificity of 79%-91% for diagnosing esophageal motility disorders (eg, achalasia, diffuse esophageal spasm) in comparison to esophageal manometry [18,19]. Occasionally, barium studies may even reveal dysmotility not seen at manometry (eg, some patients with the beaklike distal esophageal narrowing of achalasia are found to have complete relaxation of the lower esophageal sphincter on manometry) [20]. In any case, when a significant esophageal motility disorder is detected on a barium study, manometry may be performed to further elucidate the nature of this motility disorder. Alternatively, radionuclide esophageal transit scintigraphy is a simple, noninvasive, and quantitative test of esophageal motility and emptying [21-23].

Substernal Dysphagia in Immunocompromised Patients

The major consideration in immunocompromised patients with dysphagia or odynophagia (painful swallowing) is infectious esophagitis, most commonly due to *Candida albicans* or herpes simplex virus. In HIV-positive patients, *Candida* is the cause of esophageal symptoms in the majority of cases, with cytomegalovirus (CMV), herpes

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

simplex, and idiopathic ulcers (also known as HIV ulcers) the other most common etiologies [24-27]. HIV-positive patients with esophageal symptoms are generally treated empirically with antifungal therapy without undergoing a diagnostic examination. Most gastroenterologists prefer that those with persistent symptoms (or severe symptoms at presentation) be evaluated by endoscopy [28,29]. Endoscopy is preferred because of the ability to obtain specimens (eg, histology, cytology, immunostaining, and culture).

The endoscopic or radiographic appearance alone does not accurately predict diseases other than Candida esophagitis; diagnosis requires the acquisition of specimens for laboratory study [26,27]. Barium esophagography is preferred in some centers and can be useful in guiding management. Double-contrast esophagography is more accurate than single-contrast esophagography for detecting ulcers or plaques associated with infectious esophagitis [30-35]. However, single-contrast esophagrams may be performed if the patient is too sick or debilitated to tolerate a double-contrast examination. Patients with radiographically diagnosed Candida or herpes esophagitis may be treated with antifungal or antiviral agents without endoscopic evaluation, but endoscopy is warranted for patients with giant esophageal ulcers in order to differentiate CMV and HIV, so that appropriate therapy can be started [34].

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations	
Relative Radiation Level	Effective Dose Estimate Range
None	0
Minimal	< 0.1 mSv
Low	0.1-1 mSv
Medium	1-10 mSv
High	10-100 mSv

References

1. Wilcox CM, Alexander LN, Clark WS. Localization of an obstructing esophageal lesion. Is the patient accurate? *Dig Dis Sci* 1995; 40(10):2192-2196.
2. Logemann JA. Role of the modified barium swallow in management of patients with dysphagia. *Otolaryngol Head Neck Surg* 1997; 116(3):335-338.
3. Pikus L, Levine MS, Yang YX, et al. Videofluoroscopic studies of swallowing dysfunction and the relative risk of pneumonia. *AJR* 2003; 180(6):1613-1616.
4. Levine MS, Rubesin SE. Radiologic investigation of dysphagia. *AJR* 1990; 154(6):1157-1163.
5. Rubesin SE. Oral and pharyngeal dysphagia. *Gastroenterol Clin North Am* 1995; 24(2):331-352.
6. Schima W, Pokieser P, Schober E, et al. Globus sensation: value of static radiography combined with videofluoroscopy of the pharynx and oesophagus. *Clin Radiol* 1996; 51(3):177-185.
7. Levine MS, Chu P, Furth EE, Rubesin SE, Laufer I, Herlinger H. Carcinoma of the esophagus and esophagogastric junction: sensitivity of radiographic diagnosis. *AJR* 1997; 168(6):1423-1426.
8. DiPalma JA, Prechter GC, Brady CE, 3rd. X-ray-negative dysphagia: is endoscopy necessary? *J Clin Gastroenterol* 1984; 6(5):409-411.
9. Halpert RD, Feczko PJ, Spickler EM, Ackerman LV. Radiological assessment of dysphagia with endoscopic correlation. *Radiology* 1985; 157(3):599-602.
10. Chen YM, Ott DJ, Gelfand DW, Munitz HA. Multiphasic examination of the esophagogastric region for strictures, rings, and hiatal hernia: evaluation of the individual techniques. *Gastrointest Radiol* 1985; 10(4):311-316.
11. Ott DJ, Chen YM, Wu WC, Gelfand DW, Munitz HA. Radiographic and endoscopic sensitivity in detecting lower esophageal mucosal ring. *AJR* 1986; 147(2):261-265.
12. Ott DJ, Chen YM, Wu WC, Gelfand DW. Endoscopic sensitivity in the detection of esophageal strictures. *J Clin Gastroenterol* 1985; 7(2):121-125.
13. Ott DJ, Gelfand DW, Lane TG, Wu WC. Radiologic detection and spectrum of appearances of peptic esophageal strictures. *J Clin Gastroenterol* 1982; 4(1):11-15.
14. Barloon TJ, Bergus GR, Lu CC. Diagnostic imaging in the evaluation of dysphagia. *Am Fam Physician* 1996; 53(2):535-546.
15. Kahrlas PJ. Clinical approach to dysphagia. In: Gore RM, Levine MS, Laufer I, eds. *Textbook of gastrointestinal radiology*. Philadelphia, Pa: Saunders. 1992:2464-2473.
16. Phillips AJ, Nolan DJ. Radiology of oesophageal dysphagia. *Br J Hosp Med* 1995; 53(9):458-466.
17. Richter JE. Approach to the patient with dysphagia. In: Kelley WN, ed. *Textbook of Internal Medicine*, 3rd ed. Philadelphia, Pa: Lippincott-Raven. 1997:590-593.
18. Ott DJ, Richter JE, Chen YM, Wu WC, Gelfand DW, Castell DO. Esophageal radiography and manometry: correlation in 172 patients with dysphagia. *AJR* 1987; 149(2):307-311.
19. Schima W, Stacher G, Pokieser P, et al. Esophageal motor disorders: videofluoroscopic and manometric evaluation--prospective study in 88 symptomatic patients. *Radiology* 1992; 185(2):487-491.
20. Amaravadi R, Levine MS, Rubesin SE, Laufer I, Redfern RO, Katzka DA. Achalasia with complete relaxation of lower esophageal sphincter: radiographic-manometric correlation. *Radiology* 2005; 235(3):886-891.
21. Maurer AH. Scintigraphic evaluation of the upper gastrointestinal tract. In: Gore RM, Levine MS, Laufer I, eds. *Textbook of gastrointestinal radiology*. Philadelphia, Pa: Saunders. 1992:316-332.
22. Paramsothy M, Goh KL, Kannan P. Oesophageal motility disorders: rapid functional diagnosis using computerised radionuclide oesophageal transit study. *Singapore Med J* 1995; 36(3):309-313.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

23. Stacey B, Patel P. Oesophageal scintigraphy for the investigation of dysphagia: in and out of favour - underused when available. *Eur J Nucl Med Mol Imaging* 2002; 29(9):1216-1220.
24. Bonacini M, Young T, Laine L. The causes of esophageal symptoms in human immunodeficiency virus infection. A prospective study of 110 patients. *Arch Intern Med* 1991; 151(8):1567-1572.
25. Connolly GM, Hawkins D, Harcourt-Webster JN, Parsons PA, Husain OA, Gazzard BG. Oesophageal symptoms, their causes, treatment, and prognosis in patients with the acquired immunodeficiency syndrome. *Gut* 1989; 30(8):1033-1039.
26. Smith PD, Eisner MS, Manischewitz JF, Gill VJ, Masur H, Fox CF. Esophageal disease in AIDS is associated with pathologic processes rather than mucosal human immunodeficiency virus type 1. *J Infect Dis* 1993; 167(3):547-552.
27. Wilcox CM, Schwartz DA, Clark WS. Esophageal ulceration in human immunodeficiency virus infection. Causes, response to therapy, and long-term outcome. *Ann Intern Med* 1995; 123(2):143-149.
28. Rabeneck L, Laine L. Esophageal candidiasis in patients infected with the human immunodeficiency virus. A decision analysis to assess cost-effectiveness of alternative management strategies. *Arch Intern Med* 1994; 154(23):2705-2710.
29. Wilcox CM, Alexander LN, Clark WS, Thompson SE, 3rd. Fluconazole compared with endoscopy for human immunodeficiency virus-infected patients with esophageal symptoms. *Gastroenterology* 1996; 110(6):1803-1809.
30. Balthazar EJ, Megibow AJ, Hulnick D, Cho KC, Beranbaum E. Cytomegalovirus esophagitis in AIDS: radiographic features in 16 patients. *AJR* 1987; 149(5):919-923.
31. Levine MS, Loevner LA, Saul SH, Rubesin SE, Herlinger H, Laufer I. Herpes esophagitis: sensitivity of double-contrast esophagography. *AJR* 1988; 151(1):57-62.
32. Levine MS, Macones AJ, Jr., Laufer I. Candida esophagitis: accuracy of radiographic diagnosis. *Radiology* 1985; 154(3):581-587.
33. Levine MS, Woldenberg R, Herlinger H, Laufer I. Opportunistic esophagitis in AIDS: radiographic diagnosis. *Radiology* 1987; 165(3):815-820.
34. Sor S, Levine MS, Kowalski TE, Laufer I, Rubesin SE, Herlinger H. Giant ulcers of the esophagus in patients with human immunodeficiency virus: clinical, radiographic, and pathologic findings. *Radiology* 1995; 194(2):447-451.
35. Vahey TN, Maglinte DD, Chernish SM. State-of-the-art barium examination in opportunistic esophagitis. *Dig Dis Sci* 1986; 31(11):1192-1195.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.