

**American College of Radiology
ACR Appropriateness Criteria®**

Clinical Condition: **Chronic Elbow Pain**

Variant 1: **Initial evaluation for chronic elbow pain. First test.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
X-ray elbow	9		Min
MRI elbow without contrast	1		None
MR arthrography elbow	1		None
CT elbow without contrast	1		Min
CT arthrography elbow	1		Min
US elbow	1		None
Tc-99m bone scan elbow	1		Med
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 2: **Suspect intra-articular osteocartilaginous body; radiographs nondiagnostic.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
MRI elbow without contrast	9	Either routine MRI or MR arthrogram is appropriate. Depends on availability, expertise, and local conditions. If effusion is present, without contrast is preferred.	None
MR arthrography elbow	9	Either routine MRI or MR arthrogram is appropriate. Depends on availability, expertise, and local conditions. See statement regarding contrast in text under "Anticipated Exceptions."	None
CT elbow without contrast	8		Min
CT arthrography elbow	8	If double contrast is used, dose of less than 0.5 cc of contrast should be used.	Min
US elbow	6	With appropriate expertise.	None
Tc-99m bone scan elbow	1		Med
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Clinical Condition:**Chronic Elbow Pain****Variant 3:****Suspect occult injury; eg, osteochondral injury; radiographs nondiagnostic.**

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without contrast	9		None
CT elbow without contrast	2		Min
MR arthrography elbow	2		None
CT arthrography elbow	2		Min
Tc-99m bone scan elbow	2		Med
US elbow	1		None
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 4:**Suspect unstable osteochondral injury; radiographs nondiagnostic.**

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without contrast	9	Either routine MRI or MR arthrogram is appropriate. Depends on availability, expertise, and local conditions.	None
MR arthrography elbow	9	Either routine MRI or MR arthrogram is appropriate. Depends on availability, expertise, and local conditions. See statement regarding contrast in text under “Anticipated Exceptions.”	None
CT arthrography elbow	8	If MR is contraindicated or not available.	Min
CT elbow without contrast	2		Min
US elbow	1		None
Tc-99m bone scan elbow	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 5:**Suspect mass; radiographs nondiagnostic.**

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow with contrast	9	See statement regarding contrast in text under “Anticipated Exceptions.”	None
US elbow	5	An alternative to MRI if expertise is available.	None
Tc-99m bone scan elbow	2		Med
CT elbow without and with contrast	2		Min
CT arthrography elbow	1		Min
MR arthrography elbow	1		None
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Clinical Condition:**Chronic Elbow Pain****Variant 6:****Suspect chronic epicondylitis; radiographs nondiagnostic.**

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without contrast	9		None
US elbow	8	An alternative to MRI if expertise is available.	None
MR arthrography elbow	2		None
CT elbow without contrast	1		Min
CT arthrography elbow	1		Min
Tc-99m bone scan elbow	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 7:**Suspect collateral ligament tear; radiographs nondiagnostic.**

Radiologic Procedure	Rating	Comments	RRL*
MR arthrography elbow	9	Either routine MRI or MR arthrogram is appropriate. Depends on availability, expertise, and local conditions. See statement regarding contrast in text under “Anticipated Exceptions.”	None
MRI elbow without contrast	9	Either routine MRI or MR arthrogram is appropriate. Depends on availability, expertise, and local conditions.	None
US elbow	6	An alternative to MRI if expertise is available.	None
CT arthrography elbow	5		Min
CT elbow without contrast	2		Min
Tc-99m bone scan elbow	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 8:**Suspect biceps tendon tear and/or bursitis; radiographs nondiagnostic.**

Radiologic Procedure	Rating	Comments	RRL*
MRI elbow without contrast	9		None
US elbow	8	An alternative to MRI if expertise is available.	None
MR arthrography elbow	1		None
CT elbow without contrast	1		Min
CT arthrography elbow	1		Min
Tc-99m bone scan elbow	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Clinical Condition:**Chronic Elbow Pain****Variant 9:****Suspect nerve abnormality; radiographs nondiagnostic.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
MRI elbow without contrast	9		None
US elbow	8	An alternative to MRI if expertise is available. Dynamic US is ideal for assessing ulnar nerve dislocation and snapping triceps syndrome.	None
MR arthrography elbow	1		None
CT elbow without contrast	1		Min
CT arthrography elbow	1		Min
Tc-99m bone scan elbow	1		Med
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

CHRONIC ELBOW PAIN

Expert Panel on Musculoskeletal Imaging: Jon A. Jacobson, MD¹; Richard H. Daffner, MD²; Barbara N. Weissman, MD³; D. Lee Bennett, MD⁴; Judy S. Blebea, MD⁵; William B. Morrison, MD⁶; Charles S. Resnik, MD⁷; Catherine C. Roberts, MD⁸; David A. Rubin, MD⁹; Mark E. Schweitzer, MD¹⁰; Leanne L. Seeger, MD¹¹; Mihra S. Taljanovic, MD¹²; James N. Wise, MD¹³; Robert H. Haralson III, MD¹⁴; William K. Payne III, MD, MPH.¹⁵

Summary of Literature Review

Chronic elbow pain may be caused by a variety of osseous abnormalities, soft-tissue abnormalities, or both. Exclusion of an osseous abnormality with radiographs may be helpful when conservative therapy is planned. In some cases, radiographs may reveal the cause of the problem (eg, intra-articular osteocartilaginous body, calcification around the joint in the form of hydroxyapatite deposition or calcium pyrophosphate crystal deposition). When the etiology of the chronic pain is uncertain and the patient has failed appropriate conservative therapeutic trials, (eg, anti-inflammatory medication, physical therapy, and/or corticosteroid injection), other imaging studies may be considered. While imaging modalities such as computed tomography (CT) and ultrasound (US) may be used for specific indications, magnetic resonance imaging (MRI) can be used to display most abnormalities in the elbow. The success of US varies depending on the training and experience of the person performing the examination, as well as the ultrasound equipment. Imaging choices will be considered for a variety of clinical conditions.

Osteochondral Lesion or Intra-articular Body

Radiographs are required before other imaging studies and may be diagnostic for osteochondral fracture, osteochondritis dissecans, and osteocartilaginous intra-articular body (IAB). CT, as well as CT arthrography with single-contrast (iodinated contrast or air) and double-contrast (iodinated contrast and air) techniques have been

used for detecting an osteochondral lesion or IAB [1]. All of these studies have limitations; a small IAB may be obscured by contrast or confused with air bubbles (double-contrast arthrography). A CT air arthrogram can avoid confusion of air bubbles with IABs. MRI has been advocated as the initial study for suspected osteochondral fracture or IAB [2-4]. Regardless of method, detection of an IAB is limited by its size and location within the elbow joint, although detection is enhanced by the presence of joint effusion [5]. Both CT and MRI can assess for osteochondral fragment stability [6]. MRI following direct intra-articular contrast administration is preferred to routine MRI for diagnosing IAB and may also play a role in improving diagnosis of stability of an osteochondral lesion [7,8]. While US may show osteochondral abnormalities in some situations [9], MRI offers a more comprehensive evaluation of them.

Other Osseous Abnormalities

There are a number of other osseous abnormalities about the elbow that may cause chronic elbow pain; initial evaluation should begin with radiography. Both traumatic and stress fractures may be identified with MRI and bone scan [10]. Osteoid osteomas may produce synovitis if intra-articular and can be identified with MRI, CT, or bone scan [11]. Primary bone tumors are characterized with radiography and MRI before and after intravenous gadolinium administration. While the extent of osseous metastatic disease is assessed with bone scan, MRI will evaluate local extent.

Tendon, Ligament, Muscle, Nerve, or Other Soft-Tissue Abnormality

MRI may provide important diagnostic information for evaluating the elbow in many different conditions, including collateral ligament injury, epicondylitis, injury to the biceps and triceps tendons, and abnormality of the ulnar, radial, or median nerve, and for evaluating masses about the elbow joint [2-5,12-28]. There is a lack of studies showing the sensitivity and specificity of MRI in many of these conditions; most of the studies demonstrate MRI findings in patients either known or highly likely to have a specific condition. US has been shown to be helpful for diagnosing abnormalities of the distal biceps tendon, flexor and extensor tendons, and ligaments, providing an alternative to MRI [29-34].

Radiographs can be useful to identify heterotopic calcification (ossification) of the ulnar collateral ligament [35]. This finding may be associated with partial or complete tears of that structure. Avulsion of the ulnar collateral ligament at the insertion site on the ulna is a source of chronic medial elbow pain in the throwing athlete. While US has been shown to detect medial epicondylar fragmentation of the humerus in throwing athletes [9], this finding is optimally evaluated with a combination of radiographs and coronal MRI [36]. MR arthrography has been advocated to distinguish complete tears from partial tears of the medial collateral ligament

¹Principal Author, University of Michigan Medical Center, Ann Arbor, Michigan.

²Panel Chair, Allegheny General Hospital, Pittsburgh, Pennsylvania.

³Panel Vice-chair, Brigham & Women's Hospital, Boston, Massachusetts.

⁴University of Iowa Health Center, Iowa City, Iowa.

⁵University of Pennsylvania Hospital, Philadelphia, Pennsylvania.

⁶Thomas Jefferson University Hospital, Philadelphia, Pennsylvania.

⁷University of Maryland School of Medicine, Baltimore, Maryland.

⁸Mayo Clinic, Phoenix, Arizona.

⁹Mallinckrodt Institute of Radiology, Saint Louis, Missouri.

¹⁰Hospital for Joint Diseases, New York, New York.

¹¹David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California.

¹²University of Arizona Health Sciences Center, Tucson, Arizona.

¹³University of Arkansas for Medical Sciences, Little Rock, Arkansas.

¹⁴American Academy of Orthopaedic Surgeons, Chicago, Illinois.

¹⁵American Academy of Orthopaedic Surgeons, Chicago, Illinois.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply society endorsement of the final document.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

[22,37], and it improves detection of ulnar collateral ligament tears [28].

With use of appropriate pulse sequences, MRI is an effective tool in the preoperative diagnosis of posterolateral rotatory instability. This includes assessment of the ulnar band of the lateral collateral ligament [25]. Epicondylitis — caused by tendon degeneration and tear of the common extensor tendon laterally (“tennis elbow”) or common flexor tendon medially (in pitchers, golfers, and tennis players) — is a common clinical diagnosis, and imaging is usually not necessary [13]. MRI or US may be useful for confirming the diagnosis in refractory cases and to exclude associated tendon and ligament tear [5,16,20,24,29].

Bicipitoradial and interosseous bursitis around the distal biceps tendon is a source of elbow pain that can be assessed with MRI or US [34,38]. MRI also demonstrates the effects of the bursa on adjacent structures, including the posterior interosseous and median nerves [38]. MRI effectively characterizes a soft-tissue mass, showing its extent and differentiating between intra-articular mass, lymph node (as in cat scratch disease), pseudomass from tendon tear, and other soft-tissue masses.

The ulnar nerve is particularly vulnerable to trauma from a direct blow in the region of its superficial location in the restricted space of the cubital tunnel. Anatomic variations of the cubital tunnel retinaculum may contribute to ulnar neuropathy [39]. Axial T1-weighted images have been shown to depict the size and shape of the nerve, and axial T2-weighted or STIR images may show increased signal in the presence of neuritis [40], and both are more sensitive than conventional nerve conduction studies [41]. US may also show ulnar nerve enlargement and when added to electrodiagnostic tests, increases sensitivity for the diagnosis of ulnar neuropathy at the elbow from 78%-98% [42]. A snapping of the medial head of the triceps can cause recurrent dislocation of the ulnar nerve. This diagnosis can be confirmed with MRI or CT using axial images with the elbow in flexion and extension [40,43,44]. US is ideal for dynamic assessment of ulnar nerve subluxation and dislocation, as well as for confirmation of snapping triceps syndrome [30,45,46]. Radial nerve and median nerve entrapment syndromes may also be evaluated with MRI [5,13,26,40].

Chronic elbow pain may also be caused by a number of joint-related processes, such as inflammatory arthritis (and chronic infection), as well as other synovial proliferative disorders. Evaluation begins with radiography to assess for joint distention and erosions. MRI can also show erosions, and is effective in characterizing synovitis (low signal suggests hemosiderin) and the extent and activity of disease [47]. In the setting of rheumatoid arthritis, US can be used to detect joint effusion, synovitis, and erosions [48].

Summary

Initial evaluation of chronic elbow pain should begin with radiography. Chondral and osteochondral abnormalities can be further evaluated with MRI or CT, although the

addition of arthrography is helpful. Radiographically occult bone abnormalities can be detected with MRI. Soft-tissue abnormalities (tendon, ligament, nerve, joint recess) are well-demonstrated with MRI or US. Dynamic assessment with US is effective for diagnosing nerve or muscle subluxation.

Anticipated Exceptions

Nephrogenic systemic fibrosis (NSF) is a disorder with a scleroderma-like presentation and a spectrum of manifestations that can range from limited clinical sequelae to fatality. It appears to be related to both underlying severe renal dysfunction and the administration of gadolinium-based contrast agents. It has occurred primarily in patients on dialysis, rarely in patients with very limited glomerular filtration rate (GFR) (ie, <30 mL/min/1.73m²), and almost never in other patients. There is growing literature regarding NSF. Although some controversy and lack of clarity remain, there is a consensus that it is advisable to avoid all gadolinium-based contrast agents in dialysis-dependent patients unless the possible benefits clearly outweigh the risk, and to limit the type and amount in patients with estimated GFR rates <30 mL/min/1.73m². For more information, please see the [ACR Manual on Contrast Media](#) [49].

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations	
Relative Radiation Level	Effective Dose Estimate Range
None	0
Minimal	< 0.1 mSv
Low	0.1-1 mSv
Medium	1-10 mSv
High	10-100 mSv

Supporting Document(s)

- [ACR Appropriateness Criteria® Overview](#)
- [Evidence Table](#)

References

1. Singson RD, Feldman F, Rosenberg ZS. Elbow joint: assessment with double-contrast CT arthrography. *Radiology* 1986; 160(1):167-173.
2. Ho CP. Sports and occupational injuries of the elbow: MR imaging findings. *AJR* 1995; 164(6):1465-1471.
3. Quinn SF, Haberman JJ, Fitzgerald SW, Traugher PD, Belkin RI, Murray WT. Evaluation of loose bodies in the elbow with MR imaging. *J Magn Reson Imaging* 1994; 4(2):169-172.

4. Sonin AH, Tutton SM, Fitzgerald SW, Peduto AJ. MR imaging of the adult elbow. *Radiographics* 1996; 16(6):1323-1336.
5. Fritz RC, Steinbach LS. Magnetic resonance imaging of the musculoskeletal system: Part 3. The elbow. *Clin Orthop Relat Res* 1996; (324):321-339.
6. Ouellette H, Kassarian A, Tetreault P, Palmer W. Imaging of the overhead throwing athlete. *Semin Musculoskelet Radiol* 2005; 9(4):316-333.
7. Grainger AJ, Elliott JM, Campbell RS, Tirman PF, Steinbach LS, Genant HK. Direct MR arthrography: a review of current use. *Clin Radiol* 2000; 55(3):163-176.
8. Steinbach LS, Palmer WE, Schweitzer ME. Special focus session. MR arthrography. *Radiographics* 2002; 22(5):1223-1246.
9. Harada M, Takahara M, Sasaki J, Mura N, Ito T, Ogino T. Using sonography for the early detection of elbow injuries among young baseball players. *AJR* 2006; 187(6):1436-1441.
10. Anderson MW. Imaging of upper extremity stress fractures in the athlete. *Clin Sports Med* 2006; 25(3):489-504, vii.
11. Ebrahim FS, Jacobson JA, Lin J, Housner JA, Hayes CW, Resnick D. Intraarticular osteoid osteoma: sonographic findings in three patients with radiographic, CT, and MR imaging correlation. *AJR* 2001; 177(6):1391-1395.
12. Kijowski R, De Smet AA. Magnetic resonance imaging findings in patients with medial epicondylitis. *Skeletal Radiol* 2005; 34(4):196-202.
13. Beltran J, Rosenberg ZS. Diagnosis of compressive and entrapment neuropathies of the upper extremity: value of MR imaging. *AJR* 1994; 163(3):525-531.
14. Bredella MA, Tirman PF, Fritz RC, Feller JF, Wischer TK, Genant HK. MR imaging findings of lateral ulnar collateral ligament abnormalities in patients with lateral epicondylitis. *AJR* 1999; 173(5):1379-1382.
15. Chung CB, Chew FS, Steinbach L. MR imaging of tendon abnormalities of the elbow. *Magn Reson Imaging Clin N Am* 2004; 12(2):233-245, vi.
16. Coel M, Yamada CY, Ko J. MR imaging of patients with lateral epicondylitis of the elbow (tennis elbow): importance of increased signal of the anconeus muscle. *AJR* 1993; 161(5):1019-1021.
17. Fritz RC, Steinbach LS, Tirman PF, Martinez S. MR imaging of the elbow. An update. *Radiol Clin North Am* 1997; 35(1):117-144.
18. Gaary EA, Potter HG, Altchek DW. Medial elbow pain in the throwing athlete: MR imaging evaluation. *AJR* 1997; 168(3):795-800.
19. Ho CP. MR imaging of tendon injuries in the elbow. *Magn Reson Imaging Clin N Am* 1997; 5(3):529-543.
20. Mirowitz SA, London SL. Ulnar collateral ligament injury in baseball pitchers: MR imaging evaluation. *Radiology* 1992; 185(2):573-576.
21. Murphy BJ. MR imaging of the elbow. *Radiology* 1992; 184(2):525-529.
22. Nakanishi K, Masatomi T, Ochi T, et al. MR arthrography of elbow: evaluation of the ulnar collateral ligament of elbow. *Skeletal Radiol* 1996; 25(7):629-634.
23. Patten RM. Overuse syndromes and injuries involving the elbow: MR imaging findings. *AJR* 1995; 164(5):1205-1211.
24. Potter HG, Hannafin JA, Morwessel RM, DiCarlo EF, O'Brien SJ, Altchek DW. Lateral epicondylitis: correlation of MR imaging, surgical, and histopathologic findings. *Radiology* 1995; 196(1):43-46.
25. Potter HG, Weiland AJ, Schatz JA, Paletta GA, Hotchkiss RN. Posterolateral rotatory instability of the elbow: usefulness of MR imaging in diagnosis. *Radiology* 1997; 204(1):185-189.
26. Rosenberg ZS, Beltran J, Cheung YY, Ro SY, Green SM, Lenzo SR. The elbow: MR features of nerve disorders. *Radiology* 1993; 188(1):235-240.
27. Schwartz ML, al-Zahrani S, Morwessel RM, Andrews JR. Ulnar collateral ligament injury in the throwing athlete: evaluation with saline-enhanced MR arthrography. *Radiology* 1995; 197(1):297-299.
28. Kijowski R, Tuite M, Sanford M. Magnetic resonance imaging of the elbow. Part II: Abnormalities of the ligaments, tendons, and nerves. *Skeletal Radiol* 2005; 34(1):1-18.
29. Martinoli C, Bianchi S, Giovagnorio F, Pugliese F. Ultrasound of the elbow. *Skeletal Radiol* 2001; 30(11):605-614.
30. Martinoli C, Bianchi S, Zamorani MP, Zunzunegui JL, Derchi LE. Ultrasound of the elbow. *Eur J Ultrasound* 2001; 14(1):21-27.
31. Miller TT, Adler RS. Sonography of tears of the distal biceps tendon. *AJR* 2000; 175(4):1081-1086.
32. Miller TT, Adler RS, Friedman L. Sonography of injury of the ulnar collateral ligament of the elbow-initial experience. *Skeletal Radiol* 2004; 33(7):386-391.
33. Miller TT, Shapiro MA, Schultz E, Kalish PE. Comparison of sonography and MRI for diagnosing epicondylitis. *J Clin Ultrasound* 2002; 30(4):193-202.
34. Sofka CM, Adler RS. Sonography of cubital bursitis. *AJR* 2004; 183(1):51-53.
35. Mulligan SA, Schwartz ML, Broussard MF, Andrews JR. Heterotopic calcification and tears of the ulnar collateral ligament: radiographic and MR imaging findings. *AJR* 2000; 175(4):1099-1102.
36. Glajchen N, Schwartz ML, Andrews JR, Gladstone J. Avulsion fracture of the sublime tubercle of the ulna: a newly recognized injury in the throwing athlete. *AJR* 1998; 170(3):627-628.
37. Steinbach LS, Schwartz M. Elbow arthrography. *Radiol Clin North Am* 1998; 36(4):635-649.
38. Skaf AY, Boutin RD, Dantas RW, et al. Bicipitoradial bursitis: MR imaging findings in eight patients and anatomic data from contrast material opacification of bursae followed by routine radiography and MR imaging in cadavers. *Radiology* 1999; 212(1):111-116.
39. Sofka CM, Potter HG. Imaging of elbow injuries in the child and adult athlete. *Radiol Clin North Am* 2002; 40(2):251-265.
40. Bordalo-Rodrigues M, Rosenberg ZS. MR imaging of entrapment neuropathies at the elbow. *Magn Reson Imaging Clin N Am* 2004; 12(2):247-263, vi.
41. Vucic S, Cordato DJ, Yiannikas C, Schwartz RS, Shnier RC. Utility of magnetic resonance imaging in diagnosing ulnar neuropathy at the elbow. *Clin Neurophysiol* 2006; 117(3):590-595.
42. Beekman R, Van Der Plas JP, Uitdehaag BM, Schellens RL, Visser LH. Clinical, electrodiagnostic, and sonographic studies in ulnar neuropathy at the elbow. *Muscle Nerve* 2004; 30(2):202-208.
43. Spinner RJ, Goldner RD, Fada RA, Sotereanos DG. Snapping of the triceps tendon over the lateral epicondyle. *J Hand Surg [Am]* 1999; 24(2):381-385.
44. Spinner RJ, Hayden FR, Jr., Hipps CT, Goldner RD. Imaging the snapping triceps. *AJR* 1996; 167(6):1550-1551.
45. Park GY, Kim JM, Lee SM. The ultrasonographic and electrodiagnostic findings of ulnar neuropathy at the elbow. *Arch Phys Med Rehabil* 2004; 85(6):1000-1005.
46. Jacobson JA, Jebson PJ, Jeffers AW, Fessell DP, Hayes CW. Ulnar nerve dislocation and snapping triceps syndrome: diagnosis with dynamic sonography--report of three cases. *Radiology* 2001; 220(3):601-605.
47. Jbara M, Patnana M, Kazmi F, Beltran J. MR imaging: Arthropathies and infectious conditions of the elbow, wrist, and hand. *Radiol Clin North Am* 2006; 44(4):625-642, ix.
48. Lerch K, Borisch N, Paetzel C, Grifka J, Hartung W. Proposal for a sonographic classification of target joints in rheumatoid arthritis. *Rheumatol Int* 2005; 25(3):215-219.
49. American College of Radiology. *Manual on Contrast Media*. Available at: http://www.acr.org/SecondaryMainMenuCategories/quality_safety/contrast_manual.aspx.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.