

**American College of Radiology
ACR Appropriateness Criteria®**

Clinical Condition:

Nontraumatic Knee Pain

Variant 1:

Child or adolescent: nonpatellofemoral symptoms. Mandatory minimal initial examination.

Radiologic Procedure	Rating	Comments	RRL*
X-ray knee	9		Min
X-ray hip ipsilateral	1		Med
CT knee without contrast	1		Low
CT arthrography knee	1		Low
MRI knee without contrast	1		None
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 2:

Child or adult: patellofemoral (anterior) symptoms. Mandatory minimal initial examination.

Radiologic Procedure	Rating	Comments	RRL*
X-ray knee	9		Min
X-ray hip ipsilateral	1		Med
CT knee without contrast	1		Min
CT arthrography knee	1		Min
MRI knee without contrast	1		None
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Clinical Condition:**Nontraumatic Knee Pain****Variant 3:****Adult: nontrauma, nontumor, nonlocalized pain. Mandatory minimal initial examination.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray knee	9		Min
X-ray hip ipsilateral	1		Med
CT knee without contrast	1		Min
CT arthrography knee	1		Min
MRI knee without contrast	1		None
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 4:**Child or adolescent: nonpatellofemoral symptoms. Initial knee radiographs nondiagnostic (demonstrate normal findings or a joint effusion).**

Radiologic Procedure	Rating	Comments	RRL*
MRI knee without contrast	9		None
X-ray hip ipsilateral	2	Indicated if there is clinical evidence or concern for hip pathology causing referred pain to the knee.	Med
CT knee without contrast	1		Low
CT arthrography knee	1		Low
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 5:**Child or adult: patellofemoral (anterior) symptoms. Initial knee radiographs nondiagnostic (demonstrate normal findings or a joint effusion).**

Radiologic Procedure	Rating	Comments	RRL*
MRI knee without contrast	9	If additional imaging is necessary and if internal derangement is suspected.	None
X-ray hip ipsilateral	1		Med
CT knee without contrast	1		Min
CT arthrography knee	1		Min
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Clinical Condition:**Nontraumatic Knee Pain****Variant 6:**

Adult: nontrauma, nontumor, nonlocalized pain. Initial knee radiographs nondiagnostic (demonstrate normal findings or a joint effusion).

Radiologic Procedure	Rating	Comments	RRL*
MRI knee without contrast	9	If additional imaging is necessary and if internal derangement is suspected.	None
X-ray hip ipsilateral	1		Med
CT knee without contrast	1		Min
CT arthrography knee	1		Min
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 7:

Child or adolescent: nonpatellofemoral symptoms. Initial knee radiographs demonstrate osteochondral injuries (fracture/osteochondritis dissecans or a loose body).

Radiologic Procedure	Rating	Comments	RRL*
MRI knee without contrast	9		None
MR arthrography knee	6		None
CT arthrography knee	5	If MRI cannot be done.	Low
X-ray hip ipsilateral	1		Med
CT knee without contrast	1		Low
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 8:

Adult: patellofemoral (anterior) symptoms. Initial knee radiographs demonstrate degenerative joint disease and/or chondrocalcinosis.

Radiologic Procedure	Rating	Comments	RRL*
X-ray hip ipsilateral	1		Med
CT knee without contrast	1		Min
CT arthrography knee	1		Min
MRI knee without contrast	1		None
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Clinical Condition:**Nontraumatic Knee Pain****Variant 9:**

Adult: nontumor, nonlocalized pain. Initial knee radiographs demonstrate inflammatory, crystalline, or degenerative joint disease (uni- to tri-compartmental sclerosis, hypertrophic spurs, joint space narrowing, and/or subchondral cysts).

Radiologic Procedure	Rating	Comments	RRL*
X-ray hip ipsilateral	1		Med
CT knee without contrast	1		Min
CT arthrography knee	1		Min
MRI knee without contrast	1	Consider for preoperative assessment.	None
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 10:

Adult: nontumor, nonlocalized pain. Initial knee radiographs demonstrate avascular necrosis.

Radiologic Procedure	Rating	Comments	RRL*
MRI knee without contrast	7	If needed for therapy.	None
CT knee without contrast	1		Min
CT arthrography knee	1		Min
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 11:

Adult: nontumor, nonlocalized pain. Initial knee radiographs demonstrate evidence of internal derangement (eg, Segond fracture, deep lateral femoral notch sign).

Radiologic Procedure	Rating	Comments	RRL*
MRI knee without contrast	9		None
CT arthrography knee	2	If MRI contraindicated.	Min
CT knee without contrast	1		Min
MR arthrography knee	1		None
US knee	1		None
Tc-99m bone scan lower extremity	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

NONTRAUMATIC KNEE PAIN

Expert Panel on Musculoskeletal Imaging:
D. Lee Bennett, MD, MA¹; Richard H. Daffner, MD²;
Barbara N. Weissman, MD³; Judy S. Blebea, MD⁴;
Jon A. Jacobson, MD⁵; William B. Morrison, MD⁶;
Charles S. Resnik, MD⁷; Catherine C. Roberts, MD⁸;
David A. Rubin, MD⁹; Mark E. Schweitzer, MD¹⁰;
Leanne L. Seeger, MD¹¹; Mihra S. Taljanovic, MD¹²;
James N. Wise, MD¹³; William K. Payne III, MD, MPH.¹⁴

Summary of Literature Review

Nontraumatic knee pain in children, adolescents, and adults includes localized complaints such as anterior (patellofemoral) pain and diffuse nonlocalized symptoms. The consensus of the committee is that the initial imaging study for nontraumatic knee pain are anteroposterior (AP) and lateral radiographs [1-3]. For patients with diffuse nonlocalized symptoms, a Merchant or axial view may be useful as part of the initial examination. In children with nontraumatic knee pain, referred pain from the hip must be considered and hip radiographs may need to be obtained if there is clinical evidence or clinical concern for hip pathology.

In elderly patients, the most common source of nontraumatic knee pain is osteoarthritis. Conventional radiographic diagnosis of degenerative joint disease (osteoarthritis) includes joint space narrowing, osteophytes, subchondral cysts, and sclerosis bordering the joint. Articular cartilage is evaluated indirectly on radiographs by joint space narrowing and changes in the subchondral bone [4]. Routine radiographs are insensitive for assessing articular cartilage in the early stages of osteoarthritis, while in advanced disease, joint space narrowing on radiographs is usually an accurate assessment of cartilage loss [5,6]. Standing radiographs have been reported to more accurately reflect medial and lateral joint compartment cartilage loss than supine radiographs; however, in the presence of a severe varus or valgus deformity, significant cartilage loss in the compartment that appears wide (due to the alignment deformity) may not be evident [7-9]. A weight-bearing

posteroanterior (PA) radiograph, obtained with knee flexion, has been reported to show the cartilage width of the posterior medial and lateral joint compartments more accurately than a standing view obtained with the knee extended [10]. The standing flexed view may be indicated in elderly patients with osteoarthritis when surgical intervention is being planned. Finally, one should bear in mind that a significant portion of the joint space narrowing may be due to meniscal extrusion or degeneration rather than hyaline cartilage loss in some patients [11]. Additional imaging studies are not indicated in patients for whom radiographs are diagnostic of degenerative joint disease unless treatment or surgery or both depend on additional findings such as internal knee derangement, or when symptoms are not explained by the radiographic findings (eg, stress fractures).

Other nontraumatic causes of knee pain in adult patients include internal knee derangement (meniscal and ligament tears), stress fracture, subchondral insufficiency fracture (known as spontaneous osteonecrosis), inflammatory arthritis, transient osteoporosis, and chronic regional pain syndrome. Chronic anterolateral knee pain may also result from patellar tendon–lateral femoral condyle friction syndrome or iliotibial band syndrome (friction syndrome), both of which can be confirmed or excluded by magnetic resonance imaging (MRI) [12].

When initial radiographs are nondiagnostic (normal findings or a joint effusion) and knee symptoms are suspicious for an internal derangement, the next indicated study is an MRI examination [13]. MRI is also indicated when the patient has persistent knee pain and normal radiographs. MRI is more sensitive than radiography and provides more specific information compared with radionuclide bone scan [14,15]. MRI of nontraumatic knee pain may document a joint effusion, communicating synovial cysts, proliferative changes of the synovial membrane, osteophytes, subchondral cysts, articular cartilage loss, meniscal and/or ligamentous tears and/or degeneration, bone marrow edema, fractures, and osteonecrosis [14-16]. A secondary MRI finding with a high sensitivity for internal derangement is an AP joint fluid measurement of greater than 10 mm in the lateral suprapatellar pouch [17].

MRI is useful to identify a subchondral insufficiency fracture as the initial injury from which localized osteonecrosis may result and which was otherwise identified as spontaneous osteonecrosis [18]. MRI can also detect osteonecrosis of the medial femoral condyle or of the medial tibial plateau associated with tibial stress fracture [19].

A suprapatellar joint effusion is readily detected on a lateral radiograph of the knee; however, the extent of a joint effusion, the presence of a communicating synovial (popliteal) cyst, or synovial proliferation is readily identified on MRI [16,18-24]. Subchondral cysts are easily detected on MRI because of its tomographic

¹Principal Author, University of Iowa Health Center, Iowa City, Iowa.

²Panel Chair, Allegheny General Hospital, Pittsburgh, Pennsylvania.

³Panel Vice-chair, Brigham & Women's Hospital, Boston, Massachusetts.

⁴Cleveland Clinic, Cleveland, Ohio.

⁵University of Michigan Medical Center, Ann Arbor, Michigan.

⁶Thomas Jefferson University Hospital, Philadelphia, Pennsylvania.

⁷University of Maryland School of Medicine, Baltimore, Maryland.

⁸Mayo Clinic, Phoenix, Arizona.

⁹Mallinckrodt Institute of Radiology, Saint Louis, Missouri.

¹⁰University of Ottawa, Ottawa, Ontario, Canada.

¹¹David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California.

¹²University of Arizona Health Sciences Center, Tucson, Arizona.

¹³University of Kentucky, Lexington, Kentucky.

¹⁴American Academy of Orthopaedic Surgeons, Chicago, Illinois.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply society endorsement of the final document.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

quality, multiplaner imaging capability, and superb sensitivity to fluid- and fat-containing tissues [15,16,25]. Cartilage pathology, both articular and meniscal, can be evaluated directly on MRI, and demonstration depends on the location of the abnormality and the pulse sequences used [26-30].

Magnetic resonance arthrography (MRA) performed with an intra-articular injection of dilute gadolinium solution [31,32] or with an intravenous injection [33] of gadolinium contrast to improve cartilage evaluation has been investigated, but noncontrast MRI has been reported accurate for cartilage abnormalities [26]. Patellofemoral cartilage loss has been reported to be closely associated with chronic knee pain symptoms [34].

Transient osteoporosis is characterized by self-limited pain and radiographically demonstrable osteopenia. The osteopenia typically develops within eight weeks after the onset of pain. Spontaneous osteonecrosis of the medial femoral condyle, most often found in middle-aged and elderly females, may have normal radiographs for months, followed by subchondral collapse, fragmentation of the articular cartilage, and progressive osteoarthritis [35-37]. Bone marrow edema seen on MRI occurs in association with, or independent of, transient osteoporosis or osteonecrosis, and also in association with stress fractures; MRI is highly sensitive for detecting these abnormalities [36]. In adult patients with conventional radiograph diagnosis of an osteochondral injury such as osteochondritis dissecans or osteonecrosis, an MRI examination may be indicated if an additional injury is suspected clinically or when it is necessary to determine the status of the articular cartilage over the area of abnormality. In the child or adolescent with radiographic evidence of osteochondritis dissecans, an MRI is indicated to determine the best method of treatment [38-40]. Finally, MRI is not indicated to confirm a stress fracture that is evident on the radiographic study.

In patients with radiographic evidence of inflammatory arthritis of the knee, the consensus of the panel is that a knee MRI is usually not indicated for preoperative differentiation of pannus from effusion or for evaluation of erosion [24]. An aspiration for crystals may be indicated; however, the use of medical imaging (such as fluoroscopic guidance, ultrasound guidance, or arthrographic confirmation) is usually not necessary.

When an intra-articular abnormality is suspected in a patient with claustrophobia, with a large body habitus, or who cannot for some reason tolerate an MRI examination, or when there is contraindication to an MRI, a CT arthrogram may be used instead of the MRI to evaluate the cruciate ligaments, menisci, and articular cartilage [41,42].

Summary

- The mandatory initial imaging examination for nontraumatic knee pain is AP and lateral radiography.

- In patients with anterior patellofemoral knee pain, an axial view should be included in the initial radiographic study.
- An MRI examination for nontraumatic knee pain is indicated when the pain is persistent and conventional radiographs are nondiagnostic or when additional information is necessary before instituting treatment or surgical intervention.
- An MRI is not indicated before a physical examination or routine conventional radiographs, or when there is diagnostic radiographic evidence of severe degenerative joint diseases, inflammatory arthritis, stress fracture, osteonecrosis, or reflex sympathetic dystrophy, for which additional imaging is not going to alter the treatment plan.

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations	
Relative Radiation Level	Effective Dose Estimate Range
None	0
Minimal	< 0.1 mSv
Low	0.1-1 mSv
Medium	1-10 mSv
High	10-100 mSv

Supporting Document(s)

- [ACR Appropriateness Criteria® Overview](#)
- [Evidence Table](#)

References

1. Laurin CA, Dussault R, Levesque HP. The tangential x-ray investigation of the patellofemoral joint: x-ray technique, diagnostic criteria and their interpretation. *Clin Orthop Relat Res* 1979; (144):16-26.
2. Merchant AC, Mercer RL, Jacobsen RH, Cool CR. Roentgenographic analysis of patellofemoral congruence. *J Bone Joint Surg Am* 1974; 56(7):1391-1396.
3. Newberg AH, Seligson D. The patellofemoral joint: 30 degrees, 60 degrees, and 90 degrees views. *Radiology* 1980; 137(1 Pt 1):57-61.
4. Hayes CW, Conway WF. Evaluation of articular cartilage: radiographic and cross-sectional imaging techniques. *Radiographics* 1992; 12(3):409-428.
5. Brandt KD, Fife RS, Braunstein EM, Katz B. Radiographic grading of the severity of knee osteoarthritis: relation of the Kellgren and Lawrence grade to a grade based on joint space narrowing, and correlation with arthroscopic evidence of articular cartilage degeneration. *Arthritis Rheum* 1991; 34(11):1381-1386.

6. Messieh SS, Fowler PJ, Munro T. Anteroposterior radiographs of the osteoarthritic knee. *J Bone Joint Surg Br* 1990; 72(4):639-640.
7. Altman RD, Fries JF, Bloch DA, et al. Radiographic assessment of progression in osteoarthritis. *Arthritis Rheum* 1987; 30(11):1214-1225.
8. Leach RE, Gregg T, Siber FJ. Weight-bearing radiography in osteoarthritis of the knee. *Radiology* 1970; 97(2):265-268.
9. Marklund T, Myrneris R. Radiographic determination of cartilage height in the knee joint. *Acta Orthop Scand* 1974; 45(5):752-755.
10. Rosenberg TD, Paulos LE, Parker RD, Coward DB, Scott SM. The forty-five-degree posteroanterior flexion weight-bearing radiograph of the knee. *J Bone Joint Surg Am* 1988; 70(10):1479-1483.
11. Hunter DJ, Zhang YQ, Tu X, et al. Change in joint space width: hyaline articular cartilage loss or alteration in meniscus? *Arthritis Rheum* 2006; 54(8):2488-2495.
12. Chung CB, Skaf A, Roger B, Campos J, Stump X, Resnick D. Patellar tendon-lateral femoral condyle friction syndrome: MR imaging in 42 patients. *Skeletal Radiol* 2001; 30(12):694-697.
13. Vincken PW, ter Braak AP, van Erkel AR, et al. MR imaging: effectiveness and costs at triage of patients with nonacute knee symptoms. *Radiology* 2007; 242(1):85-93.
14. McAlindon TE, Watt I, McCrae F, Goddard P, Dieppe PA. Magnetic resonance imaging in osteoarthritis of the knee: correlation with radiographic and scintigraphic findings. *Ann Rheum Dis* 1991; 50(1):14-19.
15. Reiser MF, Vahlensieck M, Schuller H. Imaging of the knee joint with emphasis on magnetic resonance imaging. *Eur Radiol* 1992; 2:87-94.
16. Sabiston CP, Adams ME, Li DK. Magnetic resonance imaging of osteoarthritis: correlation with gross pathology using an experimental model. *J Orthop Res* 1987; 5(2):164-172.
17. Kolman BH, Daffner RH, Sciulli RL, Soehnen MW. Correlation of joint fluid and internal derangement on knee MRI. *Skeletal Radiol* 2004; 33(2):91-95.
18. Yamamoto T, Bullough PG. Spontaneous osteonecrosis of the knee: the result of subchondral insufficiency fracture. *J Bone Joint Surg Am* 2000; 82(6):858-866.
19. Le Gars L, Savy JM, Orcel P, et al. Osteonecrosis-like syndrome of the medial tibial plateau can be due to a stress fracture. MR findings in 13 patients. *Rev Rhum Engl Ed* 1999; 66(6):323-330.
20. Beltran J, Caudill JL, Herman LA, et al. Rheumatoid arthritis: MR imaging manifestations. *Radiology* 1987; 165(1):153-157.
21. Reiser MF, Naegele M. Inflammatory joint disease: static and dynamic gadolinium-enhanced MR imaging. *J Magn Reson Imaging* 1993; 3(1):307-310.
22. Shanley DJ, Auber AE, Watabe JT, Buckner AB. Pigmented villonodular synovitis of the knee demonstrated on bone scan. Correlation with US, CT, and MRI. *Clin Nucl Med* 1992; 17(11):901-902.
23. Terrier F, Hricak H, Revel D, et al. Magnetic resonance imaging and spectroscopy of the periarticular inflammatory soft-tissue changes in experimental arthritis of the rat. *Invest Radiol* 1985; 20(8):813-823.
24. Yulish BS, Montanez J, Goodfellow DB, Bryan PJ, Mulopulos GP, Modic MT. Chondromalacia patellae: assessment with MR imaging. *Radiology* 1987; 164(3):763-766.
25. Chan WP, Lang P, Stevens MP, et al. Osteoarthritis of the knee: comparison of radiography, CT, and MR imaging to assess extent and severity. *AJR* 1991; 157(4):799-806.
26. Boegard TL, Rudling O, Petersson IF, Jonsson K. Magnetic resonance imaging of the knee in chronic knee pain. A 2-year follow-up. *Osteoarthritis Cartilage* 2001; 9(5):473-480.
27. Ghelman B, Hodge JC. Imaging of the patellofemoral joint. *Orthop Clin North Am* 1992; 23(4):523-543.
28. Konig H, Sauter R, Deimling M, Vogt M. Cartilage disorders: comparison of spin-echo, CHES, and FLASH sequence MR images. *Radiology* 1987; 164(3):753-758.
29. Reiser MF, Bongartz G, Erlemann R, et al. Magnetic resonance in cartilaginous lesions of the knee joint with three-dimensional gradient-echo imaging. *Skeletal Radiol* 1988; 17(7):465-471.
30. Spritzer CE, Vogler JB, Martinez S, et al. MR imaging of the knee: preliminary results with a 3DFT GRASS pulse sequence. *AJR* 1988; 150(3):597-603.
31. Engel A. Magnetic resonance knee arthrography. Enhanced contrast by gadolinium complex in the rabbit and in humans. *Acta Orthop Scand Suppl* 1990; 240:1-57.
32. Hajek PC, Baker LL, Sartoris DJ, Neumann CH, Resnick D. MR arthrography: anatomic-pathologic investigation. *Radiology* 1987; 163(1):141-147.
33. Winalski CS, Aliabadi P, Wright RJ, Shortkroff S, Sledge CB, Weissman BN. Enhancement of joint fluid with intravenously administered gadopentetate dimeglumine: technique, rationale, and implications. *Radiology* 1993; 187(1):179-185.
34. Hunter DJ, March L, Sambrook PN. The association of cartilage volume with knee pain. *Osteoarthritis Cartilage* 2003; 11(10):725-729.
35. Ahlback S, Bauer GC, Bohne WH. Spontaneous osteonecrosis of the knee. *Arthritis Rheum* 1968; 11(6):705-733.
36. Hayes CW, Conway WF, Daniel WW. MR imaging of bone marrow edema pattern: transient osteoporosis, transient bone marrow edema syndrome, or osteonecrosis. *Radiographics* 1993; 13(5):1001-1011; discussion 1012.
37. Lotke PA, Ecker ML. Osteonecrosis of the knee. *J Bone Joint Surg Am* 1988; 70(3):470-473.
38. De Smet AA, Ilahi OA, Graf BK. Untreated osteochondritis dissecans of the femoral condyles: prediction of patient outcome using radiographic and MR findings. *Skeletal Radiol* 1997; 26(8):463-467.
39. O'Connor MA, Palaniappan M, Khan N, Bruce CE. Osteochondritis dissecans of the knee in children. A comparison of MRI and arthroscopic findings. *J Bone Joint Surg Br* 2002; 84(2):258-262.
40. Pill SG, Ganley TJ, Milam RA, Lou JE, Meyer JS, Flynn JM. Role of magnetic resonance imaging and clinical criteria in predicting successful nonoperative treatment of osteochondritis dissecans in children. *J Pediatr Orthop* 2003; 23(1):102-108.
41. Vande Berg BC, Lecouvet FE, Maldague B, Malghem J. MR appearance of cartilage defects of the knee: preliminary results of a spiral CT arthrography-guided analysis. *Eur Radiol* 2004; 14(2):208-214.
42. Vande Berg BC, Lecouvet FE, Poilvache P, Dubuc JE, Maldague B, Malghem J. Anterior cruciate ligament tears and associated meniscal lesions: assessment at dual-detector spiral CT arthrography. *Radiology* 2002; 223(2):403-409.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.