

**American College of Radiology
ACR Appropriateness Criteria®**

Clinical Condition: Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus

Variant 1: Soft-tissue edema without ulcer or neuroarthropathy.

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
X-ray foot	9	Initial study. Radiographs and MRI are complementary. Both are indicated.	⊕
MRI foot with contrast	9	Radiographs and MRI are complementary. Both are indicated. Useful for mapping devitalized areas preoperatively. See statement regarding contrast in text under "Anticipated Exceptions."	O
MRI foot without contrast	9	Radiographs and MRI are complementary. Both are indicated.	O
Tc-99m 3-phase bone scan and In-111 WBC scan foot	4	If MRI contraindicated.	⊕ ⊕ ⊕ ⊕
Tc-99m 3-phase bone scan foot	1		⊕ ⊕ ⊕
In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		⊕ ⊕ ⊕ ⊕
Tc-99m 3-phase bone scan and In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		⊕ ⊕ ⊕ ⊕
US foot	1		O
CT foot without contrast	1		⊕
FDG-PET foot	1		⊕ ⊕ ⊕ ⊕
<u>Rating Scale:</u> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition:

Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus

Variant 2:

Ulcer with no exposed bone without neuroarthropathy.

Radiologic Procedure	Rating	Comments	RRL*
X-ray foot	9	Initial study. Radiographs and MRI are complementary. Both are indicated.	⊕
MRI foot with contrast	9	Radiographs and MRI are complementary. Both are indicated. Useful for mapping devitalized areas preoperatively. See statement regarding contrast in text under “Anticipated Exceptions.”	O
MRI foot without contrast	9	Radiographs and MRI are complementary. Both are indicated.	O
Tc-99m 3-phase bone scan and In-111 WBC scan foot	4	If MRI contraindicated.	⊕ ⊕ ⊕ ⊕
Tc-99m 3-phase bone scan foot	1		⊕ ⊕ ⊕
In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		⊕ ⊕ ⊕ ⊕
Tc-99m 3-phase bone scan and In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		⊕ ⊕ ⊕ ⊕
US foot	1		O
CT foot without contrast	1		⊕
FDG-PET foot	1		⊕ ⊕ ⊕ ⊕
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 3:

Ulcer with exposed bone without neuroarthropathy.

Radiologic Procedure	Rating	Comments	RRL*
X-ray foot	9	Initial study. Radiographs and MRI are complementary. Both are indicated.	⊕
MRI foot with contrast	9	Radiographs and MRI are complementary. Both are indicated. Useful for mapping devitalized areas preoperatively. See statement regarding contrast in text under “Anticipated Exceptions.”	O
MRI foot without contrast	9	Radiographs and MRI are complementary. Both are indicated.	O
Tc-99m 3-phase bone scan and In-111 WBC scan foot	4	If MRI contraindicated.	⊕ ⊕ ⊕ ⊕
Tc-99m 3-phase bone scan foot	1		⊕ ⊕ ⊕
In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		⊕ ⊕ ⊕ ⊕
Tc-99m 3-phase bone scan and In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		⊕ ⊕ ⊕ ⊕
US foot	1		O
CT foot without contrast	1		⊕
FDG-PET foot	1		⊕ ⊕ ⊕ ⊕
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition:**Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus****Variant 4:****Neuropathy without ulcer.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
X-ray foot	9	Initial study. Radiographs and MRI are complementary. Both are indicated.	☼
MRI foot with contrast	9	Radiographs and MRI are complementary. Both are indicated. See statement regarding contrast in text under "Anticipated Exceptions."	O
MRI foot without contrast	9	Radiographs and MRI are complementary. Both are indicated.	O
CT foot without contrast	5	For neuropathy or if MRI contraindicated.	☼
Tc-99m 3-phase bone scan foot	5	Useful for pre-radiographic findings of neuropathy. Also if MRI contraindicated.	☼ ☼ ☼
Tc-99m 3-phase bone scan and In-111 WBC scan foot	2		☼ ☼ ☼ ☼
In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		☼ ☼ ☼ ☼
Tc-99m 3-phase bone scan and In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		☼ ☼ ☼ ☼
US foot	1		O
FDG-PET foot	1		☼ ☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 5:**Neuroarthropathy with ulcer without exposed bone.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
X-ray foot	9	Initial study. Radiographs and MRI are complementary. Both are indicated.	☼
MRI foot with contrast	9	Radiographs and MRI are complementary. Both are indicated. See statement regarding contrast in text under "Anticipated Exceptions."	O
MRI foot without contrast	9	Radiographs and MRI are complementary. Both are indicated.	O
Tc-99m 3-phase bone scan and In-111 WBC scan foot	4	If MRI contraindicated.	☼ ☼ ☼ ☼
Tc-99m 3-phase bone scan foot	1		☼ ☼ ☼
In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		☼ ☼ ☼ ☼
Tc-99m 3-phase bone scan and In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		☼ ☼ ☼ ☼
CT foot without contrast	1		☼
US foot	1		O
FDG- PET foot	1		☼ ☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition:**Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus****Variant 6:****Neuroarthopathy with ulcer with exposed bone.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
X-ray foot	9	Initial study. Radiographs and MRI are complementary. Both are indicated.	⊕
MRI foot with contrast	9	Radiographs and MRI are complementary. Both are indicated. See statement regarding contrast in text under "Anticipated Exceptions."	O
MRI foot without contrast	9	Radiographs and MRI are complementary. Both are indicated.	O
Tc-99m 3-phase bone scan and In-111 WBC scan foot	4	If MRI contraindicated.	⊕ ⊕ ⊕ ⊕
Tc-99m 3-phase bone scan foot	1		⊕ ⊕ ⊕
In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		⊕ ⊕ ⊕ ⊕
Tc-99m 3-phase bone scan and In-111 WBC scan and Tc-99m sulfur colloid marrow scan foot	1		⊕ ⊕ ⊕ ⊕
CT foot without contrast	1		⊕
US foot	1		O
FDG-PET foot	1		⊕ ⊕ ⊕ ⊕
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

SUSPECTED OSTEOMYELITIS OF THE FOOT IN PATIENTS WITH DIABETES MELLITUS

Expert Panel on Musculoskeletal Imaging:
Mark E. Schweitzer, MD¹; Richard H. Daffner, MD²;
Barbara N. Weissman, MD³; D. Lee Bennett, MD, MA⁴;
Judy S. Blebea, MD⁵; Jon A. Jacobson, MD⁶;
William B. Morrison, MD⁷; Charles S. Resnik, MD⁸;
Catherine C. Roberts, MD⁹; David A. Rubin, MD¹⁰;
Leanne L. Seeger, MD¹¹; Mihra Taljanovic, MD¹²;
James N. Wise, MD¹³; William K. Payne III, MD, MPH.¹⁴

Summary of Literature Review

Through the last 50 years there has been much written about the diabetic foot, with little consensus as to whether, when, and what imaging is appropriate. This overview will summarize some of the work and draw conclusions based on the available data. We will discuss several clinical situations in which osteomyelitis or diabetic pedal disease is suspected but clinical findings differ because of the presence or absence of edema ulceration and neuropathy.

Please note that although several of the variants have similar recommendations, they do present as unique clinical scenarios.

Soft-Tissue Edema without Ulceration

First, the probability of having osteomyelitis in a diabetic foot without evidence of ulceration is extremely low [1]. Whether there is or is not soft-tissue swelling, these patients have almost no incidence of osteomyelitis and a low incidence of septic arthritis, but some frequency of soft-tissue infections [2]. The only situation in which such a patient can have osteomyelitis is the presence of a “hidden” ulcer that has granulated over and may appear healed. In that situation the risk of osteomyelitis is still extremely low, since the ulcer would not have granulated over if osteomyelitis were present [3]. Therefore, without a clinically apparent ulcer, the role of imaging might be to diagnose neuropathic disease or to see if there is soft-tissue infection only [3].

Neuropathy without Ulcer

A more difficult question is whether it is the neuroarthropathy or the soft-tissue infection that is causing the soft-tissue swelling [4,5]. In a patient who has neuroarthropathy, the risk of infection is usually low without ulceration. Radiography can be used as a screening examination. Computed tomography (CT) may pick up neuroarthropathy, which may not be apparent radiographically and may be the cause of the swelling and pain (mimicking infection). CT can rarely exclude the diagnosis of osteomyelitis definitively if there is no edema in the marrow (fat is visible).

Scintigraphy is of indeterminate insensitivity and specificity, whether it is bone scan, indium or indium with sulfur colloid, or even positron emission tomography (PET) [6-9]. Flow images are the best discriminators of infection, but remain imperfect. Magnetic resonance imaging (MRI) likely has the best clinical results in this scenario with or without contrast, but the yield is going to be low in this clinical group of patients, and it is costly [10].

There is some importance in diagnosing neuropathic disease prior to radiographic changes, as these patients will be treated with altered footwear and orthotics to prevent the progression to deformity. Scintigraphy is, however, extremely sensitive to early neuropathic disease, long before radiographic changes are present. MRI is less sensitive but is a better test if there is a possibility of soft-tissue infection.

Ulcer with Exposed Bone

If an ulcer is present, the risk of infection is quite high, and almost invariable if the ulcer reaches bone. The role of imaging would be to confirm the infection and show extent. Radiography will show the infection, however late. Bone scan is quite nonspecific [7,11]. Surprisingly, indium scan, even when combined with sulfur colloid marrow imaging, has low specificity [12-14], although if the ulcer is away from the joint these techniques are better. MRI has high specificity and sensitivity both with and without contrast [15]. Ultrasound (US) may have promise in long bones but, to date, data about its utility in diagnosing the diabetic foot are quite limited. PET results are similarly poor, as this technique shows metabolic activity primarily and therefore is not specific [16].

Ulcer with Neuropathy and Exposed Bone

In patients who have diabetes and secondary neuroarthropathy, the infection is usually over an osseous abnormality with an ulcer. If the ulcer tracks down to bone, the risk of osteomyelitis is extremely high, perhaps even higher than in the preceding situation where there is an ulcer without neuropathic deformity. The overall role of imaging therefore, is more to determine the extent of the disease than to definitively diagnose it [17]. Therefore, most authors do not advocate scintigraphy in

¹Principal Author, University of Ottawa, Ottawa, Ontario, Canada.

²Panel Chair, Allegheny General Hospital, Pittsburgh, Pennsylvania.

³Panel Vice-chair, Brigham & Women's Hospital, Boston, Massachusetts.

⁴University of Iowa Health Center, Iowa City, Iowa.

⁵Cleveland Clinic, Cleveland, Ohio.

⁶University of Michigan Medical Center, Ann Arbor, Michigan.

⁷Thomas Jefferson University Hospital, Philadelphia, Pennsylvania.

⁸University of Maryland School of Medicine, Baltimore, Maryland.

⁹Mayo Clinic, Phoenix, Arizona.

¹⁰Mallinckrodt Institute of Radiology, Saint Louis, Missouri.

¹¹David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California.

¹²University of Arizona Health Sciences Center, Tucson, Arizona.

¹³University of Kentucky, Lexington, Kentucky.

¹⁴American Academy of Orthopaedic Surgeons, Chicago, Illinois.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply society endorsement of the final document.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

this situation because of its relative poor spatial resolution for extent of disease; similar conclusions apply to PET [11].

Similarly, indium-labeled WBC (white blood cell) scanning with or without bone marrow scanning has only mixed sensitivity and specificity for osteomyelitis with neuropathy and yields poor anatomic extent of infection. Radiography has a high specificity but low sensitivity. US is unproven. CT will show the neuroarthropathic disease but not much else. MRI should be performed to determine extent of disease [1]. T1 and fat-suppressed sequences are complementary, and contrast may or may not be used. The use of contrast is more to see the extent of the disease as well as the extent of vascularity, rather than to diagnose infections [10]. Contrast may also help identify necrotic or poorly perfused regions, and to aid in surgical planning [18,19].

Summary and Recommendations

If a patient has an ulcer that extends to bone, there is quite likely, but not invariably, osteomyelitis. The best way to confirm this diagnosis and determine the extent of disease is with MRI. If there is no ulcer and there is still a clinical suspicion of infection, MRI is the test of choice. However, conventional radiographs should be done simultaneously in both situations. In indeterminate cases, aspiration and biopsy would be the next step.

If there is soft-tissue swelling the question is, “Is there early neuropathic disease or infection present?” Radiographs should be performed first. If the radiographs are normal, another test should be performed. If the suspicion of infection is low, the next test should probably be a three-phase bone scan. If there is a modest risk of infection, MRI is probably indicated.

Anticipated Exceptions

Nephrogenic systemic fibrosis (NSF) is a disorder with a scleroderma-like presentation and a spectrum of manifestations that can range from limited clinical sequelae to fatality. It appears to be related to both underlying severe renal dysfunction and the administration of gadolinium-based contrast agents. It has occurred primarily in patients on dialysis, rarely in patients with very limited glomerular filtration rate (GFR) (ie, <30 mL/min/1.73m²), and almost never in other patients. There is growing literature regarding NSF. Although some controversy and lack of clarity remain, there is a consensus that it is advisable to avoid all gadolinium-based contrast agents in dialysis-dependent patients unless the possible benefits clearly outweigh the risk, and to limit the type and amount in patients with estimated GFR rates <30 mL/min/1.73m². For more information, please see the [ACR Manual on Contrast Media](#) [20].

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated

with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, both because of organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared to those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
O	0 mSv	0 mSv
⊕	<0.1 mSv	<0.03 mSv
⊕ ⊕	0.1-1 mSv	0.03-0.3 mSv
⊕ ⊕ ⊕	1-10 mSv	0.3-3 mSv
⊕ ⊕ ⊕ ⊕	10-30 mSv	3-10 mSv
⊕ ⊕ ⊕ ⊕ ⊕	30-100 mSv	10-30 mSv
*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as NS (not specified).		

Supporting Document(s)

- [ACR Appropriateness Criteria® Overview](#)
- [Procedure Contrast Information](#)
- Evidence table under review

References

1. Ledermann HP, Morrison WB, Schweitzer ME. Pedal abscesses in patients suspected of having pedal osteomyelitis: analysis with MR imaging. *Radiology* 2002; 224(3):649-655.
2. Schweitzer ME, Morrison WB. MR imaging of the diabetic foot. *Radiol Clin North Am* 2004; 42(1):61-71, vi.
3. Tomas MB, Patel M, Marwin SE, Palestro CJ. The diabetic foot. *Br J Radiol* 2000; 73(868):443-450.
4. Chatha DS, Cunningham PM, Schweitzer ME. MR imaging of the diabetic foot: diagnostic challenges. *Radiol Clin North Am* 2005; 43(4):747-759, ix.
5. Ledermann HP, Morrison WB. Differential diagnosis of pedal osteomyelitis and diabetic neuroarthropathy: MR Imaging. *Semin Musculoskelet Radiol* 2005; 9(3):272-283.
6. Hopfner S, Krolak C, Kessler S, et al. Preoperative imaging of Charcot neuroarthropathy in diabetic patients: comparison of ring PET, hybrid PET, and magnetic resonance imaging. *Foot Ankle Int* 2004; 25(12):890-895.
7. Jay PR, Michelson JD, Mizel MS, Magid D, Le T. Efficacy of three-phase bone scans in evaluating diabetic foot ulcers. *Foot Ankle Int* 1999; 20(6):347-355.

8. Melkun ET, Lewis VL, Jr. Evaluation of (111) indium-labeled autologous leukocyte scintigraphy for the diagnosis of chronic osteomyelitis in patients with grade IV pressure ulcers, as compared with a standard diagnostic protocol. *Ann Plast Surg* 2005; 54(6):633-636.
9. Vesco L, Boulahdour H, Hamissa S, et al. The value of combined radionuclide and magnetic resonance imaging in the diagnosis and conservative management of minimal or localized osteomyelitis of the foot in diabetic patients. *Metabolism* 1999; 48(7):922-927.
10. Ledermann HP, Schweitzer ME, Morrison WB. Nonenhancing tissue on MR imaging of pedal infection: characterization of necrotic tissue and associated limitations for diagnosis of osteomyelitis and abscess. *AJR* 2002; 178(1):215-222.
11. Keidar Z, Militianu D, Melamed E, Bar-Shalom R, Israel O. The diabetic foot: initial experience with 18F-FDG PET/CT. *J Nucl Med* 2005; 46(3):444-449.
12. Becker W. Imaging osteomyelitis and the diabetic foot. *Q J Nucl Med* 1999; 43(1):9-20.
13. Kapoor A, Page S, Lavalley M, Gale DR, Felson DT. Magnetic resonance imaging for diagnosing foot osteomyelitis: a meta-analysis. *Arch Intern Med* 2007; 167(2):125-132.
14. Termaat MF, Raijmakers PG, Scholten HJ, Bakker FC, Patka P, Haarman HJ. The accuracy of diagnostic imaging for the assessment of chronic osteomyelitis: a systematic review and meta-analysis. *J Bone Joint Surg Am* 2005; 87(11):2464-2471.
15. Al-Khawari HA, Al-Saeed OM, Jumaa TH, Chishti F. Evaluating diabetic foot infection with magnetic resonance imaging: Kuwait experience. *Med Princ Pract* 2005; 14(3):165-172.
16. Giurato L, Uccioli L. The diabetic foot: Charcot joint and osteomyelitis. *Nucl Med Commun* 2006; 27(9):745-749.
17. Ledermann HP, Morrison WB, Schweitzer ME, Raikin SM. Tendon involvement in pedal infection: MR analysis of frequency, distribution, and spread of infection. *AJR* 2002; 179(4):939-947.
18. Durham JR, Lukens ML, Campanini DS, Wright JG, Smead WL. Impact of magnetic resonance imaging on the management of diabetic foot infections. *Am J Surg* 1991; 162(2):150-153; discussion 153-154.
19. Horowitz JD, Durham JR, Nease DB, Lukens ML, Wright JG, Smead WL. Prospective evaluation of magnetic resonance imaging in the management of acute diabetic foot infections. *Ann Vasc Surg* 1993; 7(1):44-50.
20. American College of Radiology. *Manual on Contrast Media*. Available at: http://www.acr.org/SecondaryMainMenuCategories/quality_safety/contrast_manual.aspx.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.