

American College of Radiology ACR Appropriateness Criteria®

Clinical Condition:

Suspected Ankle Fractures

Variant 1:

Patient Meeting Ottawa Rules.

1. Inability to bear weight immediately after the injury OR
2. Point tenderness over the medial malleolus, or the posterior edge or inferior tip of the lateral malleolus or talus or calcaneus OR
3. Inability to ambulate for four steps in the emergency department.

Radiologic Procedure	Rating	Comments	RRL*
X-ray ankle	9	AP, lateral, and mortise views.	Min
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

SUSPECTED ANKLE FRACTURES

Expert Panel on Musculoskeletal Imaging: D. Lee Bennett, MD, MA¹; Richard H. Daffner, MD²; Barbara N. Weissman, MD³; Laura Bancroft, MD⁴; Judy S. Blebea, MD⁵; Ian Blair Fries, MD⁶; Jon A. Jacobson, MD⁷; William B. Morrison, MD⁸; William K. Payne III, MD, MPH⁹; Charles S. Resnik, MD¹⁰; Catherine C. Roberts, MD¹¹; Mark E. Schweitzer, MD¹²; Leanne L. Seeger, MD¹³; Mihra S. Taljanovic, MD¹⁴; James N. Wise, MD.¹⁵

Summary of Literature Review

The musculoskeletal expert panel has reviewed pertinent articles dealing with adult patients with ankle injuries. The reviewed papers were primarily concerned with missed fractures and improving fracture detection [1-5] or with the establishment of clinical criteria that would decrease the number of ankle radiographs without missing significant injuries [6-17].

Radiography

In one large series, radiographs were obtained in 89% of all patients who presented to the emergency department with a history of extremity trauma; 17% of these cases had abnormalities that altered treatment [7]. Ankle radiographs accounted for approximately 10% of all radiographs ordered in the emergency department [9]; they are the third most common study ordered and are exceeded in frequency only by chest and cervical spine radiographs [9]. Stiell and colleagues reported that more than 92% of patients with ankle trauma in the ER setting had radiographs ordered [18]. In a retrospective review of more than 600 patients, Vargish et al [17] found that fewer than 25% had adequate physical examinations, and more than 99% had radiographs. In another study, all patients for whom radiographs were ordered were subjected to a physical examination by the radiology resident; there were no significant differences in the percentages of indicated studies ordered by triage personnel and residents in the emergency department [2].

The percentage of significant injuries detected on the radiographs was equivalent for the two groups [2]. It is, therefore, not surprising that radiographs taken by nurse practitioners, nurses, and medical students had similar percentages of abnormal findings because radiographs were ordered for almost everyone seen with ankle trauma.

Gleadhill et al [10] concluded that it is possible to establish guidelines that would increase the quality and efficiency of service and influence the diagnostic skills and referral habits of physicians ordering ankle radiographs in the emergency department. In a prospective study of 500 patients with inversion injuries of the ankle, Dunlop et al [9] concluded that radiographs should be performed only for patients with distal fibula tenderness or inability to bear weight, or who are older than age 60. In these older patients, material swelling was absent in 11% of malleolar fractures and in two of four calcaneal fractures [17]. Sujitkumar et al [19] analyzed 2,000 ankle injuries and concluded that swelling alone is an unreliable indicator of injury and that patients with minimal pain and swelling who are able to bear weight do not require radiographs. Stiell et al [14-16,18,20], in a number of well-designed, elaborate papers, concluded that focal tenderness over the malleolus and the inability to bear weight will detect virtually 100% of patients with significant ankle fractures. They evaluated 1,032 patients prospectively and validated their criteria on 453 new patients [14]. They believed that if this rule is used, significant fractures could be detected with a sensitivity of 1 (100%) and a confidence level of 95% [14]. Foot and ankle radiographs could be reduced 30% without missing any significant injuries [14]. When these rules were implemented there was a decrease in the number of ankle radiographs ordered, which decreased patient waiting times and costs without patient dissatisfaction or missed fractures [20]. This study was confirmed at an independent site by Pigman et al who reported a 19% reduction in ankle and midfoot radiographs [21].

In the clinical setting, radiographs of the foot and ankle are often obtained together, even though the pain can almost always be localized to one area or another. Ordering may reflect an inadequate clinical examination; on the rare occasions when fifth-metatarsal fractures occur in association with inversion injuries of the ankle, they can be detected clinically. In the presence of an inversion injury of the ankle, foot radiographs have no role in management [22]. It is widely accepted that an adequate radiograph of the ankle should include the base of the fifth metatarsal bone distal to the tuberosity.

An evaluation of the traumatized ankle should consist of anteroposterior (AP), lateral, and mortise views of the ankle [3]. Additional views can be added to the minimal series in questionable cases. The fifth metatarsal base distal to the tuberosity should be seen on at least one projection. The use of a pertinent clinical history for the site of point tenderness will decrease the miss rate for subtle fractures by approximately 50% [12].

¹Principal Author, University of Iowa Health Center, Iowa City, Iowa.

²Panel Chair, Allegheny General Hospital, Pittsburgh, Pennsylvania.

³Panel Vice-chair, Brigham & Women's Hospital, Boston, Massachusetts.

⁴Florida Hospital, Orlando, Florida.

⁵Cleveland Clinic, Cleveland, Ohio.

⁶Bone, Spine and Hand Surgery, Chartered, Brick, New Jersey, American Academy of Orthopaedic Surgeons.

⁷University of Michigan Medical Center, Ann Arbor, Michigan.

⁸Thomas Jefferson University Hospital, Philadelphia, Pennsylvania.

⁹American Academy of Orthopaedic Surgeons, Chicago, Illinois.

¹⁰University of Maryland School of Medicine, Baltimore, Maryland.

¹¹Mayo Clinic, Phoenix, Arizona.

¹²University of Ottawa, Ottawa, Ontario, Canada.

¹³David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California.

¹⁴University of Arizona Health Sciences Center, Tucson, Arizona.

¹⁵University of Kentucky, Lexington, Kentucky.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply society endorsement of the final document.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

Ottawa Ankle Rules

The committee believed that the guidelines established by Stiell et al [14-16] and confirmed by Pijnenburg et al [13] should be adopted in the evaluation of patients with ankle trauma. These guidelines suggest obtaining ankle radiographs in patients with the following clinical findings: 1) inability to bear weight immediately after the injury, or 2) point tenderness over the medial malleolus, or the posterior edge or inferior tip of the lateral malleolus or talus or calcaneus, or 3) inability to ambulate for four steps in the emergency department. It has been convincingly demonstrated that one can approach a sensitivity of 100% in excluding significant ankle fractures using these simple criteria [10,14-16]. Limiting ankle radiographs to patients who meet these criteria can eliminate a considerable number of ankle and midfoot radiographs (estimated range 19%-36%) without missing significant injuries [10]. This would result in a considerable savings in patient cost and waiting time.

The validation and cost effectiveness of these rules have been confirmed in multiple subsequent series [6,11,13,23].

Occult Ankle Fractures

Clark et al [4,5] have shown that occult fractures of the ankle may present with a large ankle effusion (>15 mm) in the absence of a visible fracture. However, this is an uncommon imaging scenario in that it occurred in less than 1% of all the radiographs taken in the study. The vast majority of ankle radiographs with a large joint effusion had a visible fracture on the radiograph. In those rare cases in which a large joint effusion is seen on the radiograph but no fracture is visible, a computed tomography (CT) scan will demonstrate a fracture in a third of these cases.

Computed Tomography

Haapamaki et al [24] used multidetector CT (MDCT) of the ankle in multitrauma patients and compared the MDCT findings with the radiographs. When compared to MDCT, radiographs were 87% sensitive in detecting calcaneal fractures, 78% sensitive in detecting talar fractures, and 25%-33% sensitive in detecting midfoot fractures. Only 5 of 21 Lisfranc fracture dislocations were detected on radiographs. They recommended MDCT for patients with high-energy polytrauma and in those with complex foot and ankle fractures.

Magnetic Resonance Imaging

Remplik et al [25] compared low-field (0.2 Tesla) magnetic resonance imaging (MRI) and conventional radiography and found no statistical difference in the detection of acute fractures of the distal extremities. Nikken et al [26] compared clinical outcomes (need for eventual treatment of an injury) between radiography and MRI in the setting of acute ankle trauma and found that a positive radiograph was a better positive predictor of the need for treatment than a positive MRI. However, neither a negative radiograph nor a negative MRI was good at predicting lack of need for future treatment of an injury.

In summary, the three-view ankle radiographic examination is good at identifying fractures that will need immobilization and/or surgical intervention for treatment. A negative radiographic or MRI examination is not sufficient to exclude those patients who may eventually need immobilization and/or surgical intervention; therefore, clinical follow-up is essential in the patient who has suffered an acute ankle injury but has negative imaging studies.

Summary

- In a patient who meets the Ottawa Rules for a suspected ankle fracture, a three-view (AP, lateral and mortise) ankle radiographic study is indicated.
- If the radiograph is negative, clinical follow-up is warranted for ruling out an ankle injury that may eventually need treatment.

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations	
Relative Radiation Level	Effective Dose Estimate Range
None	0
Minimal	< 0.1 mSv
Low	0.1-1 mSv
Medium	1-10 mSv
High	10-100 mSv

Supporting Document(s)

- [ACR Appropriateness Criteria® Overview](#)
- [Evidence Table](#)

References

1. Berbaum KS, el-Khoury GY, Franken EA, Jr., Kathol M, Montgomery WJ, Hesson W. Impact of clinical history on fracture detection with radiography. *Radiology* 1988; 168(2):507-511.
2. Berbaum KS, Franken EA, Jr., el-Khoury GY. Impact of clinical history on radiographic detection of fractures: a comparison of radiologists and orthopedists. *AJR* 1989; 153(6):1221-1224.
3. Brandser EA, Berbaum KS, Dorfman DD, et al. Contribution of individual projections alone and in combination for radiographic detection of ankle fractures. *AJR* 2000; 174(6):1691-1697.
4. Clark TW, Janzen DL, Ho K, Grunfeld A, Connell DG. Detection of radiographically occult ankle fractures following acute trauma: positive predictive value of an ankle effusion. *AJR* 1995; 164(5):1185-1189.
5. Clark TW, Janzen DL, Logan PM, Ho K, Connell DG. Improving the detection of radiographically occult ankle fractures: positive predictive value of an ankle joint effusion. *Clin Radiol* 1996; 51(9):632-636.

6. Anis AH, Stiell IG, Stewart DG, Laupacis A. Cost-effectiveness analysis of the Ottawa Ankle Rules. *Ann Emerg Med* 1995; 26(4):422-428.
7. Auletta AG, Conway WF, Hayes CW, Guisto DF, Gervin AS. Indications for radiography in patients with acute ankle injuries: role of the physical examination. *AJR* 1991; 157(4):789-791.
8. Brand DA, Frazier WH, Kohlhepp WC, et al. A protocol for selecting patients with injured extremities who need x-rays. *N Engl J Med* 1982; 306(6):333-339.
9. Dunlop MG, Beattie TF, White GK, Raab GM, Doull RI. Guidelines for selective radiological assessment of inversion ankle injuries. *Br Med J (Clin Res Ed)* 1986; 293(6547):603-605.
10. Gleadhill DN, Thomson JY, Simms P. Can more efficient use be made of x ray examinations in the accident and emergency department? *Br Med J (Clin Res Ed)* 1987; 294(6577):943-947.
11. Keogh SP, Shafi A, Wijetunge DB. Comparison of Ottawa ankle rules and current local guidelines for use of radiography in acute ankle injuries. *J R Coll Surg Edinb* 1998; 43(5):341-343.
12. Montague AP, McQuillan RF. Clinical assessment of apparently sprained ankle and detection of fracture. *Injury* 1985; 16(8):545-546.
13. Pijnenburg AC, Glas AS, De Roos MA, et al. Radiography in acute ankle injuries: the Ottawa Ankle Rules versus local diagnostic decision rules. *Ann Emerg Med* 2002; 39(6):599-604.
14. Stiell IG, Greenberg GH, McKnight RD, et al. Decision rules for the use of radiography in acute ankle injuries. Refinement and prospective validation. *JAMA* 1993; 269(9):1127-1132.
15. Stiell IG, Greenberg GH, McKnight RD, Nair RC, McDowell I, Worthington JR. A study to develop clinical decision rules for the use of radiography in acute ankle injuries. *Ann Emerg Med* 1992; 21(4):384-390.
16. Stiell IG, McKnight RD, Greenberg GH, Nair RC, McDowell I, Wallace GJ. Interobserver agreement in the examination of acute ankle injury patients. *Am J Emerg Med* 1992; 10(1):14-17.
17. Vargish T, Clarke WR, Young RA, Jensen A. The ankle injury--indications for the selective use of X-rays. *Injury* 1983; 14(6):507-512.
18. Stiell IG, McDowell I, Nair RC, et al. Use of radiography in acute ankle injuries: physicians' attitudes and practice. *CMAJ* 1992; 147(11):1671-1678.
19. Sujitkumar P, Hadfield JM, Yates DW. Sprain or fracture? An analysis of 2000 ankle injuries. *Arch Emerg Med* 1986; 3(2):101-106.
20. Stiell IG, McKnight RD, Greenberg GH, et al. Implementation of the Ottawa ankle rules. *JAMA* 1994; 271(11):827-832.
21. Pigman EC, Klug RK, Sanford S, Jolly BT. Evaluation of the Ottawa clinical decision rules for the use of radiography in acute ankle and midfoot injuries in the emergency department: an independent site assessment. *Ann Emerg Med* 1994; 24(1):41-45.
22. Diehr P, Highley R, Dehkordi F, et al. Prediction of fracture in patients with acute musculoskeletal ankle trauma. *Med Decis Making* 1988; 8(1):40-47.
23. Wynn-Thomas S, Love T, McLeod D, et al. The Ottawa ankle rules for the use of diagnostic X-ray in after hours medical centres in New Zealand. *N Z Med J* 2002; 115(1162):U184.
24. Haapamaki VV, Kiuru MJ, Koskinen SK. Ankle and foot injuries: analysis of MDCT findings. *AJR* 2004; 183(3):615-622.
25. Remplik P, Stabler A, Merl T, Roemer F, Bohndorf K. Diagnosis of acute fractures of the extremities: comparison of low-field MRI and conventional radiography. *Eur Radiol* 2004; 14(4):625-630.
26. Nikken JJ, Oei EH, Ginai AZ, et al. Acute ankle trauma: value of a short dedicated extremity MR imaging examination in prediction of need for treatment. *Radiology* 2005; 234(1):134-142.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.