

**American College of Radiology
ACR Appropriateness Criteria®**

Clinical Condition: **Head Trauma**

Variant 1: **Minor or mild acute closed head injury (GCS ≥13), without risk factors or neurologic deficit.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
CT head without contrast	7	Known to be low yield.	Med
X-ray and/or CT cervical spine without contrast	5	See ACR Appropriateness Criteria® on <u>Suspected Spine Trauma.</u>	Med
MRI head without contrast	4		None
MRA head and neck without contrast	3	Rarely indicated with mild trauma.	None
MRA head and neck without and with contrast	3		None
CT head without and with contrast	3		Med
CTA head and neck	3	Rarely indicated with mild trauma.	Med
MRI head without and with contrast	2		None
X-ray head	1		Min
FDG-PET head	1		High
US transcranial with Doppler	1		None
INV arteriography cervicocerebral	1		Med
NUC Tc-99m HMPAO SPECT head	1		High
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Head Trauma****Variant 2:****Minor or mild acute closed head injury, focal neurologic deficit, and/or risk factors.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
CT head without contrast	9		Med
MRI head without contrast	6	For problem solving.	None
X-ray and/or CT cervical spine without contrast	6	See ACR Appropriateness Criteria® on <u>Suspected Spine Trauma.</u>	Med
MRA head and neck without contrast	5	If vascular injury is suspected. For problem solving.	None
MRA head and neck without and with contrast	5	If vascular injury is suspected. For problem solving. See comments regarding contrast in text under “Anticipated Exceptions.”	None
CTA head and neck	5	If vascular injury is suspected. For problem solving.	Med
MRI head without and with contrast	3		None
CT head without and with contrast	2		Med
NUC Tc-99m HMPAO SPECT head	1		High
FDG-PET head	1		High
US transcranial with Doppler	1		None
X-ray head	1		Min
INV arteriography cervicocerebral	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Head Trauma****Variant 3:****Moderate or severe acute closed head injury.**

Radiologic Procedure	Rating	Comments	<u>RRL</u>*
CT head without contrast	9		Med
X-ray and/or CT cervical spine without contrast	8	See ACR Appropriateness Criteria® on <i>Suspected Spine Trauma.</i>	Med
MRI head without contrast	6		None
MRA head and neck without contrast	5		None
MRA head and neck without and with contrast	5	See comments regarding contrast in text under “Anticipated Exceptions.”	None
CTA head and neck	5		Med
CT head without and with contrast	2		Med
MRI head without and with contrast	2		None
X-ray head	2		Min
US transcranial with Doppler	1		None
FDG-PET head	1		High
INV arteriography cervicocerebral	1		Med
NUC Tc-99m HMPAO SPECT head	1		High
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Head Trauma****Variant 4:****Mild or moderate acute closed head injury, child <2 years old.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
CT head without contrast	9		High
X-ray and/or CT cervical spine without contrast	7	See ACR Appropriateness Criteria® on Suspected Spine Trauma .	High
MRI head without contrast	7	Diffusion-weighted imaging especially helpful for nonaccidental trauma.	None
X-ray head	5		Min
MRI head without and with contrast	4	Potentially useful in suspected nonaccidental trauma. See comments regarding contrast in text under “Anticipated Exceptions.”	None
MRA head and neck without contrast	4	If vascular abnormality is suspected.	None
MRA head and neck without and with contrast	4	If vascular abnormality is suspected. See comments regarding contrast in text under “Anticipated Exceptions.”	None
CTA head and neck	4	If vascular abnormality is suspected.	Med
CT head without and with contrast	2		High
FDG-PET head	1		High
NUC Tc-99m HMPAO SPECT head	1		High
US transcranial with Doppler	1		None
INV arteriography cervicocerebral	1		High
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Head Trauma****Variant 5:****Subacute or chronic closed head injury with cognitive and/or neurologic deficit(s).**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
MRI head without contrast	8		None
CT head without contrast	6		Med
NUC Tc-99m HMPAO SPECT head	4	For selected cases.	High
FDG-PET head	4	For selected cases.	High
MRA head and neck without contrast	4	For selected cases.	None
MRA head and neck without and with contrast	4	For selected cases. See comments regarding contrast in text under "Anticipated Exceptions."	None
CTA head and neck	4	For selected cases.	Med
MRI head without and with contrast	3		None
CT head without and with contrast	2		Med
X-ray and/or CT cervical spine without contrast	2	Assuming there are no spinal neurologic deficits. See ACR Appropriateness Criteria® on Suspected Spine Trauma .	Med
X-ray head	2		Min
MRI functional (fMRI) head	2		None
US transcranial with Doppler	1		None
INV arteriography cervicocerebral	1		Med
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Head Trauma****Variant 6:****Closed head injury; rule out carotid or vertebral artery dissection.**

Radiologic Procedure	Rating	Comments	<u>RRL</u>*
MRA head and neck without contrast	8	Add T1 neck images.	None
MRA head and neck without and with contrast	8	Add T1 neck images. See comments regarding contrast in text under "Anticipated Exceptions."	None
MRI head without contrast	8	Include diffusion-weighted images.	None
CT head without contrast	8		Med
CTA head and neck	8		Med
CT head without and with contrast	6	Consider perfusion.	Med
INV arteriography cervicocerebral	6	For problem solving.	Med
MRI head without and with contrast	6	See comments regarding contrast in text under "Anticipated Exceptions."	None
X-ray and/or CT cervical spine without contrast	5	See ACR Appropriateness Criteria® on Suspected Spine Trauma .	Med
X-ray head	2		Min
NUC Tc-99m HMPAO SPECT head	1		High
US transcranial with Doppler	1		None
FDG-PET head	1		High
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Head Trauma****Variant 7:****Penetrating injury, stable, neurologically intact.**

Radiologic Procedure	Rating	Comments	<u>RRL*</u>
CT head without contrast	9		Med
X-ray and/or CT cervical spine without contrast	8	If neck or C-spine is site of injury. See ACR Appropriateness Criteria® on <u>Suspected Spine Trauma.</u>	Med
X-ray head	8	If calvarium is site of injury.	Min
CTA head and neck	7		Med
MRA head and neck without contrast	6	If no MRI contraindication.	None
MRA head and neck without and with contrast	6	If no MRI contraindication. See comments regarding contrast in text under “Anticipated Exceptions.”	None
INV arteriography cervicocerebral	5	If vascular injury is suspected.	Med
MRI head without contrast	5	If no MRI contraindication.	None
CT head without and with contrast	4	Consider perfusion.	Med
MRI head without and with contrast	4	If no MRI contraindication. See comments regarding contrast in text under “Anticipated Exceptions.”	None
US transcranial with Doppler	1		None
NUC Tc-99m HMPAO SPECT head	1		High
FDG-PET head	1		High
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:

Head Trauma

Variant 8:

Skull fracture.

Radiologic Procedure	Rating	Comments	RRL*
CT head without contrast	9		Med
CTA head and neck	7	If vascular injury is suspected.	Med
MRI head without contrast	6		None
X-ray and/or CT cervical spine without contrast	6	See ACR Appropriateness Criteria® on Suspected Spine Trauma .	Med
X-ray head	5	For selected cases.	Min
MRI head without and with contrast	4	Useful if infection suspected. See comments regarding contrast in text under “Anticipated Exceptions.”	None
CT head without and with contrast	4		Med
MRA head and neck without contrast	4		None
MRA head and neck without and with contrast	4	See comments regarding contrast in text under “Anticipated Exceptions.”	None
US transcranial with Doppler	1		None
NUC Tc-99m HMPAO SPECT head	1		High
INV arteriography cervicocerebral	1		Med
FDG-PET head	1		High
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

HEAD TRAUMA

Expert Panel on Neurologic Imaging: Patricia C. Davis, MD¹; James A. Brunberg, MD²; Robert Louis De La Paz, MD³; Pr. Didier Dormont⁴; John E. Jordan, MD⁵; Suresh Kumar Mukherji, MD⁶; David J. Seidenwurm, MD⁷; Patrick A Turski, MD⁸; Franz J Wippold II, MD⁹; Robert D Zimmerman, MD¹⁰; Michael A. Sloan, MD, MS.¹¹

Summary of Literature Review

Cranio-cerebral injuries are a common cause of hospital admission following trauma, and are associated with significant long-term morbidity and mortality, particularly in the adolescent and young adult populations. Neuroimaging plays an essential role in identifying and characterizing these brain injuries. Computed tomography (CT) remains essential for detecting lesions that require immediate neurosurgical intervention (ie, acute subdural hematoma) as well as those that require in-hospital observation and medical management [1]. For patients with minor head injury (Glasgow Coma Scale [GSC] score of 13-15), the New Orleans Criteria [2] and the Canadian CT Head Rule [3] are clinical guidelines with high sensitivity for detecting injuries that require neurosurgical intervention, and they offer a potential reduction in unnecessary CT scans [4-6].

Other imaging modalities such as magnetic resonance imaging (MRI) depict nonsurgical pathology not visible on CT. Single photon emission computed tomography (SPECT), positron emission tomography (PET), and transcranial Doppler (TCD) have a complementary role in the assessment of brain injury. Because cervical spine trauma may accompany a head injury, cervical spine imaging is indicated for patients with head injury who have signs, symptoms, or a mechanism of injury that might result in spinal injury, and in those who are neurologically impaired.

Skull Radiography

Masters et al [7] developed and prospectively tested a management strategy for selecting patients who may benefit from skull radiography following head trauma, and offered recommendations for selecting patients who should receive CT scanning following head injury. The effect of that study was to shift the focus of neuroimaging of head trauma away from skull radiography and toward recognition of intracranial pathology as demonstrated by

CT scanning. Skull radiography is useful for imaging of calvarial fractures, penetrating injuries, and radiopaque foreign bodies.

Computed Tomography

CT's advantages for evaluation of the head-injured patient include its sensitivity for demonstrating mass effect, ventricular size and configuration, bone injuries, and acute hemorrhage regardless of location (ie, parenchymal, subarachnoid, subdural, or epidural spaces). Other advantages include its widespread availability, rapidity of scanning, and compatibility with other medical and life support devices. Its limitations include insensitivity in detecting small and predominantly nonhemorrhagic lesions associated with trauma such as contusion, particularly when adjacent to bony surfaces (eg, frontal lobes adjacent to the orbital roof, anterior temporal lobe adjacent to the greater sphenoid wing). Likewise, diffuse axonal injuries (DAIs) that result in small focal lesions throughout the cerebral hemispheres, corpus callosum, and upper brainstem and cerebellum often go undetected on CT. CT is relatively insensitive for detecting increased intracranial pressure or cerebral edema and for early demonstration of hypoxic-ischemic encephalopathy (HIE) that may accompany moderate or severe head injury. Potential risks of unnecessary exposure to ionizing radiation warrant judicious patient selection for CT scanning as well as radiation dose management [8].

There is now a general consensus that patients identified as having moderate or high risk for intracranial injury should undergo early postinjury noncontrast CT for evidence of intracerebral hematoma, midline shift, or increased intracranial pressure. A number of clinical criteria similar to those of Masters et al [7] are used to predict patient risk categories for intracranial injury. There is an inverse relationship between declining clinical or neurologic status as described by the GCS [9] and the incidence and severity of CT abnormalities related to head injury [10-12].

Although experienced physicians can often predict the likelihood of an abnormal CT scan in moderate or severe head injury, clinical selection criteria of patients with minor or mild injury (ie, GCS score ≥ 13) who harbor significant intracranial pathology and/or require acute surgical intervention have been problematic. Rapid CT scanning is readily available in most hospitals that treat head-injured patients; thus the routine use of CT has been advocated as a screening tool to triage minor or mild head-injured patients who require hospital admission or surgical intervention from those who can be safely discharged without hospital admission [13-15]. Although CT triage of head-injured patients who require hospital admission offers a reduced burden on inpatient hospital

¹Principal Author and Panel Chair, Northwest Radiology Consultants, Atlanta, Ga; ²University of California-Davis Medical Center, Sacramento, Calif; ³Columbia University Medical Center, New York, NY; ⁴Hôpital de la Salpêtrière, Assistance-Publique-Hôpitaux de Paris, France; ⁵Memrad Medical Group, Inc., Long Beach, Calif; ⁶University of Michigan Health System, Ann Arbor, Mich; ⁷Radiological Associates of Sacramento, Sacramento, Calif; ⁸University of Wisconsin, Madison, Wis; ⁹Mallinckrodt Institute of Radiology, Saint Louis, MO; ¹⁰New York Hospital-Cornell University Medical Center, New York, NY; ¹¹Carolinas Medical Center, Charlotte, NC, American Academy of Neurology.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

services at lower cost than routine hospital admission for observation, the result is greater CT use in the emergency setting [11,13-15]. In the minor head injury setting with a GCS score of 15, the New Orleans Criteria [2] found a 100% sensitivity for CT identification of an acute trauma lesion using risk factors of headache, vomiting, drug or alcohol intoxication, older than age 60, short-term memory deficit, physical findings of supraclavicular trauma, and/or seizure. Stiell et al [3] reported 100% sensitivity for detecting neurosurgical and/or clinically important brain injury in subjects with a GCS score of 13-15 based on high-risk factors of failure to reach a GCS score of 15 within 2 hours, suspected open skull fracture, two or more vomiting episodes, sign of basal skull fracture, or age ≥ 65 years.

Clinical criteria for scanning of children with head injury have been less reliable than those for adults, particularly for children younger than age two [16,17]. For this reason, more liberal use of CT scanning has been suggested for pediatric patients. On the other hand, this must be balanced with the higher risk of radiation exposure in childhood via judicious patient selection for scanning as well as management of radiation dose [8,18,19]. Noncontrast head CT plays an essential role in the evaluation of children with suspected physical injury from child abuse; appropriateness criteria for imaging of child abuse have already been described (see the pediatric sections of the ACR Appropriateness Criteria[®]).

Early and sometimes repeated CT scanning may be required in cases of clinical or neurologic deterioration, especially in the first 72 hours after head injury, to detect delayed hematoma, hypoxic-ischemic lesions, or cerebral edema [20]. CT has a role in subacute or chronic head injury for depicting atrophy, focal encephalomalacia, hydrocephalus, and chronic subdural hematoma.

Cerebral Angiography

Since the development of CT in the mid-1970s, the need for cerebral angiography for head injury has dramatically declined. Cerebral angiography has a role in demonstrating and managing traumatic vascular injuries such as pseudoaneurysm, dissection, or diagnosis and neurointerventional treatment of uncontrolled hemorrhage. Vascular injuries typically occur with penetrating trauma (eg, gunshot wound or stabbing), basal skull fracture, or trauma to the neck [21-23].

Dynamic spiral CT angiography (CTA) and magnetic resonance angiography (MRA) have a role as less invasive screening tools for detecting traumatic vascular lesions. MRA and fat-suppressed T1-weighted MR [22] or CTA may reveal carotid or vertebral dissection, although angiography remains the gold standard for depicting dissection. Cerebral infarction is an infrequent accompaniment to head injury, and patterns of infarction suggest that direct vascular compression related to

intracranial mass lesions is the most common underlying mechanism [24].

Magnetic Resonance Imaging

Although the role of MRI in imaging of head trauma is growing, its use is hindered by its limited availability in the acute trauma setting, long imaging times, sensitivity to patient motion, incompatibility with various medical and life support devices, and relative insensitivity to subarachnoid hemorrhage. Other factors include the need for MRI-specific monitoring equipment and ventilators, and the risk of scanning patients with certain indwelling devices (eg, cardiac pacemaker, cerebral aneurysm clip) or occult foreign bodies. In part, these limitations can be overcome by situating MRI scanners close to emergency care areas with appropriate design and equipment for managing acutely injured patients [25,26]. MRI advances such as open-bore geometry, faster imaging sequences, and improved patient monitoring equipment allow a greater role for MRI in closed head injuries.

MRI is very sensitive for detecting and characterizing subacute and chronic brain injuries. The number, size, and location of MR abnormalities in subacute head injury have been used to predict the recovery outcome of post-traumatic vegetative state [27]. While CT is sensitive for detecting injuries requiring a change in treatment [28], MRI also is used for acute head-injured patients with nonsurgical, medically stable pathology. Hemosiderin-sensitive T2-weighted gradient echo sequences are helpful for imaging small or subacute or chronic hemorrhages. Diffusion sequences improve detection of acute infarction associated with head injury. Fluid-attenuated inversion recovery (FLAIR) images are more sensitive than conventional MRI sequences for depicting subarachnoid hemorrhage and for lesions bordered by cerebrospinal fluid (CSF) [29]. MRA is helpful for screening vascular lesions such as thromboses, pseudoaneurysms, or dissection. Lang et al [30] found that the addition of gadolinium enhancement offered no significant advantage for lesion detection or characterization compared with noncontrast MRI images in head-injury patients.

The soft-tissue detail offered by MRI is superior to that of CT for depicting nonhemorrhagic primary lesions such as contusions, for detecting secondary effects of trauma such as edema and hypoxic-ischemic encephalopathy, and for imaging DAI [31-33]. DAI results from a shear-strain pattern of acceleration-deceleration with characteristic lesions in increasing order of injury severity in the: 1) cerebral white matter and gray-white matter junction; 2) corpus callosum, particularly the splenium; and 3) dorsal upper brain stem and cerebellum [31,34].

Although management of surgical injuries is not likely to be altered by the substitution of MRI for CT [28], superior depiction of nonsurgical lesions with MRI may affect medical management and predict the degree of

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

neurologic recovery [31,35]. Diffusion-weighted MRI and apparent diffusion coefficient (ADC) mapping depict cytotoxic injury almost immediately. In acute brain trauma, focal contusion and DAI may show restricted diffusion and evolve over time to atrophy or encephalomalacia [36,37]. Perfusion imaging with CT or MRI may prove helpful as a marker for disorders of vascular autoregulation or ischemia [38]. Diffusion tensor imaging and MR spectroscopy (MRS) are ancillary tools that may offer additional insight into the biochemical and structural patterns of injury following head trauma, as well as prognosis [34,39].

Other Imaging Modalities

A few reports of selected head-injury subjects suggest a role for functional imaging techniques (SPECT, PET, xenon-enhanced CT, functional MRI) to assess cognitive and neuropsychologic disturbances as well as recovery following head trauma [40-44]. SPECT studies may reveal focal areas of hypoperfusion that are discordant with findings of MRI or CT [42-44]. On the basis of these results, some investigators suggest that these functional imaging techniques may explain or predict post-injury neuropsychologic and cognitive deficits that are not explained by MRI or CT abnormalities [40,41,43]. Furthermore, focal lesions demonstrated by SPECT offer objective evidence of organic injury in patients whose neuroimaging studies are otherwise normal [41]. Oder et al [45] found that a pattern of global reduction of cerebral blood flow detected by SPECT predicted a poor likelihood of recovery for patients who are in a persistent vegetative state due to head injury. SPECT and PET do not provide the anatomic detail or image resolution of CT or MRI for demonstrating acute or neurosurgical lesions of closed head injury, so their use is generally limited to subacute or chronic patients.

Transcranial Doppler sonography (TCD) offers a noninvasive bedside evaluation of cerebral blood flow velocity and resistance in the major proximal vessels of the circle of Willis. Several investigators have suggested that TCD can be used to monitor early changes in blood flow velocities that may relate to vasospasm, hypervolemia, low velocity state, or edema [46-48], especially in management of the acutely brain-injured patient.

Anticipated Exceptions

Nephrogenic systemic fibrosis (NSF), also known as nephrogenic fibrosing dermopathy) was first identified in 1997 and has recently generated substantial concern among radiologists, referring doctors and lay people. Until the last few years, gadolinium-based MR contrast agents were widely believed to be almost universally well tolerated, extremely safe and non-nephrotoxic, even when used in patients with impaired renal function. All available experience suggests that these agents remain generally very safe, but recently some patients with renal

failure who have been exposed to gadolinium contrast agents (the percentage is unclear) have developed NSF [49-51], a syndrome that can be fatal. Further studies are necessary to determine what the exact relationships are between gadolinium-containing contrast agents, their specific components and stoichiometry, patient renal function and NSF. Current theory links the development of NSF to the administration of relatively high doses (eg, >0.2mM/kg) and to agents in which the gadolinium is least strongly chelated. The FDA has recently issued a “black box” warning concerning these contrast agents (http://www.fda.gov/cder/drug/InfoSheets/HCP/gcca_200705HCP.pdf).

This warning recommends that, until further information is available, gadolinium contrast agents should not be administered to patients with either acute or significant chronic kidney disease (estimated GFR <30 mL/min/1.73m²), recent liver or kidney transplant or hepato-renal syndrome, unless a risk-benefit assessment suggests that the benefit of administration in the particular patient clearly outweighs the potential risk(s) [50].

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations	
Relative Radiation Level	Effective Dose Estimate Range
None	0
Minimal	< 0.1 mSv
Low	0.1-1 mSv
Medium	1-10 mSv
High	10-100 mSv

References

1. Saul TG, and Joint Section on Neurotrauma and Critical Care of the American Association of Neurological Surgeons and Congress of Neurological Surgeons. Management of Head Injury, 1998.
2. Haydel MJ, Preston CA, Mills TJ, Luber S, Blaudeau E, DeBlieux PM. Indications for computed tomography in patients with minor head injury. *N Engl J Med* 2000; 343(2):100-105.
3. Stiell IG, Wells GA, Vandemheen K, et al. The Canadian CT Head Rule for patients with minor head injury. *Lancet* 2001; 357(9266):1391-1396.
4. Haydel MJ. Clinical decision instruments for CT scanning in minor head injury. *JAMA* 2005; 294(12):1551-1553.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

5. Smits M, Dippel DW, de Haan GG, et al. External validation of the Canadian CT Head Rule and the New Orleans Criteria for CT scanning in patients with minor head injury. *JAMA* 2005; 294(12):1519-1525.
6. Stiell IG, Clement CM, Rowe BH, et al. Comparison of the Canadian CT Head Rule and the New Orleans Criteria in patients with minor head injury. *JAMA* 2005; 294(12):1511-1518.
7. Masters SJ, McClean PM, Arcarese JS, et al. Skull x-ray examinations after head trauma. Recommendations by a multidisciplinary panel and validation study. *N Engl J Med* 1987; 316(2):84-91.
8. National Cancer Institute USNIoH. Radiation risks and pediatric computed tomography (CT): a guide for health care providers. August 20,2002; <http://www.nci.nih.gov/cancertopics/causes/radiation-risks-pediatric-CT>.
9. Teasdale G, Jennett B. Assessment of coma and impaired consciousness. A practical scale. *Lancet* 1974; 2(7872):81-84.
10. Kido DK, Cox C, Hamill RW, Rothenberg BM, Woolf PD. Traumatic brain injuries: predictive usefulness of CT. *Radiology* 1992; 182(3):777-781.
11. Reinius WR, Zwemer FL, Jr., Fornoff JR. Prospective optimization of patient selection for emergency cranial computed tomography: univariate and multivariate analyses. *Invest Radiol* 1996; 31(2):101-108.
12. Shackford SR, Wald SL, Ross SE, et al. The clinical utility of computed tomographic scanning and neurologic examination in the management of patients with minor head injuries. *J Trauma* 1992; 33(3):385-394.
13. Livingston DH, Loder PA, Hunt CD. Minimal head injury: is admission necessary? *Am Surg* 1991; 57(1):14-17.
14. Nagy KK, Joseph KT, Krosner SM, et al. The utility of head computed tomography after minimal head injury. *J Trauma* 1999; 46(2):268-270.
15. Stein SC, O'Malley KF, Ross SE. Is routine computed tomography scanning too expensive for mild head injury? *Ann Emerg Med* 1991; 20(12):1286-1289.
16. Dietrich AM, Bowman MJ, Ginn-Pease ME, Kosnik E, King DR. Pediatric head injuries: can clinical factors reliably predict an abnormality on computed tomography? *Ann Emerg Med* 1993; 22(10):1535-1540.
17. Homer CJ, Kleinman L. Technical report: minor head injury in children. *Pediatrics* 1999; 104(6):e78.
18. The ALARA (as low as reasonably achievable) concept in pediatric CT intelligent dose reduction. Multidisciplinary conference organized by the Society of Pediatric Radiology. August 18-19, 2001. *Pediatr Radiol* 2002; 32(4):217-313.
19. Paterson A, Frush DP, Donnelly LF. Helical CT of the body: are settings adjusted for pediatric patients? *AJR* 2001; 176(2):297-301.
20. Stein SC, Spettell C, Young G, Ross SE. Delayed and progressive brain injury in closed-head trauma: radiological demonstration. *Neurosurgery* 1993; 32(1):25-30; discussion 30-21.
21. Gaskill-Shiple MF, Tomsick TA. Angiography in the evaluation of head and neck trauma. *Neuroimaging Clin N Am* 1996; 6(3):607-624.
22. Ozdoba C, Sturzenegger M, Schroth G. Internal carotid artery dissection: MR imaging features and clinical-radiologic correlation. *Radiology* 1996; 199(1):191-198.
23. Showalter W, Esekogwu V, Newton KI, Henderson SO. Vertebral artery dissection. *Acad Emerg Med* 1997; 4(10):991-995.
24. Mirvis SE, Wolf AL, Numaguchi Y, Corradino G, Joslyn JN. Posttraumatic cerebral infarction diagnosed by CT: prevalence, origin, and outcome. *AJNR* 1990; 11(2):355-360.
25. Gentry LR, Godersky JC, Thompson B, Dunn VD. Prospective comparative study of intermediate-field MR and CT in the evaluation of closed head trauma. *AJR* 1988; 150(3):673-682.
26. Mittl RL, Grossman RI, Hiehle JF, et al. Prevalence of MR evidence of diffuse axonal injury in patients with mild head injury and normal head CT findings. *AJNR* 1994; 15(8):1583-1589.
27. Kampfl A, Schmutzhard E, Franz G, et al. Prediction of recovery from post-traumatic vegetative state with cerebral magnetic-resonance imaging. *Lancet* 1998; 351(9118):1763-1767.
28. Fiser SM, Johnson SB, Fortune JB. Resource utilization in traumatic brain injury: the role of magnetic resonance imaging. *Am Surg* 1998; 64(11):1088-1093.
29. Ashikaga R, Araki Y, Ishida O. MRI of head injury using FLAIR. *Neuroradiology* 1997; 39(4):239-242.
30. Lang DA, Hadley DM, Teasdale GM, Macpherson P, Teasdale E. Gadolinium DTPA enhanced magnetic resonance imaging in acute head injury. *Acta Neurochir (Wien)* 1991; 109(1-2):5-11.
31. Gentry LR. Imaging of closed head injury. *Radiology* 1994; 191(1):1-17.
32. Gentry LR, Godersky JC, Thompson B. MR imaging of head trauma: review of the distribution and radiopathologic features of traumatic lesions. *AJR* 1988; 150(3):663-672.
33. Gentry LR, Thompson B, Godersky JC. Trauma to the corpus callosum: MR features. *AJNR* 1988; 9(6):1129-1138.
34. Arfanakis K, Haughton VM, Carew JD, Rogers BP, Dempsey RJ, Meyerand ME. Diffusion tensor MR imaging in diffuse axonal injury. *AJNR* 2002; 23(5):794-802.
35. Doezema D, King JN, Tandberg D, Espinosa MC, Orrison WW. Magnetic resonance imaging in minor head injury. *Ann Emerg Med* 1991; 20(12):1281-1285.
36. Rugg-Gunn FJ, Symms MR, Barker GJ, Greenwood R, Duncan JS. Diffusion imaging shows abnormalities after blunt head trauma when conventional magnetic resonance imaging is normal. *J Neurol Neurosurg Psychiatry* 2001; 70(4):530-533.
37. Smith DH, Meaney DF, Lenkinski RE, et al. New magnetic resonance imaging techniques for the evaluation of traumatic brain injury. *J Neurotrauma* 1995; 12(4):573-577.
38. Wintermark M, Chiolero R, van Melle G, et al. Relationship between brain perfusion computed tomography variables and cerebral perfusion pressure in severe head trauma patients. *Crit Care Med* 2004; 32(7):1579-1587.
39. Brooks WM, Friedman SD, Gasparovic C. Magnetic resonance spectroscopy in traumatic brain injury. *J Head Trauma Rehabil* 2001; 16(2):149-164.
40. Ichise M, Chung DG, Wang P, Wortzman G, Gray BG, Franks W. Technetium-99m-HMPAO SPECT, CT and MRI in the evaluation of patients with chronic traumatic brain injury: a correlation with neuropsychological performance. *J Nucl Med* 1994; 35(2):217-226.
41. Jacobs A, Put E, Ingels M, Bossuyt A. Prospective evaluation of technetium-99m-HMPAO SPECT in mild and moderate traumatic brain injury. *J Nucl Med* 1994; 35(6):942-947.
42. Jantzen KJ, Anderson B, Steinberg FL, Kelso JA. A prospective functional MR imaging study of mild traumatic brain injury in college football players. *AJNR* 2004; 25(5):738-745.
43. Kinuya K, Kakuda K, Nobata K, et al. Role of brain perfusion single-photon emission tomography in traumatic head injury. *Nucl Med Commun* 2004; 25(4):333-337.
44. Moritz CH, Rowley HA, Haughton VM, Swartz KR, Jones J, Badie B. Functional MR imaging assessment of a non-responsive brain injured patient. *Magn Reson Imaging* 2001; 19(8):1129-1132.
45. Oder W, Goldenberg G, Podreka I, Deecke L. HM-PAO-SPECT in persistent vegetative state after head injury: prognostic indicator of the likelihood of recovery? *Intensive Care Med* 1991; 17(3):149-153.
46. Muttaqin Z, Uozumi T, Kuwabara S, et al. Hyperaemia prior to acute cerebral swelling in severe head injuries: the role of transcranial Doppler monitoring. *Acta Neurochir (Wien)* 1993; 123(1-2):76-81.
47. Steiger HJ, Aaslid R, Stooss R, Seiler RW. Transcranial Doppler monitoring in head injury: relations between type of injury, flow velocities, vasoreactivity, and outcome. *Neurosurgery* 1994; 34(1):79-85; discussion 85-76.
48. van Santbrink H, Schouten JW, Steyerberg EW, Avezaat CJ, Maas AI. Serial transcranial Doppler measurements in traumatic brain injury with special focus on the early posttraumatic period. *Acta Neurochir (Wien)* 2002; 144(11):1141-1149.
49. Broome DR, Girguis MS, Baron PW, Cottrell AC, Kjellin I, Kirk GA. Gadodiamide-associated nephrogenic systemic fibrosis: why radiologists should be concerned. *AJR* 2007; 188(2):586-592.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

50. Kanal E, Barkovich AJ, Bell C, et al. ACR guidance document for safe MR practices: 2007. *AJR* 2007; 188(6):1447-1474.

51. Sadowski EA, Bennett LK, Chan MR, et al. Nephrogenic systemic fibrosis: risk factors and incidence estimation. *Radiology* 2007; 243(1):148-157.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.