

American College of Radiology ACR Appropriateness Criteria®

LOCAL EXCISION IN EARLY-STAGE RECTAL CANCER

Expert Panel on Radiation Oncology–Rectal/Anal Cancer:
A. William Blackstock, MD¹; Suzanne M. Russo, MD²;
Andre A. Konski, MD³; W. Warren Suh, MD⁴;
Bard C. Cosman, MD⁵; Joseph Herman, MD, MSc⁶;
Mohammed Mohiuddin, MD⁷; Matthew M. Poggi, MD⁸;
William F. Regine, MD⁹; Leonard Saltz, MD¹⁰;
William Small Jr, MD¹¹; Jennifer Zook, MD.¹²

Summary of Literature Review

Background

Thirty-four percent of patients diagnosed with rectal cancer present with American Joint Commission on Cancer (AJCC) stage I disease [1]. Historically these patients have been treated with low anterior resection (LAR) or abdominoperineal resection (APR) with excellent local control (LC) and survival rates [2-4]. Postulating that early-stage lesions may not warrant such aggressive treatment as well as acknowledging the mortality and morbidity of these procedures, investigators have examined less morbid sphincter-sparing approaches such as local excision (LE). In addition, LE has been presented as an option to patients whose other comorbid conditions would not allow them to tolerate more extensive surgery. Most of the data supporting the use of LE are from single-institution, retrospective reviews [5-9]. Recently, a few prospective multi-institutional trials have investigated the efficacy of LE with or without radiation therapy (RT) in these patients [10-11].

Workup

All patients should receive a full colonoscopy with biopsy, pathology review, proctoscopy, carcinoembryonic antigen (CEA), and computerized tomography of the chest, abdomen, and pelvis. In addition, patients being considered for LE should have an endorectal ultrasound

(EUS) to evaluate depth of penetration. EUS is 62%-92% accurate for T staging and 64%-88% accurate for N staging but is highly operator dependent [12-13].

Surgical Techniques

There are three operative approaches for LE of a distal rectal lesion: transanal, posterior trans-sphincteric (York-Mason procedure), or posterior proctotomy (Kraske procedure). Transanal excision (TAE) is the most commonly used approach. Under direct visualization, the lesion is excised with a 1 cm margin including the perirectal fat. The mural defect is then closed. The posterior trans-sphincteric and posterior proctotomy approaches are used less commonly and involve posterior approaches with dissection above or below the levator ani to the rectum [14]. It is important to note that none of these procedures include lymph node evaluation. Transanal endoscopic microsurgery (TEM) allows locally complete excision of rectal neoplasms and has recently been evaluated for curative treatment of invasive cancer. TEM has been shown to be as effective, and possibly better than, conventional TAE and safe to be following chemoradiation [15-21]. (See [Variant 1](#).)

Patient Selection

In general, the best candidates for LE include small (<4 cm), low-lying tumors confined to the muscularis propria. Patients with adverse pathologic features (signet ring histology, poor differentiation, lymphovascular space invasion) or whose tumors occupy >40% of the rectum are at high risk for local recurrence, and LE is not recommended [22-23]. Patients with positive margins after LE or piecemeal resections are at very high risk of local recurrence and should be offered immediate APR or LAR. Patients with tumors invading through the muscularis propria (T3) are at very high risk (>30%) for local recurrence following LE and should not be treated with LE. Palliative LE may be performed in advanced-stage patients [24]. (See [Variant 2](#).)

Local Excision with or without Radiation Therapy

Single-institution reviews have reported failure rates of 7%-40% and 25%-62% for LE alone in T1 and T2 tumors, respectively [5-9,17-18,20,25-33]. Postoperative RT may lower these rates to 10%-20% [5,9,11,23,25], however there is little data to suggest the role of prognostic factors to select patients who are at risk for adjuvant treatment. Retrospective studies have investigated the risk of lymph node metastasis according to the depth of tumor invasion in patients with T1 and T2 tumors undergoing resection for rectal cancer, which may have important implications for patients with superficial early rectal cancers in whom LE is being considered. These studies indicate that the risk of lymph node metastases cannot be predicted by radiological staging or

¹Principal Author, Wake Forest University, Winston Salem, North Carolina.

²Research Author, East Carolina University Brody School of Medicine, Greenville, North Carolina.

³Panel Chair, Wayne State University School of Medicine, Detroit, Michigan.

⁴Panel Vice-chair, Dana-Farber Cancer Institute/Brigham & Women's Hospital, Boston, Massachusetts.

⁵VA Medical Center/University of California-San Diego, San Diego, California, American College of Surgeons.

⁶Sidney Kimmel Cancer Center at Johns Hopkins, Baltimore, Maryland.

⁷University of Kentucky Medical Center, Lexington, Kentucky.

⁸National Navy Medical Center, Bethesda, Maryland.

⁹University of Maryland Medical Center, Baltimore, Maryland.

¹⁰Memorial-Sloan Kettering Cancer Center, New York, NY, American Society of Clinical Oncology.

¹¹The Robert H. Lurie Comprehensive Cancer Center of Northwestern University, Chicago, Illinois.

¹²Indiana University School of Medicine, Indianapolis, Indiana.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

tumor size; however, multivariate analysis identified the extramural vascular invasion and tumor grade as independent predictors of lymph node metastasis, but not the depth of submucosal invasion [34-35].

An initial phase II study by the Radiation Oncology Therapy Group[®] (ROG[®]) assigned patients observation (low-grade T1 tumors with negative margins) or chemoradiation (54-65 Gy with 5-fluorouracil (5-FU) 1,000 mg/m² IV d1-3, d29-31) based on postexcision pathology [10]. Local recurrence rates were 7%, 8%, and 23% for T1, T2, and T3 tumors, respectively. Cancer and Leukemia Group B study (CALGB 8984) evaluated the role of LE with or without chemotherapy and RT in 177 patients with T1 and T2 adenocarcinomas of the rectum [11]. T1 patients underwent LE followed by observation. T2 patients underwent LE followed by RT (54 Gy/30 fractions) and chemotherapy (5-FU 500 mg/m² IV d1-3, d29-31). At 48 months of median follow-up, the 6-year overall survival rate was 85%, and the disease-free survival rate was 78% for all patients. Three of the 59 eligible T1 patients and seven of the 51 eligible T2 patients had experienced local failure. It is important to note, however, that these were highly selected patients and that one-third of patients were excluded after surgery due to large tumor size and/or questionable margin status. (See [Variant 3.](#))

More recently, LE or TEM following neoadjuvant radiation with or without chemotherapy has been reported. Data from four retrospective studies [17,19,32,36] and one prospective study [18] have demonstrated safety and LC rates ranging from 2.0%-13.2%. No studies to date have prospectively evaluated whether or not the use of neoadjuvant therapy reduces recurrence rates compared to LE or TEM alone.

Future Directions

A multi-institutional phase II trial by the American College of Surgeons Oncology Group (ACSOG Z6041) is investigating neoadjuvant chemoradiation and LE in T2 patients. As of October 2009, accrual to ACSOG Z6041 has been temporarily suspended due to a recent data review that revealed the toxicity threshold written into the toxicity protocol stopping rule has been crossed in the cohort of patients receiving the revised treatment regimen. Nevertheless, the study team has determined that current patients are not at undue risk, and there are continuing plans for the study once all the adverse events have been reviewed.

Supporting Document(s)

- [ACR Appropriateness Criteria[®] Overview](#)
- [Evidence Table](#)

References

1. American Cancer Society. *Cancer Facts & Figures 2009*. Atlanta: American Cancer Society; 2009.
2. Heald RJ, Moran BJ, Ryall RD, Sexton R, MacFarlane JK. Rectal cancer: the Basingstoke experience of total mesorectal excision, 1978-1997. *Arch Surg* 1998; 133(8):894-899.
3. Martling AL, Holm T, Rutqvist LE, Moran BJ, Heald RJ, Cedemark B. Effect of a surgical training programme on outcome

- of rectal cancer in the County of Stockholm. Stockholm Colorectal Cancer Study Group, Basingstoke Bowel Cancer Research Project. *Lancet* 2000; 356(9224):93-96.
4. Wibe A, Rendedal PR, Svensson E, et al. Prognostic significance of the circumferential resection margin following total mesorectal excision for rectal cancer. *Br J Surg* 2002; 89(3):327-334.
5. Chakravarti A, Compton CC, Shellito PC, et al. Long-term follow-up of patients with rectal cancer managed by local excision with and without adjuvant irradiation. *Ann Surg* 1999; 230(1):49-54.
6. Endreseth BH, Myrvold HE, Romundstad P, Hestvik UE, Bjerkeset T, Wibe A. Transanal excision vs. major surgery for T1 rectal cancer. *Dis Colon Rectum* 2005; 48(7):1380-1388.
7. Madbouly KM, Remzi FH, Erkek BA, et al. Recurrence after transanal excision of T1 rectal cancer: should we be concerned? *Dis Colon Rectum* 2005; 48(4):711-719; discussion 719-721.
8. Paty PB, Nash GM, Baron P, et al. Long-term results of local excision for rectal cancer. *Ann Surg* 2002; 236(4):522-529; discussion 529-530.
9. Wentworth S, Russell GB, Tuner, II, et al. Long-term results of local excision with and without chemoradiation for adenocarcinoma of the rectum. *Clin Colorectal Cancer* 2005; 4(5):332-335.
10. Russell AH, Harris J, Rosenberg PJ, et al. Anal sphincter conservation for patients with adenocarcinoma of the distal rectum: long-term results of radiation therapy oncology group protocol 89-02. *Int J Radiat Oncol Biol Phys* 2000; 46(2):313-322.
11. Greenberg JA, Shibata D, Herndon JE, 2nd, Steele GD, Jr., Mayer R, Bleday R. Local excision of distal rectal cancer: an update of cancer and leukemia group B 8984. *Dis Colon Rectum* 2008; 51(8):1185-1191; discussion 1191-1184.
12. Kim HJ, Wong WD. Role of endorectal ultrasound in the conservative management of rectal cancers. *Semin Surg Oncol* 2000; 19(4):358-366.
13. Schaffzin DM, Wong WD. Endorectal ultrasound in the preoperative evaluation of rectal cancer. *Clin Colorectal Cancer* 2004; 4(2):124-132.
14. Rothenberger DA, Ricciardi R. Procedures for Rectal Cancer. In: Souba WW, Fink MP, Furkovich GJ, et al., eds. *ACS Surgery: Principles & Practice*. Vol 4: WebMD; 2004:A.D.:1-16.
15. Bach SP, Hill J, Monson JR, et al. A predictive model for local recurrence after transanal endoscopic microsurgery for rectal cancer. *Br J Surg* 2009; 96(3):280-290.
16. Christoforidis D, Cho HM, Dixon MR, Mellgren AF, Madoff RD, Finne CO. Transanal endoscopic microsurgery versus conventional transanal excision for patients with early rectal cancer. *Ann Surg* 2009; 249(5):776-782.
17. Guerrieri M, Baldarelli M, Organetti L, et al. Transanal endoscopic microsurgery for the treatment of selected patients with distal rectal cancer: 15 years experience. *Surg Endosc* 2008; 22(9):2030-2035.
18. Lezoche G, Baldarelli M, Guerrieri M, et al. A prospective randomized study with a 5-year minimum follow-up evaluation of transanal endoscopic microsurgery versus laparoscopic total mesorectal excision after neoadjuvant therapy. *Surg Endosc* 2008; 22(2):352-358.
19. Marks JH, Valsdottir EB, DeNittis A, et al. Transanal endoscopic microsurgery for the treatment of rectal cancer: comparison of wound complication rates with and without neoadjuvant radiation therapy. *Surg Endosc* 2009; 23(5):1081-1087.
20. Moore JS, Cataldo PA, Osler T, Hyman NH. Transanal endoscopic microsurgery is more effective than traditional transanal excision for resection of rectal masses. *Dis Colon Rectum* 2008; 51(7):1026-1030; discussion 1030-1021.
21. Palma P, Horisberger K, Joos A, Rothenhoefer S, Willeke F, Post S. Local excision of early rectal cancer: is transanal endoscopic microsurgery an alternative to radical surgery? *Rev Esp Enferm Dig* 2009; 101(3):172-178.
22. Willett CG, Compton CC, Shellito PC, Efrid JT. Selection factors for local excision or abdominoperineal resection of early stage rectal cancer. *Cancer* 1994; 73(11):2716-2720.
23. Willett CG, Tepper JE, Donnelly S, et al. Patterns of failure following local excision and local excision and postoperative radiation therapy for invasive rectal adenocarcinoma. *J Clin Oncol* 1989; 7(8):1003-1008.
24. Chen H, George BD, Kaufman HS, Malaki MB, Mortensen NJ, Kettlewell MG. Endoscopic transanal resection provides palliation

- equivalent to transabdominal resection in patients with metastatic rectal cancer. *J Gastrointest Surg* 2001; 5(3):282-286.
25. Borschitz T, Gockel I, Kiesslich R, Junginger T. Oncological outcome after local excision of rectal carcinomas. *Ann Surg Oncol* 2008; 15(11):3101-3108.
 26. Borschitz T, Heintz A, Junginger T. Transanal endoscopic microsurgical excision of pT2 rectal cancer: results and possible indications. *Dis Colon Rectum* 2007; 50(3):292-301.
 27. Bretagnol F, Merrie A, George B, Warren BF, Mortensen NJ. Local excision of rectal tumours by transanal endoscopic microsurgery. *Br J Surg* 2007; 94(5):627-633.
 28. Folkesson J, Johansson R, Pahlman L, Gunnarsson U. Population-based study of local surgery for rectal cancer. *Br J Surg* 2007; 94(11):1421-1426.
 29. Garcia-Aguilar J, Mellgren A, Sirivongs P, Buie D, Madoff RD, Rothenberger DA. Local excision of rectal cancer without adjuvant therapy: a word of caution. *Ann Surg* 2000; 231(3):345-351.
 30. Lezoche E, Baldarelli M, De Sanctis A, Lezoche G, Guerrieri M. Early rectal cancer: definition and management. *Dig Dis* 2007; 25(1):76-79.
 31. Min BS, Kim NK, Ko YT, et al. Long-term oncologic results of patients with distal rectal cancer treated by local excision with or without adjuvant treatment. *Int J Colorectal Dis* 2007; 22(11):1325-1330.
 32. Nash GM, Weiser MR, Guillem JG, et al. Long-term survival after transanal excision of T1 rectal cancer. *Dis Colon Rectum* 2009; 52(4):577-582.
 33. You YN, Baxter NN, Stewart A, Nelson H. Is the increasing rate of local excision for stage I rectal cancer in the United States justified?: a nationwide cohort study from the National Cancer Database. *Ann Surg* 2007; 245(5):726-733.
 34. Perez RO, Habr-Gama A, Proscurshim I, et al. Local excision for ypT2 rectal cancer--much ado about something. *J Gastrointest Surg* 2007; 11(11):1431-1438; discussion 1438-1440.
 35. Rasheed S, Bowley DM, Aziz O, et al. Can depth of tumour invasion predict lymph node positivity in patients undergoing resection for early rectal cancer? A comparative study between T1 and T2 cancers. *Colorectal Dis* 2008; 10(3):231-238.
 36. Borschitz T, Wachtlin D, Mohler M, Schmidberger H, Junginger T. Neoadjuvant chemoradiation and local excision for T2-3 rectal cancer. *Ann Surg Oncol* 2008; 15(3):712-720.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:

Local Excision in Early Stage Rectal Cancer

Variant 1:

57-year-old male with preoperative stage uT1N0, freely mobile, moderately differentiated adenocarcinoma. Tumor is 2 cm in diameter, involves <25% of circumference, and is located 6 cm from anal verge. No lymphovascular space invasion is noted.

Treatment	Rating	Comments
Local Excision, pT1N0 and Negative Margins		
Observation	9	
RT alone	2	
RT + chemotherapy	1	
Local Excision, pT1N0 and Positive Margins		
LAR or APR	9	
RT alone	2	
RT + chemotherapy	2	
Observation	1	
<u>Rating Scale:</u> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate		

Clinical Condition:**Local Excision in Early Stage Rectal Cancer****Variant 2:**

65-year-old otherwise healthy female with preoperative stage uT2N0, moderately differentiated adenocarcinoma. Tumor is 3 cm in diameter, freely mobile, and is located 4 cm from anal verge. No lymphovascular space invasion is noted.

Treatment	Rating	Comments
Surgery		
LAR or APR	9	
Local excision	2	
If Local Excision, then		
RT + chemotherapy	8	
RT alone	2	
Observation	1	
If RT + Chemo: RT Dose to Primary		
45 Gy/1.8 Gy	2	
50.4 Gy/1.8 Gy	8	
54 Gy/1.8 Gy	8	If small bowel can be excluded.
59.4 Gy/1.8 Gy	2	
Simulation		
Patient prone	9	
Small-bowel contrast at simulation	9	
Patient immobilized	9	
Use belly board	9	If patient is prone.
Anal marker	9	
Bladder full at simulation	8	
Patient supine	5	
If RT + Chemo: RT Volume		
L5/S1 pelvis to include perineum	9	In some cases where the lesion is 4 cm above the anal verge, the perineum may be spared as long as 3 cm of inferior margin can be maintained.
RT Technique		
3 field with photons	8	
4 field with photons	8	
AP/PA	2	
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate		

Clinical Condition:**Local Excision in Early Stage Rectal Cancer****Variant 3:****60-year-old female with uT3Nx adenocarcinoma, located 4 cm from anal verge.**

Treatment	Rating	Comments
Neoadjuvant RT + chemotherapy	9	See the ACR Appropriateness Criteria® topic on “ Resectable Rectal Cancer .”
LAR or APR	9	
Local excision	1	
<u>Rating Scale:</u> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate		