

American College of Radiology ACR Appropriateness Criteria®

Clinical Condition: Acute Respiratory Illness in Immunocompetent Patients

Variant 1: Older than age 40.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	8		☼
CT chest without contrast	4		☼ ☼ ☼
CT chest with contrast	3		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 2: Dementia, any age.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	8		☼
CT chest without contrast	6	In patients without reliable follow-up or with a likelihood of morbidity if disease is not detected initially, a CT may be required in the setting of a negative chest x-ray.	☼ ☼ ☼
CT chest with contrast	3		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 3: Younger than age 40, negative physical examination, and no other signs, symptoms, or risk factors.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	4		☼
CT chest without contrast	1		☼ ☼ ☼
CT chest with contrast	1		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition:**Acute Respiratory Illness in Immunocompetent Patients****Variant 4:****Younger than age 40 and positive physical examination or other risk factors.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9		☼
CT chest without contrast	4		☼ ☼ ☼
CT chest with contrast	3		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 5:**Complicated pneumonia.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9		☼
CT chest without contrast	8	If pneumonia is not resolving or intervention is contemplated.	☼ ☼ ☼
CT chest with contrast	5		☼ ☼ ☼
CT chest without and with contrast	2		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 6:**Suspected SARS.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9		☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 7:**Suspected SARS. Chest radiograph normal or equivocal.**

Radiologic Procedure	Rating	Comments	RRL*
CT chest without contrast	9		☼ ☼ ☼
CT chest with contrast	3		☼ ☼ ☼
CT chest without and with contrast	3		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 8:**Suspected anthrax.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9		☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition:

Acute Respiratory Illness in Immunocompetent Patients

Variant 9:

Suspected anthrax. Chest radiograph normal or equivocal.

Radiologic Procedure	Rating	Comments	RRL*
CT chest without contrast	8		☼ ☼ ☼
CT chest with contrast	5		☼ ☼ ☼
CT chest without and with contrast	3		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 10:

Acute asthma, uncomplicated.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	4		☼
CT chest without contrast	1		☼ ☼ ☼
CT chest with contrast	1		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 11:

Acute asthma and suspected pneumonia or pneumothorax.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9		☼
CT chest without contrast	2		☼ ☼ ☼
CT chest with contrast	1		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 12:

Acute exacerbation of COPD, “uncomplicated” (no history of coronary artery disease or congestive heart failure, no leukocytosis, fever, or chest pain).

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	4		☼
CT chest without contrast	2		☼ ☼ ☼
CT chest with contrast	1		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition:

Acute Respiratory Illness in Immunocompetent Patients

Variant 13:

Acute exacerbation of COPD with one or more of the following: leukocytosis, pain, history of coronary artery disease or congestive heart failure.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9		☼
CT chest without contrast	4		☼ ☼ ☼
CT chest with contrast	3		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 14:

Suspected H1N1.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9		☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 15:

Suspected H1N1. Chest radiograph normal or equivocal.

Radiologic Procedure	Rating	Comments	RRL*
CT chest without contrast	8	Chest x-rays may be normal in a large percentage of patients. If clinical suspicion is present, a chest CT should follow to better assess.	☼ ☼ ☼
CT chest with contrast	3		☼ ☼ ☼
CT chest without and with contrast	1		☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

ACUTE RESPIRATORY ILLNESS IN IMMUNOCOMPETENT PATIENTS

Expert Panel on Thoracic Imaging: Jacobo Kirsch, MD¹; José Ramirez, MD²; Tan-Lucien H. Mohammed, MD³; Judith K. Amorosa, MD⁴; Kathleen Brown, MD⁵; Debra Sue Dyer, MD⁶; Mark E. Ginsburg, MD⁷; Darel E. Heitkamp, MD⁸; Jean Jeudy, MD⁹; Heber MacMahon, MB, BCh¹⁰; James G. Ravenel, MD¹¹; Anthony G. Saleh, MD¹²; Rakesh D. Shah, MD.¹³

Summary of Literature Review

Introduction/Background

Acute respiratory illness (ARI) is defined as one or more of the following: cough, sputum production, chest pain, or dyspnea (with or without fever). The workup of a patient with ARI, including the need for chest radiography and computed tomography (CT), depends on many factors, including severity of the illness; age of patient; presence of fever, leukocytosis, or hypoxemia; clinical history; presence of other risk factors; and physical examination. Not all studies concur as to which patients with ARI should have chest radiographs.

Benacerraf et al [1] in a study of 1,102 outpatients with ARI, found patient age, the physical examination, and the presence or absence of hemoptysis to be important factors. Only 4% (7/175) of patients younger than age 40 with symptoms of ARI, a negative physical examination, and no hemoptysis had acute significant radiographic findings, whereas patients either older than age 40 with hemoptysis or with a positive physical examination, such as the presence of wheezing, had a much higher incidence of chest radiograph abnormalities. In a study of 464 patients with ARI, Heckerling [2] also found a low incidence (3%) of pneumonia in patients with negative physical examinations. A notable exception was found for patients with dementia, in whom the incidence of pneumonia was very high regardless of the results of the physical examination. Okimoto et al [3] studied 79 outpatients presenting with clinical suspicion of pneumonia and concluded that radiographs should be ordered only when patients present with fever, cough, sputum production, and coarse crackles on physical

examination. Conversely, Butcher et al [4] in a study of 221 patients with ARI found that 77 (35%) had new clinically important findings. Furthermore, the clinical findings did not differ significantly between those with positive radiographic findings and those with negative findings (ie, clinical history and physical examination were poor predictors of radiography-detected abnormalities). Speets et al [5] evaluated 192 patients with a clinical suspicion of pneumonia by general practitioners and found that the probability of pneumonia was changed by chest radiographic results in 53% of patients, with a decrease in probability in 47% and an increase in probability in 6%.

Pneumonia

In a series of 300 patients with acute cough illness, Aagaard et al [6] found that for patients with a high pretest probability of pneumonia, a radiograph was not always obtained in clinical practice; they infer that when the clinical probability of pneumonia exceeds a certain level, a negative radiograph would not alter treatment decisions by clinicians. A series by Basi et al [7] that included 2,706 patients hospitalized with community-acquired pneumonia (CAP) similarly showed that 911 (one-third) of them had radiographs initially interpreted as negative for pneumonia, with minimal change in this interpretation on retrospective review of a random subgroup. The groups with positive and negative radiographs had similar rates of positive sputum cultures and blood cultures. These two studies call into question the utility of radiographs in patients with high pretest probability of pneumonia. O'Brien et al [8] developed a prediction rule for the use of chest radiographs in evaluating for CAP and in a 70-patient series, in both the outpatient and emergency setting. They concluded that chest radiographs are unnecessary in patients with normal vital signs and physical examination findings. However, because approximately 5% of cases would be missed, their criteria are only useful for patients with reliable follow-up and a low likelihood of morbidity if CAP is not detected initially. Hagaman et al [9] showed that 21% of 105 patients hospitalized with a clinical diagnosis of CAP initially had negative chest radiographs, with over one-half of the 9 patients who had subsequent radiographic studies within 48 hours of hospitalization developing opacities.

Patients with substance abuse have an increased risk of ARI due to two potential conditions: respiratory pump failure and pulmonary pathology [10]. Respiratory pump failure generally does not have radiographic manifestations. However pulmonary pathology includes multiple diagnosis with chest radiographic manifestations, including aspiration, pulmonary edema, pneumonia, hemorrhage, and septic emboli.

Jochelson et al [11] found a low incidence (4%) of pneumonia in febrile, but otherwise asymptomatic, neutropenic patients with a normal physical examination.

¹Principal Author, Cleveland Clinic, Weston, Florida.

²Research Author, Cleveland Clinic, Weston, Florida.

³Panel Chair, Cleveland Clinic Foundation, Cleveland, Ohio.

⁴Robert Wood Johnson Medical School, New Brunswick, New Jersey.

⁵David Geffen School of Medicine at UCLA, Los Angeles, California.

⁶National Jewish Health, Denver, Colorado.

⁷Columbia University, New York, New York, Society of Thoracic Surgeons.

⁸Indiana University, Indianapolis, Indiana.

⁹University of Maryland Medical Center, Baltimore, Maryland.

¹⁰University of Chicago Hospital, Chicago, Illinois.

¹¹Medical University of South Carolina, Charleston, South Carolina.

¹²New York Methodist Hospital, Brooklyn, New York, The American College of Chest Physicians.

¹³North Shore University Hospital, Manhasset, New York.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

Navigante et al [12] found a similarly low incidence (2.3%) of pneumonia at chest radiography in febrile neutropenic patients without clinical suspicion of pneumonia from history or physical examination. Heussel et al [13] evaluated the utility of thin-section CT in a group of febrile neutropenic patients with normal or nonspecific chest radiographs. There were 146 episodes in 87 patients. Among the 14% with nonspecific chest radiographs, CT suggested pneumonia in all. Forty-eight percent had a normal chest radiograph, but CT findings were suggestive of pneumonia. Of these, a specific pathogen was identified in 43%. Both chest radiograph and CT findings were normal in 38%. The CT findings changed patient's therapy in 18%. Maertens et al [14] have proposed an algorithm employing CT in conjunction with galactomannan assays to select patients for high-dose antifungal therapy, and they performed a feasibility study assessing patients with 117 episodes of neutropenic fever. When the algorithm employing CT was used, only 4.4% of patients received antifungal therapy. Hachem et al [15] retrospectively analyzed CT scans and autopsy results in 96 cancer patients who died of pneumonia and found that the presence of nodules in neutropenic patients and cavitory lesions in nonneutropenic patients was highly associated with fungal infection.

According to the guidelines of the Infectious Diseases Society of America and the American Thoracic Society, chest radiography should be obtained whenever pneumonia is suspected in adults to establish the diagnosis and to aid in differentiating CAP from other common causes of cough and fever, such as acute bronchitis [16]. Findings on chest radiographs are one of several parameters used to determine: 1) which patients should be hospitalized (presence of pleural effusion); 2) which patients should be classified as having severe pneumonia (multilobar involvement); and 3) which patients may require additional diagnostic testing (cavitation, pleural effusion), including thoracentesis (pleural effusions >5 cm on lateral upright radiograph) [16]. CT may show findings in patients with normal radiographs, but the significance of these findings and therefore the utility of CT in patients with clinically suspected pneumonia and negative radiographs is unclear [16]. Hayden et al [17] found that in 27% of patients (out of 97) presenting to the emergency department with both a chest radiograph study and a CT scan performed in the workup of varied chief complaints, pneumonia was demonstrated on a CT scan in the face of a negative or nondiagnostic chest radiograph study. As in the study by O'Brien mentioned above, they recommend that when the chief complaint is fever, cough or chest pain in patients with serious comorbidities, obtaining a CT scan in the setting of a negative chest radiograph study would seem to be necessary [8,17].

Additionally, CT may play a role in the management of severe pneumonia. It can serve as a guide for pleural drainage or to localize an appropriate site for biopsy [18]. However, its use for potentially differentiating bacterial and nonbacterial pneumonias remains limited [19]. Severe

pneumonias bear a strong relationship to etiologic pathogens and have implications for antimicrobial treatment. Patients with severe pneumonia should be considered as candidates for admission to an intensive care unit.

Asthma

The need for chest radiographs in adult patients with acute asthma is controversial. Petheram et al [20] found clinically important (ie, affecting patient management) radiographic findings in 9% of their patients and concluded that chest radiography is indicated. However, Findley and Sahn [21] observed that 99% of their patients either had normal chest radiographic examinations or showed only slightly prominent markings or hyperinflation. Heckerling [2] reported that patients with acute asthma rarely have pneumonia. Findley and Sahn [21] recommended chest radiographs only when pneumonia or pneumothorax is suspected. White et al [22] found significant chest radiographic abnormalities in 34% of adults whose asthma exacerbation warranted admission to the hospital.

Chronic Obstructive Pulmonary Disease

Sherman et al [23] studied the utility of chest radiography in 242 patients with acute exacerbations of chronic obstructive pulmonary disease (COPD) (ie, dyspnea). Of this group, 135 patients (56%) had asthma, and 107 (44%) had emphysema and chronic bronchitis. Chest radiographs were abnormal in 14% but resulted in significant change in management in only 4.5%. They concluded that the chest radiograph is indicated only if the worsening dyspnea is accompanied by leukocytosis, chest pain, or edema or by a history of coronary artery disease or congestive heart failure (CHF).

Emerging infections and biological warfare agents have come to recent attention as causes of ARI. Two infections that received a great deal of attention recently, but which have subsequently become less active concerns, are severe acute respiratory syndrome (SARS) and anthrax.

SARS

SARS emerged in China in late 2002. The etiologic agent is a novel coronavirus (SARS-CoV) that appears to have originated in Himalayan palm civets and crossed the species barrier. In February 2003 the Program for Monitoring Emerging Diseases identified this novel presentation of pneumonia which because of air travel, rapidly spread across continents to involve patients in at least 27 countries. There is literature supporting the utility of chest radiography in patients with known or suspected SARS. Wong et al [24] and Paul et al [25] described the chest radiographic findings of SARS during the Hong Kong and Toronto epidemics. Chest radiographs were abnormal in 78%-80% of patients at presentation. The most common chest radiographic finding was unifocal opacity with a peripheral and basilar predominance. Multifocal or diffuse opacities could be present initially or develop as the disease progressed. Patients whose disease progressed were generally older, had more comorbidities, and had a higher fatality rate. Cavitation, pleural effusion,

and lymphadenopathy were not features of SARS. Antonio et al [26] studied 1,373 patients in Hong Kong with SARS and found a sensitivity for disease of 82.4% on initial chest radiographs; they also concluded that the initial extent of radiographic opacification had prognostic value and that the rate of radiographic progression could be used as a prognostic indicator. However, a retrospective study by Chan et al [27] with a cohort of 1,311 patients who attended an Emergency Department SARS screening clinic during the 2003 outbreak in Hong Kong found that the sensitivity (50.3%) of the initial chest radiograph was much lower than previously expected. In contrast, the sensitivity of serial chest radiography in that study was significantly improved to 94.4% with a negative predictive value of serial chest radiography of 99.0%.

Thin-section chest CT findings of SARS have been described by Wong et al [28], Chan et al [29], and Paul et al [25]. The most common findings are ground-glass opacities and crazy paving. More extensive findings include focal or multifocal consolidation. Chan et al [29] described pleural effusions and pneumomediastinum developing in 26% of patients scanned during the course of their illness. Hui et al [30] suggest that high-resolution computed tomography (HRCT) is useful for early diagnosis of SARS in patients with negative chest radiographs. They studied 47 patients with suspected SARS and normal chest radiographs; 25 of 27 patients with serologic confirmation of SARS had abnormal findings at HRCT and developed clinical SARS, while the two with negative HRCT scans did not develop pulmonary infection.

Anthrax

Anthrax is endemic in the soil of Texas, Oklahoma, and the Mississippi Valley. During the 20th century a number of countries developed weapons-grade anthrax to be used as a biological warfare agent [31]. Much of modern medical experience with it arises from a Soviet military accident in 1979 in which 42 people died of anthrax, and from cases of anthrax that developed in the U.S. in 2001 as a result of biological warfare. Anthrax comes in three forms: cutaneous, gastrointestinal, and inhalational. Ninety-five percent of anthrax is cutaneous, but the inhalational form is the most deadly. Inhalation of anthrax spores leads to hemorrhagic lymphadenitis and mediastinitis, sometimes accompanied by necrotizing pneumonia. The chest radiographic findings [31-32] include widened mediastinum and hila, often accompanied by pleural effusions and parenchymal opacities. Earls et al [32] described the CT findings in 2 patients who survived inhalational anthrax. The CT characteristics were very suggestive of the diagnosis and included hyperattenuating hilar and mediastinal lymphadenopathy and hemorrhagic pleural effusion. Less specific findings included mediastinal edema, peribronchial thickening, and pleural effusion.

H1N1

In 2009, a new strain of human influenza A (H1N1) virus was identified in Mexico that grew to be declared a level 6 pandemic by the World Health Organization. The clinical symptoms in the greatest number of patients consist of mild clinical presentation with influenza-like illness. However, a small percentage of patients with H1N1 virus pneumonia may develop acute respiratory distress syndrome (ARDS) or severe cardiopulmonary failure despite the use of intensive-care treatment [33].

A review of chest radiograph findings in 97 patients with proven H1N1 infection by Aviram et al [34] described an abnormal chest radiograph in 40% of them. Ground-glass opacities, often central, were identified in most of the patients who had an abnormal chest radiograph upon admission to the emergency department. Overt consolidation, frequently seen in patchy distribution was also a common finding. Abbo et al [35] reviewed the imaging characteristics of 123 patients receiving radiographs and 21 patients receiving CT examinations, all with proven H1N1 influenza. In this study, up to 67% of initial radiographs were negative. When present, the most common chest radiograph finding was consolidation. On CT examinations, 90% of patients showed consolidation, ground-glass opacity, nodules, or a combination of these findings; 4 of the 21 patients imaged by both modalities had a negative chest radiograph with a positive CT. A study by Agarawal et al [36] in 222 patients with suspected or proven H1N1 influenza showed normal chest radiographs in over 58% of patients and progress to bilateral extensive air-space disease in severely ill patients. Additionally, in a study by McEwen et al [37] of 42 adults with proven H1N1, only 12 had abnormal chest radiographs. Interestingly, this study showed a trend (not validated due to the small number of patients) that those who required hospital admission had an abnormal chest radiograph appeared to be at greater risk of requiring intubation and ventilation than those with normal chest radiographs. Similarly, the study by Aviram et al [34] mentioned earlier suggests that the findings at initial chest radiography may have significance in the prediction of clinical outcome, with extensive involvement of both lungs being associated with an adverse prognosis.

Several small case series presenting the CT imaging findings of H1N1 infection show unilateral or bilateral ground-glass opacities with or without associated focal or multifocal areas of consolidation, both revealing a predominantly peribronchovascular and subpleural distribution [36,38-39].

Summary

- Based on these studies, the chest radiograph seems warranted in ARI when one or more of the following are present: older than age 40; dementia; a positive physical examination; hemoptysis; associated abnormalities (leukocytosis, hypoxemia); or other risk factors, including coronary artery disease, CHF, or drug-induced acute respiratory failure.

- Chest radiography also seems warranted for any adult patient with clinical suspicion of pneumonia, although some clinicians may choose not to perform radiography if clinical suspicion of respiratory infection is sufficiently high to warrant treatment if a radiograph were to be negative.
- It appears that in patients with ARI who are younger than age 40, chest radiography is not routinely indicated unless there are other abnormalities, a positive physical examination, or other risk factors.
- It also appears that chest radiographic examination is not indicated in most patients with exacerbations of COPD (including asthma) unless there is a suspected complication such as pneumonia or pneumothorax or unless one or more of the following are present: leukocytosis, chest pain, edema, or a history of coronary artery disease or CHF.
- Chest CT may be warranted in complicated cases of severe pneumonia and in febrile neutropenic patients with normal or nonspecific chest radiographic findings.
- In patients with a normal chest radiograph and a high clinical suspicion of either SARS or H1N1, CT can be helpful in suggesting the diagnosis.

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, both because of organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared to those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
0	0 mSv	0 mSv
☼	<0.1 mSv	<0.03 mSv
☼ ☼	0.1-1 mSv	0.03-0.3 mSv
☼ ☼ ☼	1-10 mSv	0.3-3 mSv
☼ ☼ ☼ ☼	10-30 mSv	3-10 mSv
☼ ☼ ☼ ☼ ☼	30-100 mSv	10-30 mSv

*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as NS (not specified).

Supporting Document(s)

- [ACR Appropriateness Criteria® Overview](#)
- [Procedure Information](#)
- [Evidence Table](#)

References

1. Benacerraf BR, McLoud TC, Rhea JT, Tritschler V, Libby P. An assessment of the contribution of chest radiography in outpatients with acute chest complaints: a prospective study. *Radiology* 1981; 138(2):293-299.
2. Heckerling PS. The need for chest roentgenograms in adults with acute respiratory illness. Clinical predictors. *Arch Intern Med* 1986; 146(7):1321-1324.
3. Okimoto N, Yamato K, Kurihara T, et al. Clinical predictors for the detection of community-acquired pneumonia in adults as a guide to ordering chest radiographs. *Respirology* 2006; 11(3):322-324.
4. Butcher BL, Nichol KL, Parenti CM. High yield of chest radiography in walk-in clinic patients with chest symptoms. *J Gen Intern Med* 1993; 8(3):115-119.
5. Speets AM, Hoes AW, van der Graaf Y, Kalmijn S, Sachs AP, Mali WP. Chest radiography and pneumonia in primary care: diagnostic yield and consequences for patient management. *Eur Respir J* 2006; 28(5):933-938.
6. Aagaard E, Maselli J, Gonzales R. Physician practice patterns: chest x-ray ordering for the evaluation of acute cough illness in adults. *Med Decis Making* 2006; 26(6):599-605.
7. Basi SK, Marrie TJ, Huang JQ, Majumdar SR. Patients admitted to hospital with suspected pneumonia and normal chest radiographs: epidemiology, microbiology, and outcomes. *Am J Med* 2004; 117(5):305-311.
8. O'Brien WT, Sr., Rohweder DA, Lattin GE, Jr., et al. Clinical indicators of radiographic findings in patients with suspected community-acquired pneumonia: who needs a chest x-ray? *J Am Coll Radiol* 2006; 3(9):703-706.
9. Hagaman JT, Rouan GW, Shipley RT, Panos RJ. Admission chest radiograph lacks sensitivity in the diagnosis of community-acquired pneumonia. *Am J Med Sci* 2009; 337(4):236-240.
10. Wilson KC, Saukkonen JJ. Acute respiratory failure from abused substances. *J Intensive Care Med* 2004; 19(4):183-193.
11. Jochelson MS, Altschuler J, Stomper PC. The yield of chest radiography in febrile and neutropenic patients. *Ann Intern Med* 1986; 105(5):708-709.
12. Navigante AH, Cerchietti LC, Costantini P, et al. Conventional chest radiography in the initial assessment of adult cancer patients with fever and neutropenia. *Cancer Control* 2002; 9(4):346-351.

13. Heussel CP, Kauczor HU, Heussel G, Fischer B, Mildenerger P, Thelen M. Early detection of pneumonia in febrile neutropenic patients: use of thin-section CT. *AJR* 1997; 169(5):1347-1353.
14. Maertens J, Theunissen K, Verhoef G, et al. Galactomannan and computed tomography-based preemptive antifungal therapy in neutropenic patients at high risk for invasive fungal infection: a prospective feasibility study. *Clin Infect Dis* 2005; 41(9):1242-1250.
15. Hachem R, Sumoza D, Hanna H, Girgawy E, Munsell M, Raad I. Clinical and radiologic predictors of invasive pulmonary aspergillosis in cancer patients: should the European Organization for Research and Treatment of Cancer/Mycosis Study Group (EORTC/MSG) criteria be revised? *Cancer* 2006; 106(7):1581-1586.
16. Mandell LA, Wunderink RG, Anzueto A, et al. Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis* 2007; 44 Suppl 2:S27-72.
17. Hayden GE, Wrenn KW. Chest radiograph vs. computed tomography scan in the evaluation for pneumonia. *J Emerg Med* 2009; 36(3):266-270.
18. Baber CE, Hedlund LW, Oddson TA, Putman CE. Differentiating empyemas and peripheral pulmonary abscesses: the value of computed tomography. *Radiology* 1980; 135(3):755-758.
19. Ito I, Ishida T, Togashi K, et al. Differentiation of bacterial and non-bacterial community-acquired pneumonia by thin-section computed tomography. *Eur J Radiol* 2009; 72(3):388-395.
20. Petheram IS, Kerr IH, Collins JV. Value of chest radiographs in severe acute asthma. *Clin Radiol* 1981; 32(3):281-282.
21. Findley LJ, Sahn SA. The value of chest roentgenograms in acute asthma in adults. *Chest* 1981; 80(5):535-536.
22. White CS, Cole RP, Lubetsky HW, Austin JH. Acute asthma. Admission chest radiography in hospitalized adult patients. *Chest* 1991; 100(1):14-16.
23. Sherman S, Skoney JA, Ravikrishnan KP. Routine chest radiographs in exacerbations of chronic obstructive pulmonary disease. Diagnostic value. *Arch Intern Med* 1989; 149(11):2493-2496.
24. Wong KT, Antonio GE, Hui DS, et al. Severe acute respiratory syndrome: radiographic appearances and pattern of progression in 138 patients. *Radiology* 2003; 228(2):401-406.
25. Paul NS, Roberts H, Butany J, et al. Radiologic pattern of disease in patients with severe acute respiratory syndrome: the Toronto experience. *Radiographics* 2004; 24(2):553-563.
26. Antonio GE, Ooi CG, Wong KT, et al. Radiographic-clinical correlation in severe acute respiratory syndrome: study of 1373 patients in Hong Kong. *Radiology* 2005; 237(3):1081-1090.
27. Chan SS, Mak PS, Shing KK, Chan PN, Ng WH, Rainer TH. Chest radiograph screening for severe acute respiratory syndrome in the ED. *Am J Emerg Med* 2005; 23(4):525-530.
28. Wong KT, Antonio GE, Hui DS, et al. Thin-section CT of severe acute respiratory syndrome: evaluation of 73 patients exposed to or with the disease. *Radiology* 2003; 228(2):395-400.
29. Chan MS, Chan IY, Fung KH, Poon E, Yam LY, Lau KY. High-resolution CT findings in patients with severe acute respiratory syndrome: a pattern-based approach. *AJR* 2004; 182(1):49-56.
30. Hui JY, Hon TY, Yang MK, et al. High-resolution computed tomography is useful for early diagnosis of severe acute respiratory syndrome-associated coronavirus pneumonia in patients with normal chest radiographs. *J Comput Assist Tomogr* 2004; 28(1):1-9.
31. Shafazand S, Doyle R, Ruoss S, Weinacker A, Raffin TA. Inhalational anthrax: epidemiology, diagnosis, and management. *Chest* 1999; 116(5):1369-1376.
32. Earls JP, Cerva D, Jr., Berman E, et al. Inhalational anthrax after bioterrorism exposure: spectrum of imaging findings in two surviving patients. *Radiology* 2002; 222(2):305-312.
33. Henzler T, Meyer M, Kalenka A, et al. Image findings of patients with H1N1 virus pneumonia and acute respiratory failure. *Acad Radiol* 2010; 17(6):681-685.
34. Aviram G, Bar-Shai A, Sosna J, et al. H1N1 influenza: initial chest radiographic findings in helping predict patient outcome. *Radiology* 2010; 255(1):252-259.
35. Abbo L, Quartin A, Morris MI, et al. Pulmonary imaging of pandemic influenza H1N1 infection: relationship between clinical presentation and disease burden on chest radiography and CT. *Br J Radiol* 2010; 83(992):645-651.
36. Agarwal PP, Cinti S, Kazerooni EA. Chest radiographic and CT findings in novel swine-origin influenza A (H1N1) virus (S-OIV) infection. *AJR* 2009; 193(6):1488-1493.
37. McEwen RE, Scriven JE, Green CA, Bailey MS, Banerjee AK. Chest radiography findings in adults with pandemic H1N1 2009 influenza. *Br J Radiol* 2010; 83(990):499-504.
38. Ajlan AM, Quiney B, Nicolaou S, Muller NL. Swine-origin influenza A (H1N1) viral infection: radiographic and CT findings. *AJR* 2009; 193(6):1494-1499.
39. Marchiori E, Zanetti G, Hochegger B, et al. High-resolution computed tomography findings from adult patients with Influenza A (H1N1) virus-associated pneumonia. *Eur J Radiol* 2010; 74(1):93-98.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.