

**American College of Radiology
ACR Appropriateness Criteria®**

Clinical Condition: Routine Chest Radiograph

Variant 1: Monitoring stable patient.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest portable routine admission	9	Not for coronary artery bypass graft. Not for acute upper gastrointestinal bleeding.	Min
X-ray chest portable clinical indications only	8	Clinical worsening only.	Min
X-ray chest portable routine monitoring	2		Min
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 2: Respiratory failure. Patient receiving mechanical ventilation.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest portable daily	9		Min
X-ray chest portable clinical indications only	9		Min
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 3: Compromised respiratory function. Patient with endotracheal tubes.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest portable after catheter/tube insertion	9		Min
X-ray chest portable clinical indications only	9		Min
X-ray chest portable follow-up catheter/tube	2		Min
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 4: Central venous pressure catheter (CVP) insertion.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest portable after catheter/tube insertion	9		Min
X-ray chest portable clinical indications only	8		Min
X-ray chest portable follow-up catheter/tube	2		Min
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Routine Chest Radiograph****Variant 5:****Cardiopulmonary compromise. Swan-Ganz catheter insertion.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest portable after catheter/tube insertion	9		Min
X-ray chest portable clinical indications only	8		Min
X-ray chest portable follow-up catheter/tube	2		Min
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 6:**Potential cardiopulmonary compromise. Nasogastric (NG) tube insertion.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest portable after catheter/tube insertion	9	Feeding NG tube.	Min
X-ray chest portable after catheter/tube insertion	6	Nonfeeding NG tube.	Min
X-ray chest portable follow-up catheter/tube	2		Min
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

Variant 7:**Respiratory compromise. Chest tube insertion.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest portable after catheter/tube insertion	8		Min
X-ray chest portable clinical indications only	8		Min
X-ray chest portable follow-up catheter/tube	2		Min
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

ROUTINE CHEST RADIOGRAPH

Expert Panel on Thoracic Imaging: Suzanne L. Aquino, MD¹; Arfa Khan, MD²; Poonam V. Batra MD³; Jud W. Gurney, MD⁴; Linda B. Haramati, MD⁵; Heber MacMahon, MD⁶; Tan-Lucien H. Mohammed, MD⁷; Anna Rozenshtein, MD⁸; Kay H. Vydareny, MD⁹; Lacey Washington, MD¹⁰; Helen T. Winer-Muram, MD¹¹; Pamela K. Woodard, MD¹²; Larry Kaiser, MD¹³; Suhail Raof, MBBS.¹⁴

Summary of Literature Review

Portable Radiographs in the ICU Setting

Portable radiographs can be categorized as:

1. daily or routine radiographs for patient monitoring;
2. radiographs obtained after specific procedures; and
3. radiographs documenting the presence or course of disease.

This narrative concerns the daily routine radiograph only and does not include postprocedural or postsurgical radiographs. Studies that included large volumes of chest radiographs (CXR) (>200) were included in this discussion. The most recent two studies evaluated the clinically relevant use of daily routine versus non-routine clinically indicated chest radiographs. A large study by Graat et al [1] prospectively evaluated the clinical value of 2,457 routine CXRs in a combined surgical/medical intensive care unit (SICU/MICU). In their study, 5.8% of routine CXRs showed new or unexpected findings; but only 2.2% warranted a change in therapy. No difference was found between the medical and surgical patients. A randomized control study of MICU patients by Krivopal et al [2], prospectively divided MICU patients into those who received routine CXRs and those who only received non-routine (clinically indicated) CXRs. They found a greater percentage of radiographs with significant findings (requiring intervention) in the non-routine group (26.5%) than in the routine group (13.3%). Significant interventions included diuresis, antibiotic administration, or an invasive procedure. The non-routine group also received significantly less radiographs per person than the routine group (4.4 vs 6.8). There was no significant difference in outcome between the groups in length of intubation, ICU stay, hospital stay or mortality.

Older studies reviewed only daily radiographs in ICU patients without comparison to the outcome of those without routine radiographs. The yield in routine radiographs was overall higher, especially in MICU patients and especially those who were intubated. If one looks at a broader definition of radiographic value such as “important or unexpected findings,” the range in results was large, from 3% to 60%. Studies by Brainsky et al [3] and Marik et al [4] reported 20% and 37% changes in management, respectively from routine portable CXRs obtained in the MICU. Patients who were intubated were the most affected. Hall et al [5] reported the lowest rate of significant abnormal CXR findings at 3% of all radiographs in 18% of the MICU patients. They still recommended routine studies on all critically ill patients. In a study by Strain et al [6] although a high yield was found in MICU patients who had acute cardiopulmonary disease, the yield was very low in patients with stable cardiac disease (usually myocardial infarction) and in patients in the ICU who had extrathoracic disease only.

For cardiothoracic ICU patients, two prospective nonrandomized studies showed a low incidence of significant findings on routine daily radiographs (4.5% in both studies) resulting in minimal impact on patient management. The results support the recommendation to obtain CXRs in cardiothoracic ICU for clinical findings and not for routine follow-up [7,8].

Recommendation

Routine daily portable radiographs are indicated for patients with acute cardiopulmonary problems. In stable patients admitted for cardiac monitoring, or in stable patients admitted for extrathoracic disease only, an initial admission radiograph is recommended, with follow-up radiographs obtained only for specific clinical indications.

Endotracheal Tubes

There are nine studies described in the literature since 1980 [4,6,7,9-14] that evaluate the significance of the chest radiograph in assessing endotracheal tube placement following insertion. In five of them, between 12% and 15% of patients had malpositioned endotracheal tubes, many of which required repositioning. Two studies found 28% and 46% of tubes malpositioned upon insertion, and the single dissenting paper found 2% malpositioned. Two studies compared x-ray with physical exam [9,15]. In both studies, physical exam predicted malpositioned tubes in 3% of patients, whereas the x-rays showed malpositioning in 14% of patients in one study and 28% in the other. Kollef et al [16] found that the vast majority of malpositioned tubes were discovered in the first three days.

¹Review Author, Massachusetts General Hospital, Boston, Mass; ²Panel Chair, Long Island Jewish Medical Center, New Hyde Park, NY; ³David Geffen School of Medicine, Los Angeles, Calif; ⁴University of Nebraska, Omaha, Neb; ⁵Albert Einstein College of Medicine, Montefiore Medical Center, Bronx, NY; ⁶University of Chicago Hospital, Chicago, Ill; ⁷Cleveland Clinic Foundation, Cleveland, Ohio; ⁸Columbia Presbyterian Medical Center, New York, NY; ⁹Emory University Hospital, Atlanta, Ga; ¹⁰Froedtert Memorial Lutheran Hospital, Milwaukee, Wis; ¹¹Richard Roudebush Veterans Medical Center, Indianapolis, Ind; ¹²Mallinckrodt Institute of Radiology, Saint Louis, Mo; ¹³University of Pennsylvania Medical Center, Philadelphia, Pa, The Society of Thoracic Surgeons; ¹⁴New York Methodist Hospital, Brooklyn, NY, The American College of Chest Physicians.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Recommendation

Very few malpositioned tubes are detected by physical examination. Routine radiographs immediately postintubation are indicated to insure proper positioning.

CVP Catheters

Eight studies were reviewed regarding central venous pressure (CVP) catheters [6,9-14,16]. The vast majority came to the same conclusion. Approximately 10% of the CXRs demonstrated malpositioned catheters. Pneumothoraces were present in only a small percentage of patients. Gray et al [9] separated jugular and subclavian catheters. Complications were twice as common with subclavian catheters (17% vs 8%), although unsuspected complications were infrequent.

Recommendation

A CXR after insertion of a CVP catheter is recommended to demonstrate proper placement and detect any complications. Beyond the initial insertion, follow-up radiographs have a low yield for revealing complications. Follow-up radiographs are suggested only when complications are suspected clinically.

Swanz-Ganz Catheters

Previously mentioned studies incorporated the position and potential complications of Swan-Ganz catheter placements on CXRs obtained immediately post-procedure. The majority of complications, which occur in approximately 10% of catheter insertions, are minor and require catheter repositioning [9,10,12,17]. Pneumothorax rate was approximately 2% [10,17].

Recommendation

Portable radiography is suggested after catheter insertion. Once pneumothorax has been excluded and proper positioning has been assured, follow-up radiographs are not required except for specific clinical indications.

Nasogastric Tubes

There are no large prospective studies that consider the utility of obtaining CXR immediately after the insertion of a nasogastric suction tube or a small-bore feeding tube. Routine radiographs revealed important tube malpositioning in 1% of cases [6,10,12]. Other complications have also been reported [18-21]. Clearly, a patient with a functioning nasogastric tube that has already been documented to be in satisfactory position needs no imaging unless a clinical problem arises.

Recommendation

Based on limited evidence, small-bore feeding tubes may, in a small but significant number of patients, be inadvertently placed in the lungs. This error is not always detected clinically and may lead to injection of feeding material into the lung, or tube penetration of the pleura, with subsequent pneumothorax. A CXR is warranted after initial nasogastric tube insertion and before the first

feeding. Beyond the initial radiograph, follow-up radiographs are not required for management of stable tubes.

Chest Tubes

Few studies have been performed to evaluate the efficacy of the initial CXR after the insertion of a chest tube. The three available studies show that approximately 10% of tubes are malpositioned [6,11,14]. Many of the x-ray abnormalities detected are minor and do not result in changes of tube positions.

Recommendation

After insertion of a chest tube, a CXR is recommended to show the position of the tube, any success in drainage, and possible complications from insertion. Beyond this point, evaluation of tube position and function is warranted based on management of the pleural space and clinical indications.

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations	
Relative Radiation Level	Effective Dose Estimate Range
None	0
Minimal	< 0.1 mSv
Low	0.1-1 mSv
Medium	1-10 mSv
High	10-100 mSv

References

1. Graat ME, Hendrikse KA, Spronk PE, Korevaar JC, Stoker J, Schultz MJ. Chest radiography practice in critically ill patients: a postal survey in the Netherlands. *BMC Med Imaging* 2006; 6:8.
2. Krivopal M, Shlobin OA, Schwartzstein RM. Utility of daily routine portable chest radiographs in mechanically ventilated patients in the medical ICU. *Chest* 2003; 123(5):1607-1614.
3. Brainsky A, Fletcher RH, Glick HA, Lanken PN, Williams SV, Kundel HL. Routine portable chest radiographs in the medical intensive care unit: effects and costs. *Crit Care Med* 1997; 25(5):801-805.
4. Marik PE, Janower ML. The impact of routine chest radiography on ICU management decisions: an observational study. *Am J Crit Care* 1997; 6(2):95-98.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

5. Hall JB, White SR, Karrison T. Efficacy of daily routine chest radiographs in intubated, mechanically ventilated patients. *Crit Care Med* 1991; 19(5):689-693.
6. Strain DS, Kinasewitz GT, Vereen LE, George RB. Value of routine daily chest x-rays in the medical intensive care unit. *Crit Care Med* 1985; 13(7):534-536.
7. O'Brien W, Karski JM, Cheng D, Carroll-Munro J, Peniston C, Sandler A. Routine chest roentgenography on admission to intensive care unit after heart operations: is it of any value? *J Thorac Cardiovasc Surg* 1997; 113(1):130-133.
8. Graham RJ, Meziane MA, Rice TW, et al. Postoperative portable chest radiographs: optimum use in thoracic surgery. *J Thorac Cardiovasc Surg* 1998; 115(1):45-50; discussion 50-42.
9. Gray P, Sullivan G, Ostryzniuk P, McEwen TA, Rigby M, Roberts DE. Value of postprocedural chest radiographs in the adult intensive care unit. *Crit Care Med* 1992; 20(11):1513-1518.
10. Silverstein DS, Livingston DH, Elcavage J, Kovar L, Kelly KM. The utility of routine daily chest radiography in the surgical intensive care unit. *J Trauma* 1993; 35(4):643-646.
11. Henschke CI, Pasternack GS, Schroeder S, Hart KK, Herman PG. Bedside chest radiography: diagnostic efficacy. *Radiology* 1983; 149(1):23-26.
12. Horst HM, Fagan B, Beute GH. Chest radiographs in surgical intensive care patients: a valuable "routine". *Henry Ford Hosp Med J* 1986; 34(2):84-86.
13. Brunel W, Coleman DL, Schwartz DE, Peper E, Cohen NH. Assessment of routine chest roentgenograms and the physical examination to confirm endotracheal tube position. *Chest* 1989; 96(5):1043-1045.
14. Bekemeyer WB, Crapo RO, Calhoun S, Cannon CY, Clayton PD. Efficacy of chest radiography in a respiratory intensive care unit. A prospective study. *Chest* 1985; 88(5):691-696.
15. Bhagwanjee S, Muckart DJ. Routine daily chest radiography is not indicated for ventilated patients in a surgical ICU. *Intensive Care Med* 1996; 22(12):1335-1338.
16. Kollef MH, Legare EJ, Damiano M. Endotracheal tube misplacement: incidence, risk factors, and impact of a quality improvement program. *South Med J* 1994; 87(2):248-254.
17. Sise MJ, Hollingsworth P, Brimm JE, Peters RM, Virgilio RW, Shackford SR. Complications of the flow-directed pulmonary artery catheter: A prospective analysis in 219 patients. *Crit Care Med* 1981; 9(4):315-318.
18. Kassner EG, Baumstark A, Balsam D, Haller JO. Passage of feeding catheters into the pleural space: a radiographic sign of trauma to the pharynx and esophagus in the newborn. *AJR* 1977; 128(1):19-22.
19. Hand RW, Kempster M, Levy JH, Rogol PR, Spirn P. Inadvertent transbronchial insertion of narrow-bore feeding tubes into the pleural space. *JAMA* 1984; 251(18):2396-2397.
20. Balogh GJ, Adler SJ, VanderWoude J, Glazer HS, Roper C, Weyman PJ. Pneumothorax as a complication of feeding tube placement. *AJR* 1983; 141(6):1275-1277.
21. Dorsey JS, Cogordan J. Nasotracheal intubation and pulmonary parenchymal perforation. An unusual complication of naso-enteral feeding with small-diameter feeding tubes. *Chest* 1985; 87(1):131-132.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.