

**American College of Radiology
ACR Appropriateness Criteria®**

Clinical Condition: Screening for Pulmonary Metastases

Variant 1: Primary malignancy: bone and soft-tissue sarcoma.

Radiologic Procedure	Rating	Comments	RRL*
CT chest without contrast	9	Initial evaluation or surveillance.	☼ ☼ ☼
X-ray chest	9	If performed as a baseline.	☼
FDG-PET whole body	5		☼ ☼ ☼ ☼
MRI chest with or without contrast	2		O
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 2: Primary malignancy: renal cell carcinoma.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	8		☼
CT chest without contrast	7	Depends on the stage of the disease.	☼ ☼ ☼
MRI chest with or without contrast	2		O
FDG-PET whole body	1		☼ ☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 3: Primary malignancy: testicular cancer.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	8		☼
CT chest without contrast	7	Recommended if abdominal disease is present.	☼ ☼ ☼
MRI chest with or without contrast	3		O
FDG-PET whole body	3		☼ ☼ ☼ ☼
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Clinical Condition:**Screening for Pulmonary Metastases****Variant 4:****Primary malignancy: malignant melanoma.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9	If performed as a baseline.	☼
CT chest without contrast	8	Initial evaluation or surveillance.	☼ ☼ ☼
FDG-PET whole body	5		☼ ☼ ☼ ☼
MRI chest with or without contrast	2		○
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Variant 5:**Primary malignancy: head and neck carcinoma.**

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	9	If performed as a baseline.	☼
CT chest without contrast	9	Initial evaluation or surveillance.	☼ ☼ ☼
FDG-PET whole body	5		☼ ☼ ☼ ☼
MRI chest with or without contrast	2		○
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

SCREENING FOR PULMONARY METASTASES

Expert Panel on Thoracic Imaging: Tan-Lucien Mohammed, MD¹; Aqeel A. Chowdhry, MD²; Arfa R. Khan, MD³; Judith K. Amorosa, MD⁴; Poonam V. Batra, MD⁵; Debra Sue Dyer, MD⁶; Jud W. Gurney, MD⁷; Jean Jeudy, MD⁸; Larry Kaiser, MD⁹; Heber MacMahon, MB, BCh¹⁰; Suhail Raof, MBBS¹¹; Kay H. Vydaireny, MD.¹²

Summary of Literature Review

The incidence of pulmonary metastatic disease in patients who have died of an extrathoracic malignancy (ETM) is reported as 20%-54% [1,2]. The indications for chest radiography, computed tomography (CT), magnetic resonance imaging (MRI), scintigraphic imaging, and positron emission tomography/computed tomography (PET/CT) have been discussed in the literature. There have been improvements in CT imaging quality and scan time, as well as advances in the field of nuclear medicine and MRI. In particular, there has been widespread use of PET/CT for evaluating patients with metastatic pulmonary disease, especially in patients with primary head and neck tumors.

In determining the specific imaging modality that should be performed, authors conclude that several factors should be taken into consideration: 1) the biological behavior of the tumor, 2) the sensitivity and specificity of the imaging modality, 3) radiation dose, and 4) cost-effectiveness. The relative indications for chest radiography, CT, MRI, and scintigraphy have been evaluated for various primary malignancies. Detection of pulmonary nodules, lymphangitic spread, endobronchial lesions, intravascular metastatic pulmonary disease, nodal disease, and chest wall lesions have all been discussed in the literature.

Chest Radiography

It is generally accepted that chest radiography, with posteroanterior (PA) and lateral views, should be the initial imaging test in patients without known or suspected thoracic metastatic disease [1-3]. If the chest radiograph demonstrates obvious multiple pulmonary nodules,

further imaging beyond follow-up chest radiography may not be indicated unless biopsy is planned, or unless precise quantification of disease is required in the preoperative evaluation for metastasectomy or the assessment of response to systemic radiation therapy or chemotherapy.

Some authors have questioned the role of routine chest radiographs. In one study, a review of routine chest radiographs obtained in the evaluation of patients with breast cancer revealed that fewer than 0.93% of these radiographs demonstrated previously undiagnosed pulmonary metastases [4]. In another study, 876 asymptomatic patients with localized cutaneous (stage I or intermediate thickness stage II) malignant melanoma had initial staging chest radiographs; 130 (15%) had "suspicious" findings, but on further follow-up only one (0.1%) of these patients had a true-positive study for pulmonary metastasis [5]. Another study analyzed the overall cost-effectiveness of chest radiographs in the life-long screening of patients with intermediate-thickness cutaneous melanoma. It was concluded that significant cost savings may be possible by decreasing the frequency of screening in the first 2 years and limiting screening to the first 5 to 10 years after diagnosis [6]. Yet another study involving 23 patients with untreated primary head and neck tumors demonstrated that chest radiography by itself had a lower sensitivity of detecting pulmonary metastatic disease (67%) compared to PET/CT (100%) [7]. Patients with a higher probability of pulmonary metastatic disease should be screened more frequently or with a more sensitive imaging modality such as CT.

Computed Tomography

Compared with chest radiography, CT is much more sensitive for detecting pulmonary nodules because of its lack of superimposition and its high contrast resolution [1-3]. Other abnormalities, such as lymphadenopathy, pleural involvement, chest wall lesions, endobronchial lesions, intravascular pulmonary involvement, or incidental findings in the upper abdomen, may also be revealed or better demonstrated. In patients with known ETM, chest CT is recommended if the initial chest radiograph reveals an apparent solitary pulmonary nodule or an equivocal finding. If the chest radiograph is negative, CT is recommended if the underlying ETM is one that has a high propensity for dissemination to the lungs, such as breast, renal cell, colon, and bladder carcinoma. As noted in the preceding section, CT is indicated even with multiple pulmonary nodules on the chest radiograph if biopsy or definitive treatment by metastasectomy or systemic therapy is planned.

It is now well known that spiral CT scanning is more sensitive than conventional CT, allowing the detection of a significantly larger number of nodules and also a larger number of small nodules <5 mm in diameter [8]. With further developments in technology, it is likely that the sensitivity of CT scanning will continue to improve while

¹Principal Author and Panel Vice-chair, Cleveland Clinic Foundation, Cleveland, Ohio.

²Research Author, Cleveland Clinic Foundation, Cleveland, Ohio.

³Panel Chair, Long Island Jewish Medical Center, New Hyde Park, New York.

⁴Robert Wood Johnson Medical School, New Brunswick, New Jersey.

⁵David Geffen School of Medicine, Los Angeles, California.

⁶National Jewish Health, Denver, Colorado.

⁷University of Nebraska, Omaha, Nebraska.

⁸University of Maryland Medical Center, Baltimore, Maryland.

⁹University of Pennsylvania Medical Center, Philadelphia, Pennsylvania, Society of Thoracic Surgeons.

¹⁰University of Chicago Hospital, Chicago, Illinois.

¹¹New York Methodist Hospital, Brooklyn, New York, American College of Chest Physicians.

¹²Emory University School of Medicine, Atlanta, Georgia.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply society endorsement of the final document.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

the radiation dose associated with scanning may be lowered. Nevertheless, a few studies that have correlated CT findings with surgical or pathologic findings offer some sobering results. In a retrospective review, McCormack et al [9] found that CT underestimated the surgical pathologic findings in 25% of cases. More thorough detection of metastatic nodules is possible at thoracotomy by means of manual palpation of the entire collapsed lung [10,11].

It has been suggested that the greater sensitivity of CT for detecting pulmonary nodules, as compared with chest radiography, is associated with diminished specificity [2]. Nevertheless, it is increasingly recognized that even small pulmonary nodules may represent malignant lesions. In a series of patients undergoing video-assisted thoracoscopic resection of small (≤ 1 cm) pulmonary nodules, 28 malignant lesions were diagnosed in 27 patients with a history of previous malignancy; 23 lesions (84%) were malignant, including 15 metastases (54%) and eight new lung carcinomas (29%), and five nodules (18%) were benign [12]. The specificity of CT in any given series depends on several variables: 1) the propensity of the underlying ETM to disseminate to the lungs; 2) the stage of the ETM; 3) selection factors for the study population; and 4) patient age, smoking history, history of prior treatment for the ETM, and likelihood of prior granulomatous disease. In addition, it has been reported that intraoperative palpation of the lungs is still warranted to detect metastatic lesions not detected by spiral CT. In one study, 22% (9/41) more malignant nodules were found intraoperatively than were detected by helical CT [13].

Computer-aided detection (CAD) for pulmonary metastatic disease has been adapted to chest CT from applications from mammography. Although these programs are in their developmental phases, it has been suggested that CAD can be used as a second look after the radiologist has completed reviewing the study [14]. Nevertheless, these programs require more development and currently can only be used when there is limited breathing artifact and stable lung expansion. A recent study demonstrated that CAD detected 82.4% of known pulmonary nodules under ideal conditions [14]. CAD is still in the experimental phase and currently has limited use in evaluation of patients with pulmonary metastatic disease.

The utility of CT for evaluating intravascular pulmonary metastatic disease has also been described recently. Liver, kidney, stomach, and breast carcinoma as well as sarcomas have been reported to embolize to the pulmonary vasculature [15]. Differentiation between metastatic disease and thromboembolic disease can be difficult. Ting et al [15] describe morphological features such as tubular and beaded appearance to help distinguish between the two. With improvement of CT resolution, such intravascular metastatic disease will be more readily detectable [16].

CT can be used not only to detect pulmonary metastatic disease but also to determine response to therapy. Patients can be evaluated by follow-up CT to determine if nodule size is decreasing or if there are any new pulmonary metastases. CT volumetry has been discussed in the literature and may prove to be a more accurate method for determining response to therapy, especially when there are small changes in tumor size [17].

Recommendations for the use of CT in detecting pulmonary metastases must be tailored for each ETM. Even for an individual ETM, however, it may still be difficult to arrive at a consensus for the optimal application of CT. Some guidelines for chest CT surveillance in a few common primary tumors, as determined from review of the recent literature, are summarized below.

Bone and Soft-Tissue Sarcomas

Despite multiagent chemotherapy regimens and radical resection of the primary tumor, a large number of patients with bone and soft-tissue sarcomas will have relapse, manifested by dissemination of disease to the lungs as the first site of metastasis. One review of the published literature for osteosarcoma recommends aggressive surgical resection of synchronous and metachronous pulmonary metastases, even if multiple thoracotomies are required [18]. The authors state that CT is the preferred study in the screening for such metastases, although up to twice as many lesions may be found at thoracotomy.

A more recent study on pediatric bone and soft-tissue sarcomas discusses the importance of recognizing that imaging features on CT are important in determining the presence of metastatic disease. In this review of 210 patients, 41.7% of patients who underwent biopsy or resection of a pulmonary nodule had metastasis. Those with three or more nodules, bilateral distribution of disease, and/or large nodule size were more likely to have metastasis [19].

The presence of nodules, regardless of their size, has an impact on long-term survival in patients with soft-tissue sarcomas. In a recent study, 331 sarcoma patients were followed. Of these, 71 had small, indeterminate nodules detected on CT. Of these patients (28%) developed pulmonary metastatic disease, most (90%) in the area of the original indeterminate nodule [20].

Other authors, in a study of 5-year survival after pulmonary metastasectomy for soft-tissue sarcoma, determined through multivariate analysis that the number of nodules detected by preoperative CT has prognostic value, and they recommend routine use of CT [21]. In another study of patients with high-grade soft-tissue sarcomas undergoing metastasectomy, a specific protocol for follow-up is described: routine chest radiographs and chest CT are performed for the first 5 years, with a radiograph obtained at each visit and chest CT performed every 3 months for the first year, every 4 months for the second year, every 6 months for the third year, and once yearly thereafter [22].

Renal Cell Carcinoma

Pulmonary metastases from renal cell carcinoma are seen in 25%-30% of patients at the time of initial diagnosis, and in 30%-50% of patients at a later time [23]. In patients with metastases to the lungs, surgical resection may provide the only effective treatment, in light of the fact that the 5-year survival rate is <5% for stage IV disease [24]. Based on their own experience and a review of the literature, Lim and Carter [24] recommend PA and lateral chest radiographs as an initial test. In patients with low-stage (T1) disease and a normal chest radiograph, CT is not necessary; if the chest radiograph demonstrates multiple nodules, CT is not necessary unless it is required as part of the protocol for systemic therapy. The authors propose that indications for chest CT should include: 1) a solitary pulmonary nodule on the chest radiograph; 2) symptoms suggestive of endobronchial metastasis; 3) extensive regional disease; and 4) presence of other extrathoracic metastases that might be amenable to resection. Other authors advocate a more aggressive approach, with biannual chest radiographs and chest CT examinations [23]. They recommend that such surveillance be life-long, in view of the possibility of delayed recurrent pulmonary metastases.

Testicular Cancer

See and Hoxie [25] suggest that the risk of intrathoracic metastases is correlated with the presence of abnormal findings on abdominal CT. In their study, 74 of 155 patients with seminomatous or nonseminomatous testicular germ cell tumors had imaging by both chest radiographs and chest CT scans concurrently at the time of initial staging. Findings were compared to those of patients having negative or abnormal abdominal CT scans. For the group of 42 patients with negative abdominal CT scans, results of chest CT did not increase the yield for diagnosis of metastases as compared with the chest radiograph; a 2.3% chest CT false-positive rate is in fact cited as a potential source of morbidity in the workup of patients. For the group of 32 patients with abnormal abdominal CT, however, chest CT allowed detection of pulmonary metastases not seen on the chest radiograph in 12.5% of cases. For initial staging workup, the authors therefore recommend chest radiographs for patients with a negative abdominal CT and chest CT for patients with an abnormal abdominal CT.

Malignant Melanoma

Recommendations for chest CT scanning in malignant melanoma appear to be largely determined by the stage of the primary tumor. Buzaid et al [26] retrospectively assessed the role of CT (neck, chest, abdomen, and pelvis) in detecting occult distant metastases in 89 asymptomatic patients with local-regional melanoma who had normal chest radiographs and serum lactate dehydrogenase levels. In only one case was there evidence of disease on chest CT not seen on the chest radiograph, and the authors concluded that chest CT may not be indicated. A large retrospective study of asymptomatic patients with stage III melanoma, assessing the role of CT (head, chest, abdomen, pelvis), suggests that chest CT should be used

selectively in patients with cervical adenopathy [27]. In a review of the role of surgical resection for melanoma metastatic to the lung, Ollila and Morton [28] emphasized that metastasectomy may be the only potentially curative treatment modality in stage IV disease. While noting that metastasectomy is believed to improve survival in patients with one or two pulmonary nodules, they cautioned that the number of lesions should not be an absolute contraindication to surgery. They recommended that preoperative evaluation of patients for pulmonary metastasectomy should include not only chest CT to determine the number of nodules but also whole-body imaging to detect or exclude other extrapulmonary stage IV disease.

Head and Neck Carcinoma

Although the lungs are the most common site of distant metastases in squamous cell carcinoma (SCCA) of the head and neck, there is no clear consensus as to the optimal imaging modality for surveillance. An issue of particular importance in this population is the known increased incidence (15%-30%) of second primary malignancies, including neck, lung, and esophageal cancers [29]. In one retrospective study, only two of 57 patients with head and neck SCCA (stage not specified) had malignancy in the form of synchronous tumors identified on routine chest CT, and these lesions were also evident on chest radiographs [30]. Other authors, however, have observed that chest CT demonstrates a high number of malignancies, including both pulmonary metastases and additional thoracic malignancies, in patients with advanced SCCA [29]. Among 93 patients undergoing chest CT at the time of initial presentation, during routine follow-up or at the time of local-regional neck recurrence, a total of 24 (25.8%) had identification of thoracic malignancy, including 14 (15%) with pulmonary metastases, five (5.4%) with lung carcinoma, and one (1.1%) with esophageal carcinoma. Except for two patients with initial stage I or II disease and local-regional neck recurrence, these patients all had stage III or IV disease.

Magnetic Resonance Imaging

MRI has been considered an alternative to CT for detecting pulmonary metastases, primarily because exposure to ionizing radiation would be avoided, an issue of particular concern with young patients undergoing multiple follow-up examinations. Nevertheless, it is generally accepted that MRI does not currently have a role in screening of patients for pulmonary metastases [3,31]. Motion-related artifacts, a lower spatial resolution than CT, and an inability to detect calcification within lesions all represent limitations of MRI [31]. A recent study comparing turbo-spin echo MRI with spiral CT as a gold standard demonstrated a lower sensitivity for MRI in detecting pulmonary metastases; for 340 metastases identified on CT, the overall sensitivity of MRI was 84%, but for nodules <5 mm in diameter, sensitivity was only 36% [31]. A more recent study comparing various MRI sequences in determining nodule detection accuracy supports these results. The optimal sequence (triggered

short time inversion recovery [STIR]) had 72% sensitivity for nodule detection. These nodules were all previously seen on CT and were >5 mm [32].

Scintigraphy

The use of scintigraphy in conjunction with tumor-seeking agents may offer significant incremental information, enhancing the specificity of diagnosis, as compared with conventional morphologic imaging techniques. There are preliminary reports of results for a variety of scintigraphic techniques applied to a number of different malignancies, but the ultimate role of such imaging has yet to be established.

Imaging with fluorine-18-2-fluoro-2-deoxy-D-glucose positron emission tomography (FDG-PET) is increasingly being used in the staging of patients with bronchogenic carcinoma, not only to detect any nodal involvement but also to check for possible distant metastases. Its role in detecting pulmonary metastases from known ETM is well established. One study demonstrated the utility of FDG-PET in detecting occult extrapulmonary disease in patients with pulmonary metastatic melanoma [33]. In particular, it was determined to be useful in excluding extrapulmonary metastatic melanoma prior to surgery, and the authors concluded that PET scanning should be used in patients with pulmonary metastatic melanoma prior to metastectomy [33]. Use of FDG-PET in the staging of malignant melanoma has also been investigated, but it is acknowledged that this technique has limited sensitivity for small pulmonary nodules, and that false-positive results may occur because of inflammatory processes [34]. The role of FDG-PET in staging head and neck tumors has also been discussed. Although, at times, it is helpful in detecting pulmonary metastasis, in one study of 86 patients, thoracic malignancy was suspected in 23 patients (27%) by uptake criteria. Of these suspicious lesions, 83% were found to be benign [35]. Use of FDG-PET alone does not negate the need for spiral CT in evaluating pulmonary metastatic disease. A negative FDG-PET examination cannot exclude metastatic disease [36]. This is thought to be due to small metastatic nodules.

The use of PET/CT versus CT alone for evaluating pulmonary metastatic disease has been discussed in the literature. PET/CT is commonly used for evaluating patients with primary head and neck tumors. In one study of 24 consecutive patients, there was no statistically significant difference between PET/CT and CT alone [37]. However, this study is limited in that it only looked at nodules >1 cm. Currently, high-resolution chest CT is more sensitive in evaluating pulmonary metastasis than PET/CT, especially for metastasis measuring <1 cm [37].

Other radiopharmaceuticals have also been used in the past. In one older study, encouraging results were reported for the use of ^{99m}Tc-methoxyisobutylisonitrile scintigraphy in 81 patients with a history of previously excised malignant melanoma [38]. Such whole-body scanning correctly detected 92% of 74 metastatic lesions at various sites, including eight lung lesions ranging from

1.2 to 6.0 cm in size, two of which were not previously diagnosed. Use of an indium-111-labeled monoclonal antibody (CCR 086) for detecting colorectal metastases at various sites, including lung lesions as small as 1 cm, has been reported [39]. Bone scintigraphy with single photon emission tomography (SPECT) can be useful in patients with osteosarcoma metastasis [40]. More recently, ^{99m}Tc-depreotide has shown promise in detecting pulmonary metastasis from renal cell carcinoma that is somatostatin receptor-positive [41].

Summary

- Chest radiograph should be performed as a baseline in patients with primary neoplasms known to metastasize to the pulmonary system.
- In many cases, a chest CT without contrast should be performed.
- A chest CT should be performed as an initial evaluation for patients with bone and soft-tissue sarcoma, malignant melanoma, and head and neck carcinoma.
- In patients with primary renal cell or testicular carcinoma, chest CT should be performed if based on the presence of metastatic disease elsewhere.

Relative Radiation Level Information

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, both because of organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared to those specified for adults (see Table below). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® [Radiation Dose Assessment Introduction](#) document.

Relative Radiation Level Designations		
Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
O	0 mSv	0 mSv
☼	<0.1 mSv	<0.03 mSv
☼☼	0.1-1 mSv	0.03-0.3 mSv
☼☼☼	1-10 mSv	0.3-3 mSv
☼☼☼☼	10-30 mSv	3-10 mSv
☼☼☼☼☼	30-100 mSv	10-30 mSv

*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as NS (not specified).

Supporting Document(s)

- [ACR Appropriateness Criteria® Overview](#)
- [Procedure Contrast Information](#)
- [Evidence Table](#)

References

- Davis SD. CT evaluation for pulmonary metastases in patients with extrathoracic malignancy. *Radiology* 1991; 180(1):1-12.
- Snyder BJ, Pugatch RD. Imaging characteristics of metastatic disease to the chest. *Chest Surg Clin N Am* 1998; 8(1):29-48.
- Herold CJ, Bankier AA, Fleischmann D. Lung metastases. *Eur Radiol* 1996; 6(5):596-606.
- Puglisi F, Follador A, Minisini AM, et al. Baseline staging tests after a new diagnosis of breast cancer: further evidence of their limited indications. *Ann Oncol* 2005; 16(2):263-266.
- Terhune MH, Swanson N, Johnson TM. Use of chest radiography in the initial evaluation of patients with localized melanoma. *Arch Dermatol* 1998; 134(5):569-572.
- Mooney MM, Mettlin C, Michalek AM, Petrelli NJ, Kraybill WG. Life-long screening of patients with intermediate-thickness cutaneous melanoma for asymptomatic pulmonary recurrences: a cost-effectiveness analysis. *Cancer* 1997; 80(6):1052-1064.
- Gourin CG, Watts TL, Williams HT, Patel VS, Bilodeau PA, Coleman TA. Identification of distant metastases with positron-emission tomography-computed tomography in patients with previously untreated head and neck cancer. *Laryngoscope* 2008; 118(4):671-675.
- Remy-Jardin M, Remy J, Giraud F, Marquette CH. Pulmonary nodules: detection with thick-section spiral CT versus conventional CT. *Radiology* 1993; 187(2):513-520.
- McCormack PM, Ginsberg KB, Bains MS, et al. Accuracy of lung imaging in metastases with implications for the role of thoracoscopy. *Ann Thorac Surg* 1993; 56(4):863-865; discussion 865-866.
- Peldschus K, Herzog P, Wood SA, Cheema JI, Costello P, Schoepf UJ. Computer-aided diagnosis as a second reader: spectrum of findings in CT studies of the chest interpreted as normal. *Chest* 2005; 128(3):1517-1523.
- Rubin GD, Lyo JK, Paik DS, et al. Pulmonary nodules on multi-detector row CT scans: performance comparison of radiologists and computer-aided detection. *Radiology* 2005; 234(1):274-283.
- Munden RF, Pugatch RD, Liptay MJ, Sugarbaker DJ, Le LU. Small pulmonary lesions detected at CT: clinical importance. *Radiology* 1997; 202(1):105-110.
- Parsons AM, Dettlerbeck FC, Parker LA. Accuracy of helical CT in the detection of pulmonary metastases: is intraoperative palpation

still necessary? *Ann Thorac Surg* 2004; 78(6):1910-1916; discussion 1916-1918.

- Lee KW, Kim M, Gierada DS, Bae KT. Performance of a computer-aided program for automated matching of metastatic pulmonary nodules detected on follow-up chest CT. *AJR* 2007; 189(5):1077-1081.
- Ting PT, Burrowes PW, Gray RR. Intravascular pulmonary metastases from sarcoma: appearance on computed tomography in 3 cases. *Can Assoc Radiol J* 2005; 56(4):214-218.
- Shepard JA, Moore EH, Templeton PA, McLoud TC. Pulmonary intravascular tumor emboli: dilated and beaded peripheral pulmonary arteries at CT. *Radiology* 1993; 187(3):797-801.
- Marten K, Auer F, Schmidt S, Rummeny EJ, Engelke C. Automated CT volumetry of pulmonary metastases: the effect of a reduced growth threshold and target lesion number on the reliability of therapy response assessment using RECIST criteria. *Eur Radiol* 2007; 17(10):2561-2571.
- La Quaglia MP. Osteosarcoma. Specific tumor management and results. *Chest Surg Clin N Am* 1998; 8(1):77-95.
- Absalon MJ, McCarville MB, Liu T, Santana VM, Daw NC, Navid F. Pulmonary nodules discovered during the initial evaluation of pediatric patients with bone and soft-tissue sarcoma. *Pediatr Blood Cancer* 2008; 50(6):1147-1153.
- Rissing S, Rougraff BT, Davis K. Indeterminate pulmonary nodules in patients with sarcoma affect survival. *Clin Orthop Relat Res* 2007; 459:118-121.
- Casson AG, Putnam JB, Natarajan G, et al. Five-year survival after pulmonary metastasectomy for adult soft tissue sarcoma. *Cancer* 1992; 69(3):662-668.
- Robinson MH, Sheppard M, Moskovic E, Fischer C. Lung metastasectomy in patients with soft tissue sarcoma. *Br J Radiol* 1994; 67(794):129-135.
- Cerfolio RJ, Allen MS, Deschamps C, et al. Pulmonary resection of metastatic renal cell carcinoma. *Ann Thorac Surg* 1994; 57(2):339-344.
- Lim DJ, Carter MF. Computerized tomography in the preoperative staging for pulmonary metastases in patients with renal cell carcinoma. *J Urol* 1993; 150(4):1112-1114.
- See WA, Hoxie L. Chest staging in testis cancer patients: imaging modality selection based upon risk assessment as determined by abdominal computerized tomography scan results. *J Urol* 1993; 150(3):874-878.
- Buzaid AC, Tinoco L, Ross MI, Legha SS, Benjamin RS. Role of computed tomography in the staging of patients with local-regional metastases of melanoma. *J Clin Oncol* 1995; 13(8):2104-2108.
- Kuvshinoff BW, Kurtz C, Coit DG. Computed tomography in evaluation of patients with stage III melanoma. *Ann Surg Oncol* 1997; 4(3):252-258.
- Ollila DW, Morton DL. Surgical resection as the treatment of choice for melanoma metastatic to the lung. *Chest Surg Clin N Am* 1998; 8(1):183-196.
- Mercader VP, Gatenby RA, Mohr RM, Fisher MS, Caroline DF. CT surveillance of the thorax in patients with squamous cell carcinoma of the head and neck: a preliminary experience. *J Comput Assist Tomogr* 1997; 21(3):412-417.
- Nilssen EL, Murthy P, McClymont L, Denholm S. Radiological staging of the chest and abdomen in head and neck squamous cell carcinoma--are computed tomography and ultrasound necessary? *J Laryngol Otol* 1999; 113(2):152-154.
- Kersjes W, Mayer E, Buchenroth M, Schunk K, Fouda N, Cagil H. Diagnosis of pulmonary metastases with turbo-SE MR imaging. *Eur Radiol* 1997; 7(8):1190-1194.
- Bruegel M, Gaa J, Woertler K, et al. MRI of the lung: value of different turbo spin-echo, single-shot turbo spin-echo, and 3D gradient-echo pulse sequences for the detection of pulmonary metastases. *J Magn Reson Imaging* 2007; 25(1):73-81.
- Dalrymple-Hay MJ, Rome PD, Kennedy C, Fulham M, McCaughan BC. Pulmonary metastatic melanoma -- the survival benefit associated with positron emission tomography scanning. *Eur J Cardiothorac Surg* 2002; 21(4):611-614; discussion 614-615.
- Paquet P, Hustinx R, Rigo P, Pierard GE. Malignant melanoma staging using whole-body positron emission tomography. *Melanoma Res* 1998; 8(1):59-62.
- Roh JL, Ryu CH, Kim JS, et al. Clinical significance of intrathoracic lesions detected by 18F-fluorodeoxyglucose positron

- emission tomography in the management of patients with head and neck cancer. *Oral Oncol* 2007; 43(8):757-763.
36. Franzius C, Daldrup-Link HE, Sciuk J, et al. FDG-PET for detection of pulmonary metastases from malignant primary bone tumors: comparison with spiral CT. *Ann Oncol* 2001; 12(4):479-486.
 37. De Wever W, Meylaerts L, De Ceuninck L, Stroobants S, Verschakelen JA. Additional value of integrated PET-CT in the detection and characterization of lung metastases: correlation with CT alone and PET alone. *Eur Radiol* 2007; 17(2):467-473.
 38. Alonso O, Martinez M, Mut F, et al. Detection of recurrent malignant melanoma with 99mTc-MIBI scintigraphy. *Melanoma Res* 1998; 8(4):355-360.
 39. Abdel-Nabi HH, Levine G, Lamki LM, et al. Colorectal carcinoma metastases: detection with In-111-labeled monoclonal antibody CCR 086. *Radiology* 1990; 176(1):117-122.
 40. Pevarski DJ, Drane WE, Scarborough MT. The usefulness of bone scintigraphy with SPECT images for detection of pulmonary metastases from osteosarcoma. *AJR* 1998; 170(2):319-322.
 41. Shih WJ, Romero RA, Timothy M, Milan PP. 99mTc-Depreotide chest SPECT demonstrates pulmonary metastases from renal cell carcinoma. *J Nucl Med Technol* 2004; 32(1):19-21.

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.