

<p>Table 1</p> <p>Indications for Use of Iodinated Contrast Media</p>

Intravascular

Intravenous

- Computed tomography
- Digital subtraction angiography
- Intravenous urography
- Venography (phlebography)
 - Inferior vena cava and its tributaries
 - Superior vena cava and its tributaries
 - Extremities
 - Other venous sites
- Epidural venography

Intra-arterial

- Angiocardiology
- Computed tomography
- Coronary angiography
- Pulmonary angiography
- Aortography
- Visceral and peripheral arteriography
- Digital subtraction angiography
- Central nervous system
 - Cerebral, vertebral, and spinal angiography

Intrathecal (Use U.S. Food and Drug Administration-approved contrast media only)

- Myelography (myelographic nonionic only)
- Cysternography (myelographic nonionic only)

Other

- Oral, rectal, or ostomy – gastrointestinal tract
 - Conventional fluoroscopy
 - Computed tomography
 - Therapeutic uses
- Body cavity use
 - Herniography
 - Peritoneography
 - Vaginography
- Hysterosalpingography
- Arthrography
- Endoscopic retrograde cholangiopancreatography
- Cholangiography
- Nephrostography
- Pyelography – antegrade, retrograde
- Urethrography – voiding, retrograde
- Cystography
- Sialography
- Ductography (breast)
- Miscellaneous
 - Sinus tract injection
 - Cavity delineation (including urinary diversions, such as loop and pouch)