

The American College of Radiology, with more than 30,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice guidelines and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice guidelines and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice guideline and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review, requiring the approval of the Commission on Quality and Safety as well as the ACR Board of Chancellors, the ACR Council Steering Committee, and the ACR Council. The practice guidelines and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice guideline and technical standard by those entities not providing these services is not authorized.

Revised 2010 (Resolution 17)*

ACR–SSR PRACTICE GUIDELINE FOR THE PERFORMANCE AND INTERPRETATION OF MAGNETIC RESONANCE IMAGING (MRI) OF THE SHOULDER

PREAMBLE

These guidelines are an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. They are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care. For these reasons and those set forth below, the American College of Radiology cautions against the use of these guidelines in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the physician or medical physicist in light of all the circumstances presented. Thus, an approach that differs from the guidelines, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in the guidelines when, in the reasonable judgment of the practitioner, such course of action is indicated by the condition of the patient, limitations of available resources, or advances in knowledge or technology subsequent to publication of the guidelines. However, a practitioner who employs an approach substantially different from these guidelines is advised to document in the patient record information sufficient to explain the approach taken.

The practice of medicine involves not only the science, but also the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment.

Therefore, it should be recognized that adherence to these guidelines will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The sole purpose of these guidelines is to assist practitioners in achieving this objective.

I. INTRODUCTION

This guideline was developed and written collaboratively by the American College of Radiology (ACR) and the Society of Skeletal Radiology (SSR).

Magnetic resonance imaging (MRI) is an established and proven imaging modality for the detection, evaluation, assessment, staging, and follow-up of disorders of the shoulder. Properly performed and interpreted, MRI contributes not only to diagnosis but also to treatment planning and prognostication. However, it should be performed only for a valid medical reason and after careful consideration of alternative diagnostic modalities. MRI of the shoulder may be performed without contrast, following intra-articular contrast injection (“direct” MR arthrography) to increase conspicuity of intra-articular abnormalities, or with intravenous contrast to identify hyperemic lesions or to create “indirect” arthrographic images by enhancing synovial-lined structures and their contents.

An analysis of the strengths and potential risks of MRI and other diagnostic modalities should be weighed against their suitability for specific patients and particular clinical conditions. Radiographs usually are the first imaging test

performed for most suspected abnormalities in the shoulder and will often suffice to diagnose or exclude an abnormality, or will direct further imaging evaluation. Radionuclide bone scanning can screen the entire skeleton in addition to the shoulder for radiographically occult bone disease such as metastases. Other nuclear medicine examinations have a role for specific clinical scenarios (e.g., a labeled white blood cell study for suspected osteomyelitis). Conventional single-contrast or double-contrast arthrography can accurately depict most articular surface and full-thickness tears of the rotator cuff [1,2]. Sonography can be used to evaluate the rotator cuff and biceps tendon and has the advantage of imaging during physiologic motion [3-6]. Computed tomography (CT) is used to evaluate the bone integrity of the glenoid fossa and humerus, and the alignment and congruence of the glenohumeral joint [7]. When combined with arthrography, CT can also be used for evaluating the labrum, articular cartilage, and loose bodies [8]. Lastly, arthroscopy provides a detailed examination of the internal structures of the shoulder, allowing the surgeon to treat as well as diagnose many internal derangements.

While MRI is one of the most sensitive diagnostic tests for detecting anatomic abnormalities of the extremities, findings may be misleading if not closely correlated with other imaging studies, clinical history, clinical examination, and physiologic tests. Adherence to the following guideline will enhance the probability of accurately diagnosing such abnormalities.

II. QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL

See the [ACR Practice Guideline for Performing and Interpreting Magnetic Resonance Imaging \(MRI\)](#). The interpreting physician needs a thorough knowledge and understanding of the anatomy of the shoulder, including the numerous normal variations in the glenohumeral capsular and labral configurations, and their corresponding MR imaging appearances.

III. INDICATIONS

A. Primary indications for MRI of the shoulder include, but are not limited to, diagnosis, exclusion, and grading of suspected:

1. Rotator cuff tendon abnormalities: full-thickness, partial-thickness, and recurrent (postoperative) tears, tendinopathy, tendinitis, and cuff tear arthropathy † [9-19].
2. Disorders of the long head of the biceps brachii: full-thickness, partial-thickness, and recurrent (postoperative) tears, tendinopathy, tendinitis, subluxation, and dislocation † [17,20-22].
3. Conditions affecting the supraspinatus outlet: acromial shape, os acromiale, subacromial spurs,

acromioclavicular joint disorders, coracoacromial ligament integrity, subacromial bursitis † [12,23-26].

4. Labral abnormalities: cysts, degeneration, and tears, including superior labrum anterior posterior (SLAP) lesions, Bankart lesions and their variants, and recurrent (postoperative) labral tears † [8,17,27-41].
5. Abnormalities of the rotator interval and biceps pulley † [20,42,43].
6. Muscle disorders affecting the shoulder girdle: atrophy, hypertrophy, denervation, masses, injuries [15,19,44-50].
7. Glenohumeral chondral and osteochondral abnormalities: osteochondral fractures and osteochondritis dissecans, articular cartilage degeneration, fissures, fractures, flaps, and separations † [51-53].
8. Intra-articular bodies †.
9. Synovial-based disorders: synovitis, bursitis, metaplasia, and neoplasia * [54,55].
10. Marrow abnormalities: osteonecrosis, marrow edema syndromes, and stress fractures * [56].
11. Neoplasms, masses, and cysts of bone, joint, or soft tissue * [26,36,57].
12. Infections of bone, joint, or soft tissue * [58,59].
13. Congenital and developmental conditions including dysplasia and normal variants* [60-63].
14. Vascular conditions: entrapment, aneurysm, stenosis, and occlusion * [64].
15. Neurologic conditions: entrapment, compression, masses, and peripheral neuritis * [34,39,65].

B. MRI of the shoulder may be indicated to further clarify and stage conditions diagnosed clinically and/or suggested by other imaging modalities, including, but not limited to:

1. Arthritides: inflammatory, infectious, neuropathic, degenerative, crystal-induced, post-traumatic * [26,66,67].
2. Frozen shoulder and adhesive capsulitis † [42].
3. Primary and secondary bone and soft tissue tumors * [57].
4. Fractures and dislocations [23,68,69].

C. MRI of the shoulder may be useful to evaluate specific clinical scenarios, including, but not limited to:

1. Prolonged, refractory, or unexplained shoulder pain *†.
2. Acute shoulder trauma [23,69].
3. Impingement syndromes: subacromial, subcoracoid, internal † [12,24,25,70-73].
4. Glenohumeral instability: chronic, recurrent, subacute, and acute dislocation and subluxation † [40,60,69,74-76].

5. Shoulder symptoms in the overhead or throwing athlete † [77-80].
6. Mechanical shoulder symptoms: catching, locking, snapping, crepitus †.
7. Limited or painful range of motion.
8. Swelling, enlargement, mass, or atrophy * [36].
9. Patients for whom diagnostic or therapeutic arthroscopy is planned †.
10. Patients with recurrent, residual, or new symptoms following shoulder surgery † [11,15,17,19,48,81,82].

* Conditions in which intravenous contrast may be useful.

† Conditions in which intra-articular contrast (performed by direct intra-articular injection or indirect joint opacification following intravenous administration) may be useful.

IV. SAFETY GUIDELINES AND POSSIBLE CONTRAINDICATIONS

See the [ACR Practice Guideline for Performing and Interpreting Magnetic Resonance Imaging \(MRI\)](#), the ACR Guidance Document for Safe MR Practices [83], and the [ACR Manual on Contrast Media](#) [84].

Peer-reviewed literature pertaining to MR safety should be reviewed on a regular basis [85,86].

V. SPECIFICATIONS OF THE EXAMINATION

The supervising physician must have complete understanding of the indications, risks, and benefits of the examination, as well as alternative imaging procedures. The physician must be familiar with potential hazards associated with MRI, including potential adverse reactions to contrast media. The physician should be familiar with relevant ancillary studies that the patient may have undergone. The physician performing MRI interpretation must have a clear understanding and knowledge of the anatomy and pathophysiology relevant to the MRI examination.

The written or electronic request for MRI of the shoulder should provide sufficient information to demonstrate the medical necessity of the examination and allow for its proper performance and interpretation.

Documentation that satisfies medical necessity includes 1) signs and symptoms and/or 2) relevant history (including known diagnoses). Additional information regarding the specific reason for the examination or a provisional diagnosis would be helpful and may at times be needed to allow for the proper performance and interpretation of the examination.

The request for the examination must be originated by a physician or other appropriately licensed health care

provider. The accompanying clinical information should be provided by a physician or other appropriately licensed health care provider familiar with the patient's clinical problem or question and consistent with the state's scope of practice requirements. (ACR Resolution 35, adopted in 2006)

The supervising physician must also understand the pulse sequences to be used and their effect on the appearance of the images, including the potential generation of image artifacts. Standard imaging protocols may be established and varied on a case-by-case basis when necessary. These protocols should be reviewed and updated periodically.

A. Patient Selection

The physician responsible for the examination should supervise patient selection and preparation, and be available in person or by phone for consultation. Patients must be screened and interviewed prior to the examination to exclude individuals who may be at risk by exposure to the MR environment.

Certain indications require administration of intravenous (IV) contrast media. IV contrast enhancement should be performed using appropriate injection protocols and in accordance with the institution's policy on IV contrast utilization. (See the [ACR Practice Guideline for the Use of Intravascular Contrast Media](#).)

Patients suffering from anxiety or claustrophobia may require sedation or additional assistance. Administration of moderate sedation may be needed to achieve a successful examination. If moderate sedation is necessary, refer to the [ACR-SIR Practice Guideline for Sedation/Analgesia](#).

B. Facility Requirements

Appropriate emergency equipment and medications must be immediately available to treat adverse reactions associated with administered medications. The equipment and medications should be monitored for inventory and drug expiration dates on a regular basis. The equipment, medications, and other emergency support must also be appropriate for the range of ages and sizes in the patient population.

C. Examination Technique

Shoulder MRI can be performed using a variety of magnet designs (closed or open) and field strengths (low, medium, or high) [13,35,87-91]. Since the inherent signal-to-noise ratio is reduced with lower field strength MR systems, imaging parameters may require modifications. For lower field strength systems, for example, the number of acquisitions can be increased, at

the expense of longer imaging times and increased risk of involuntary patient motion [87,89,92,93]. Alternatively, the voxel size can be increased (by a combination of larger field of view (FOV), thicker slices, and/or decreased matrix) at the expense of spatial resolution [89,90,93,94]. Fat suppression techniques that rely on the difference between fat and water frequencies (chemical shifts) are unreliable at low field strength, and substituting short-T1 inversion recovery (STIR) images may be necessary [87,90]. Even when the imaging protocol is optimized for shoulder imaging on a low-field open system, subjective image quality will likely be inferior to that obtained with a high-field system [90,93]. Various investigators using different equipment and scanning parameters have reached contradictory conclusions regarding the diagnostic performance of low-field-strength MR scanners for shoulder disorders. Some studies have found that the accuracy for complete and partial rotator cuff tears and for labral abnormalities is not significantly different for open, low-field and closed, high-field systems, with careful attention to technique [88,90,91,95]. MR arthrography can further enhance the diagnostic yield for shoulder MRI performed on low-field-strength systems [87,93]. Other investigators have found lower accuracy for evaluating disorders like SLAP tears, capsular abnormalities, and small rotator cuff tears with specific low-field systems compared to high-field ones [94-96].

Regardless of system design, a local coil is mandatory to maximize the signal-to-noise ratio. Commercially available coils appropriate for shoulder imaging include single-loop contoured or flat-surface coils [97,98], paired coils in a Helmholtz configuration [33,99], circularly polarized flexible coils [94], solenoid coils [89], and phased array designs [28,38].

Patients are positioned supine with the affected arm at the side. For evaluation of the rotator cuff and anterior labrum, internal rotation of the arm should be avoided [76,98,100]. When MR arthrography is performed, repositioning the affected arm into the abduction external rotation (ABER) position may increase sensitivity for anterior inferior labral tears [8,29,101,102] and may increase accuracy for rotator cuff tears, especially partial-thickness undersurface tears [103-105]. Images with the patient's arm in this position are obtained parallel to the humeral shaft prescribed from a coronal localizer image [29].

Shoulder MR examinations usually include images acquired in the transverse, oblique sagittal, and oblique coronal planes. The oblique sections are prescribed orthogonal to the glenoid fossa or the supraspinatus tendon. Evaluation of the rotator cuff is performed using both oblique coronal and oblique sagittal images [106]. Prescribing the oblique sagittal images in the frontal plane so that they are perpendicular to the distal supraspinatus

tendon may be useful for identifying subtle partial-thickness rotator cuff tears [107]. The oblique coronal and oblique sagittal images can be used to evaluate the labrum, biceps tendon, acromial anatomy, supraspinatus outlet, acromioclavicular joint, and rotator interval [20,24,43]. The transverse images best display the subscapularis tendon, the long head of the biceps tendon, glenohumeral articulation, and the glenoid labrum [8,21,30,33]. Transverse images may aid in detecting anterior rotator cuff tears. Radial imaging for the glenoid labrum has been reported [108], but it is not widely used.

FOV should be tailored to the size of the patient and the structures being examined, but for the standard sequences, the FOV should be 16 cm or smaller on medium-field and high-field units. Larger FOVs and smaller imaging matrices may be necessary on lower field systems but will result in lower spatial resolution, limiting the sensitivity of the examination [90,93]. Occasionally, additional sequences with a larger FOV will be appropriate to more fully evaluate a specific suspected or detected abnormality, for example, in the scapulothoracic articulation or the pectoralis major muscle. Slice thickness in the oblique sagittal and oblique coronal planes of 4 mm or less is needed to demonstrate subtle tendon pathology, but thinner sections may be advantageous for detailed analysis of other structures such as the labrum and articular cartilage. An interslice gap may be selected to decrease signal loss due to cross talk [110], but should be no more than 33% of the slice width. Two interleaved scans may allow imaging without gaps at the expense of an increase in scan time. The imaging matrix should balance intravoxel signal-to-noise ratio with desired in-plane spatial resolution and reduction of truncation artifacts, but should be at least 160 steps in the phase direction and 256 steps in the frequency direction for 2D imaging, other than when imaging a large tumor. Some practices may use higher imaging matrices (up to 512 steps) to increase spatial resolution for diagnosing labral lesions, including SLAP tears [28,30].

Shoulder MRI can be performed with a wide variety of pulse sequences [111]. The choice of sequences can be tailored to optimize the examination for specific clinical questions, and may vary due to local preferences. Conventional spin-echo, fast (turbo) spin-echo, and gradient-recalled sequences have all been used successfully for shoulder MRI. A typical imaging protocol will be composed of one or more of these pulse sequence types. The prescribed TR, TE, and flip angle will depend on the field strength of the magnet and the relative contrast weighting desired.

Fluid sensitive sequences such as long TR/moderate-to-long TE (proton-density weighted or T2-weighted) images with or without fat suppression, or STIR images are typically used for evaluating the rotator cuff, with either conventional spin-echo or fast (turbo) spin-echo

technique [18,112-114]. T2*-weighted gradient-echo recalled sequences can also be used for diagnosing rotator cuff abnormalities, but probably with lower accuracy compared with conventional spin-echo or fast spin-echo sequences [115,116]. To show labral abnormalities, long-TR (proton-density weighted or T2-weighted) spin-echo or fast spin-echo images, or T2*-weighted gradient recalled images are typically performed [30,33,117], although gradient echo imaging may be less accurate, when used in isolation for anterior labrum abnormalities compared with conventional spin-echo or fast spin-echo imaging [98]. Lesions of the superior labrum such as SLAP tears can be visualized on fast spin-echo, long-TR images [28,38], or with MR arthrography [27,31]. T1-weighted sequences (short TR/short TE) have a role in characterizing marrow abnormalities [68], various stages of hemorrhage [118,119], and muscle pathology [19,44,45,48,49].

MR arthrography using direct intra-articular injection of saline [76] or dilute gadolinium-containing contrast [74] may improve diagnostic accuracy in unstable shoulders [41]. Additionally, MR arthrography may improve diagnostic performance for some rotator cuff tendon tears, particularly partial-thickness tears, postoperative recurrent tears, and subscapularis tears [11,16,101,103,120,121]. Contrast opacification of the glenohumeral joint can also be accomplished indirectly by allowing contrast injected intravenously to diffuse across the synovial membrane; MR imaging in this circumstance is performed after a short delay (during which time the patient may be asked to move or exercise the shoulder) following intravenous injection of a gadolinium-containing agent [103,122]. T1-weighted images either without [8,74,120] or with fat suppression [27,29,121] are most frequently used when direct or indirect MR arthrography is performed with gadolinium-containing contrast. At least one fluid sensitive sequence is still necessary when performing MR arthrography to detect pathology that does not communicate with the joint.

Suppressing the signal from fat may enhance the diagnostic yield of some pulse sequences [111]. Fat suppression can be performed using spectrally selective RF pulses, selective water excitation, a STIR sequence, or a phase-dependent method (e.g., the Dixon method) [87,90,123,124]. The latter two techniques may be necessary on low-field systems [90,105]. The addition of fat suppression may increase diagnostic accuracy for rotator cuff tendon tears [113,123], especially partial-thickness tears [18]. Fat suppression is a useful adjunct to T1-weighted images when MR arthrography is performed using gadolinium-containing contrast [27,29,121].

Additional imaging techniques have specific roles for certain shoulder disorders. Applying axial traction to the affected arm via a weight attached to the wrist may aid in

the visualization of SLAP lesions [125]. The ABER position may help with the MR arthrographic diagnosis of instability lesions and partial-thickness, articular-surface rotator cuff tears [8,29,101-105].

Various techniques are used to minimize artifacts that can reduce imaging quality. Wraparound artifact should be reduced by phase oversampling [126]. Involuntary patient motion is best controlled by ensuring patient comfort combined with gentle immobilization when necessary [111]. Securing the affected arm against the thigh may further reduce motion artifacts [76]. When available, software that compensates for motion by the use of navigator echoes can be useful [127]. Flowing blood and other periodic motions produce ghosting artifacts, which can be reduced with presaturation pulses or gradient moment nulling [126,128]. Chemical shift artifact is more severe at higher field strengths and may necessitate an increase in the receiver bandwidth [13,92,126]. Susceptibility artifacts, which originate from heterogeneity of the local field, are also more severe at higher field strengths and when using gradient-recalled pulse sequences. Avoiding gradient-echo imaging and reducing the voxel size will help reduce the magnitude of susceptibility artifacts [126,127]. Vacuum phenomena in the shoulder joint can also result in artifact generation, especially when gradient-recalled pulse sequences are used [129]. Lastly, magic angle artifact can produce apparent increased signal intensity on short-TE images within the supraspinatus tendon as it curves over the humeral head, mimicking intratendinous pathology [130]. This pitfall is best avoided by confirming abnormal signal intensity in the tendon on long-TR images, and correlating apparent signal intensity abnormalities with changes in tendon thickness.

It is the responsibility of the supervising physician to determine whether additional pulse sequences or imaging techniques would confer added benefit for the diagnosis and management of the patient. Examinations that use techniques not approved by the Food and Drug Administration, such as the intra-articular injection of gadolinium chelates (direct MR arthrography) [131], can be considered when they are judged to be medically appropriate.

VI. DOCUMENTATION

Reporting should be in accordance with the [ACR Practice Guideline for Communication of Diagnostic Imaging Findings](#).

At a minimum, the report should address the condition of the rotator cuff muscles and tendons, supraspinatus outlet, biceps tendon, and labrum. In selected cases, a description of findings in the major ligaments and capsule, articular cartilage, bone marrow, synovium, and cortical bone

would be appropriate. An effort should be made to adopt a standardized lexicon of terms, and the report should use precise anatomic descriptions of identified abnormalities whenever possible [132].

VII. EQUIPMENT SPECIFICATIONS

The MRI equipment specifications and performance must meet all state and federal requirements. The requirements include, but are not limited to, specifications of maximum static magnetic strength, maximum rate of change of the magnetic field strength (dB/dt), maximum radiofrequency power deposition (specific absorption rate), and maximum acoustic noise levels.

VIII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *Position Statement on QC & Improvement, Safety, Infection Control, and Patient Education* on the ACR web page (<http://www.acr.org/guidelines>).

Specific policies and procedures related to MRI safety should be in place with documentation that is updated annually and compiled under the supervision and direction of the supervising MRI physician. Guidelines should be provided that deal with potential hazards associated with the MRI examination of the patient as well as to others in the immediate area [85,86,133]. Screening forms must also be provided to detect those patients who may be at risk for adverse events associated with the MRI examination [85,86,133,134].

Equipment monitoring should be in accordance with the [ACR Technical Standard for Diagnostic Medical Physics Performance Monitoring of Magnetic Resonance Imaging \(MRI\) Equipment](#).

ACKNOWLEDGEMENTS

This guideline was revised according to the process described under the heading *The Process for Developing ACR Practice Guidelines and Technical Standards* on the ACR web page (<http://www.acr.org/guidelines>) by the Musculoskeletal Committee of the Commission on Body Imaging in collaboration with the SSR.

Collaborative Committee

ACR

Mark J. Kransdorf, MD, FACR
Lynne S. Steinbach, MD, FACR

SSR

David A. Rubin, MD, Chair
J.H. Edmund Lee, MD, Co-Chair
Mark W. Anderson, MD
Jesse A. Davila, MD
Mihra S. Taljanovic, MD
Hilary R. Umans, MD

ACR Musculoskeletal Committee on Body Imaging

Mark J. Kransdorf, MD, FACR, Chair
Mark W. Anderson, MD
Murray K. Dalinka, MD, FACR
Arthur A. DeSmet, MD, FACR
David G. Disler, MD, FACR
Jon A. Jacobson, MD
Jeremy J. Kaye, MD
Mark D. Murphey, MD
Leanne L. Seeger, MD, FACR
Lynne S. Steinbach, MD, FACR
James A. Brink, MD, FACR, Chair, Commission

Comments Reconciliation Committee

Howard B. Fleishon, MD, MMM, FACR, Co-Chair
Rodney S. Owen, MD, FACR, Co-Chair
Mark W. Anderson, MD
Kimberly E. Applegate, MD, MS, FACR
James A. Brink, MD, FACR
Jesse A. Davila, MD
Alan D. Kaye, MD, FACR
Mark J. Kransdorf, MD, FACR
Paul A. Larson, MD, FACR
Lawrence A. Liebscher, MD, FACR
David A. Rubin, MD
G. Scott Stacy, MD
Lynne S. Steinbach, MD, FACR
Margaret A. Stull, MD
Mihra S. Taljanovic, MD
Lawrence N. Tanenbaum, MD, FACR
Hilary R. Umans, MD

REFERENCES

1. Farin PU, Kaukanen E, Jaroma H, Vaatainen U, Miettinen H, Soimakallio S. Site and size of rotator-cuff tear. Findings at ultrasound, double-contrast arthrography, and computed tomography arthrography with surgical correlation. *Invest Radiol* 1996; 31:387-394.
2. Mink JH, Harris E, Rappaport M. Rotator cuff tears: evaluation using double-contrast shoulder arthrography. *Radiology* 1985; 157:621-623.
3. Farin P, Jaroma H. Sonographic detection of tears of the anterior portion of the rotator cuff (subscapularis tendon tears). *J Ultrasound Med* 1996; 15:221-225.
4. Farin PU, Jaroma H, Harju A, Soimakallio S. Medial displacement of the biceps brachii tendon: evaluation with dynamic sonography during

- maximal external shoulder rotation. *Radiology* 1995; 195:845-848.
5. Teefey SA, Middleton WD, Bauer GS, Hildebolt CF, Yamaguchi K. Sonographic differences in the appearance of acute and chronic full-thickness rotator cuff tears. *J Ultrasound Med* 2000; 19:377-378; quiz 383.
 6. Teefey SA, Rubin DA, Middleton WD, Hildebolt CF, Leibold RA, Yamaguchi K. Detection and quantification of rotator cuff tears. Comparison of ultrasonographic, magnetic resonance imaging, and arthroscopic findings in seventy-one consecutive cases. *J Bone Joint Surg Am* 2004; 86-A:708-716.
 7. Griffith JF, Yung PS, Antonio GE, Tsang PH, Ahuja AT, Chan KM. CT compared with arthroscopy in quantifying glenoid bone loss. *AJR Am J Roentgenol* 2007; 189:1490-1493.
 8. Chandnani VP, Yeager TD, DeBerardino T, et al. Glenoid labral tears: prospective evaluation with MRI imaging, MR arthrography, and CT arthrography. *AJR Am J Roentgenol* 1993; 161:1229-1235.
 9. Bergin D, Parker L, Zoga A, Morrison W. Abnormalities on MRI of the subscapularis tendon in the presence of a full-thickness supraspinatus tendon tear. *AJR Am J Roentgenol* 2006; 186:454-459.
 10. Deutsch A, Altchek DW, Veltri DM, Potter HG, Warren RF. Traumatic tears of the subscapularis tendon. Clinical diagnosis, magnetic resonance imaging findings, and operative treatment. *Am J Sports Med* 1997; 25:13-22.
 11. Duc SR, Mengiardi B, Pfirrmann CW, Jost B, Hodler J, Zanetti M. Diagnostic performance of MR arthrography after rotator cuff repair. *AJR Am J Roentgenol* 2006; 186:237-241.
 12. Hambly N, Fitzpatrick P, MacMahon P, Eustace S. Rotator cuff impingement: correlation between findings on MRI and outcome after fluoroscopically guided subacromial bursography and steroid injection. *AJR Am J Roentgenol* 2007; 189:1179-1184.
 13. Magee T, Williams D. 3.0-T MRI of the supraspinatus tendon. *AJR Am J Roentgenol* 2006; 187:881-886.
 14. Meister K, Thesing J, Montgomery WJ, Indelicato PA, Walczak S, Fontenot W. MR arthrography of partial thickness tears of the undersurface of the rotator cuff: an arthroscopic correlation. *Skeletal Radiol* 2004; 33:136-141.
 15. Mellado JM, Calmet J, Olona M, et al. Surgically repaired massive rotator cuff tears: MRI of tendon integrity, muscle fatty degeneration, and muscle atrophy correlated with intraoperative and clinical findings. *AJR Am J Roentgenol* 2005; 184:1456-1463.
 16. Pfirrmann CW, Zanetti M, Weishaupt D, Gerber C, Hodler J. Subscapularis tendon tears: detection and grading at MR arthrography. *Radiology* 1999; 213:709-714.
 17. Probyn LJ, White LM, Salonen DC, Tomlinson G, Boynton EL. Recurrent symptoms after shoulder instability repair: direct MR arthrographic assessment--correlation with second-look surgical evaluation. *Radiology* 2007; 245:814-823.
 18. Singson RD, Hoang T, Dan S, Friedman M. MR evaluation of rotator cuff pathology using T2-weighted fast spin-echo technique with and without fat suppression. *AJR Am J Roentgenol* 1996; 166:1061-1065.
 19. Thomazeau H, Boukobza E, Morcet N, Chaperon J, Langlais F. Prediction of rotator cuff repair results by magnetic resonance imaging. *Clin Orthop Relat Res* 1997:275-283.
 20. Morag Y, Jacobson JA, Shields G, et al. MR arthrography of rotator interval, long head of the biceps brachii, and biceps pulley of the shoulder. *Radiology* 2005; 235:21-30.
 21. Spritzer CE, Collins AJ, Cooperman A, Speer KP. Assessment of instability of the long head of the biceps tendon by MRI. *Skeletal Radiol* 2001; 30:199-207.
 22. Zanetti M, Weishaupt D, Gerber C, Hodler J. Tendinopathy and rupture of the tendon of the long head of the biceps brachii muscle: evaluation with MR arthrography. *AJR Am J Roentgenol* 1998; 170:1557-1561.
 23. Alyas F, Curtis M, Speed C, Saifuddin A, Connell D. MR imaging appearances of acromioclavicular joint dislocation. *Radiographics* 2008; 28:463-479; quiz 619.
 24. Epstein RE, Schweitzer ME, Frieman BG, Fenlin JM, Jr., Mitchell DG. Hooked acromion: prevalence on MR images of painful shoulders. *Radiology* 1993; 187:479-481.
 25. Park JG, Lee JK, Phelps CT. Os acromiale associated with rotator cuff impingement: MR imaging of the shoulder. *Radiology* 1994; 193:255-257.
 26. Tshering Vogel DW, Steinbach LS, Hertel R, Bernhard J, Stauffer E, Anderson SE. Acromioclavicular joint cyst: nine cases of a pseudotumor of the shoulder. *Skeletal Radiol* 2005; 34:260-265.
 27. Bencardino JT, Beltran J, Rosenberg ZS, et al. Superior labrum anterior-posterior lesions: diagnosis with MR arthrography of the shoulder. *Radiology* 2000; 214:267-271.
 28. Connell DA, Potter HG, Wickiewicz TL, Altchek DW, Warren RF. Noncontrast magnetic resonance imaging of superior labral lesions. 102 cases confirmed at arthroscopic surgery. *Am J Sports Med* 1999; 27:208-213.

29. Cvitanic O, Tirman PF, Feller JF, Bost FW, Minter J, Carroll KW. Using abduction and external rotation of the shoulder to increase the sensitivity of MR arthrography in revealing tears of the anterior glenoid labrum. *AJR Am J Roentgenol* 1997; 169:837-844.
30. Gusmer PB, Potter HG, Schatz JA, et al. Labral injuries: accuracy of detection with unenhanced MR imaging of the shoulder. *Radiology* 1996; 200:519-524.
31. Jee WH, McCauley TR, Katz LD, Matheny JM, Ruwe PA, Daigneault JP. Superior labral anterior posterior (SLAP) lesions of the glenoid labrum: reliability and accuracy of MR arthrography for diagnosis. *Radiology* 2001; 218:127-132.
32. Jin W, Ryu KN, Kwon SH, Rhee YG, Yang DM. MR arthrography in the differential diagnosis of type II superior labral anteroposterior lesion and sublabral recess. *AJR Am J Roentgenol* 2006; 187:887-893.
33. Legan JM, Burkhard TK, Goff WB, 2nd, et al. Tears of the glenoid labrum: MR imaging of 88 arthroscopically confirmed cases. *Radiology* 1991; 179:241-246.
34. Ludig T, Walter F, Chapuis D, Mole D, Roland J, Blum A. MR imaging evaluation of suprascapular nerve entrapment. *Eur Radiol* 2001; 11:2161-2169.
35. Magee TH, Williams D. Sensitivity and specificity in detection of labral tears with 3.0-T MRI of the shoulder. *AJR Am J Roentgenol* 2006; 187:1448-1452.
36. O'Connor EE, Dixon LB, Peabody T, Stacy GS. MRI of cystic and soft-tissue masses of the shoulder joint. *AJR Am J Roentgenol* 2004; 183:39-47.
37. Stetson WB, Templin K. The crank test, the O'Brien test, and routine magnetic resonance imaging scans in the diagnosis of labral tears. *Am J Sports Med* 2002; 30:806-809.
38. Tuite MJ, Cirillo RL, De Smet AA, Orwin JF. Superior labrum anterior-posterior (SLAP) tears: evaluation of three MR signs on T2-weighted images. *Radiology* 2000; 215:841-845.
39. Tung GA, Entzian D, Stern JB, Green A. MR imaging and MR arthrography of paraglenoid labral cysts. *AJR Am J Roentgenol* 2000; 174:1707-1715.
40. Waldt S, Burkart A, Imhoff AB, Bruegel M, Rummeny EJ, Woertler K. Anterior shoulder instability: accuracy of MR arthrography in the classification of anteroinferior labroligamentous injuries. *Radiology* 2005; 237:578-583.
41. Waldt S, Burkart A, Lange P, Imhoff AB, Rummeny EJ, Woertler K. Diagnostic performance of MR arthrography in the assessment of superior labral anteroposterior lesions of the shoulder. *AJR Am J Roentgenol* 2004; 182:1271-1278.
42. Mengiardi B, Pfirrmann CW, Gerber C, Hodler J, Zanetti M. Frozen shoulder: MR arthrographic findings. *Radiology* 2004; 233:486-492.
43. Vinson EN, Major NM, Higgins LD. Magnetic resonance imaging findings associated with surgically proven rotator interval lesions. *Skeletal Radiol* 2007; 36:405-410.
44. Connell DA, Potter HG, Sherman MF, Wickiewicz TL. Injuries of the pectoralis major muscle: evaluation with MR imaging. *Radiology* 1999; 210:785-791.
45. Fuchs B, Weishaupt D, Zanetti M, Hodler J, Gerber C. Fatty degeneration of the muscles of the rotator cuff: assessment by computed tomography versus magnetic resonance imaging. *J Shoulder Elbow Surg* 1999; 8:599-605.
46. Huang CC, Ko SF, Ko JY, et al. Contracture of the deltoid muscle: sonographic evaluation with MRI correlation. *AJR Am J Roentgenol* 2005; 185:364-370.
47. Ogawa K, Takahashi M, Naniwa T. Deltoid contracture: MR imaging features. *Clin Radiol* 2001; 56:146-149.
48. Schaefer O, Winterer J, Lohrmann C, Laubenberger J, Reichelt A, Langer M. Magnetic resonance imaging for supraspinatus muscle atrophy after cuff repair. *Clin Orthop Relat Res* 2002:93-99.
49. Zanetti M, Gerber C, Hodler J. Quantitative assessment of the muscles of the rotator cuff with magnetic resonance imaging. *Invest Radiol* 1998; 33:163-170.
50. Zvijac JE, Schurhoff MR, Hechtman KS, Uribe JW. Pectoralis major tears: correlation of magnetic resonance imaging and treatment strategies. *Am J Sports Med* 2006; 34:289-294.
51. Carroll KW, Helms CA, Speer KP. Focal articular cartilage lesions of the superior humeral head: MR imaging findings in seven patients. *AJR Am J Roentgenol* 2001; 176:393-397.
52. Sanders TG, Tirman PF, Linares R, Feller JF, Richardson R. The glenolabral articular disruption lesion: MR arthrography with arthroscopic correlation. *AJR Am J Roentgenol* 1999; 172:171-175.
53. Yu JS, Greenway G, Resnick D. Osteochondral defect of the glenoid fossa: cross-sectional imaging features. *Radiology* 1998; 206:35-40.
54. Matsuzaki S, Yoneda M, Kobayashi Y, Fukushima S, Wakitani S. Dynamic enhanced MRI of the subacromial bursa: correlation with arthroscopic and histological findings. *Skeletal Radiol* 2003; 32:510-520.
55. Schraner AB, Major NM. MR imaging of the subcoracoid bursa. *AJR Am J Roentgenol* 1999; 172:1567-1571.
56. Sakai T, Sugano N, Nishii T, Hananouchi T, Yoshikawa H. Extent of osteonecrosis on MRI predicts humeral head collapse. *Clin Orthop Relat Res* 2008; 466:1074-1080.

57. Ritchie DA, Davies AM. MR imaging of tumors and tumor-like lesions of the shoulder girdle. *Magn Reson Imaging Clin N Am* 2004; 12:125-141, vii.
58. Kothari NA, Pelchovitz DJ, Meyer JS. Imaging of musculoskeletal infections. *Radiol Clin North Am* 2001; 39:653-671.
59. Struk DW, Munk PL, Lee MJ, Ho SG, Worsley DF. Imaging of soft tissue infections. *Radiol Clin North Am* 2001; 39:277-303.
60. Harper KW, Helms CA, Haystead CM, Higgins LD. Glenoid dysplasia: incidence and association with posterior labral tears as evaluated on MRI. *AJR Am J Roentgenol* 2005; 184:984-988.
61. Park YH, Lee JY, Moon SH, et al. MR arthrography of the labral capsular ligamentous complex in the shoulder: imaging variations and pitfalls. *AJR Am J Roentgenol* 2000; 175:667-672.
62. Tirman PF, Feller JF, Palmer WE, Carroll KW, Steinbach LS, Cox I. The Buford complex--a variation of normal shoulder anatomy: MR arthrographic imaging features. *AJR Am J Roentgenol* 1996; 166:869-873.
63. Tuite MJ, Blankenbaker DG, Seifert M, Ziegert AJ, Orwin JF. Sublabral foramen and buford complex: inferior extent of the unattached or absent labrum in 50 patients. *Radiology* 2002; 223:137-142.
64. Mochizuki T, Isoda H, Masui T, et al. Occlusion of the posterior humeral circumflex artery: detection with MR angiography in healthy volunteers and in a patient with quadrilateral space syndrome. *AJR Am J Roentgenol* 1994; 163:625-627.
65. Bredella MA, Tirman PF, Fritz RC, Wischer TK, Stork A, Genant HK. Denervation syndromes of the shoulder girdle: MR imaging with electrophysiologic correlation. *Skeletal Radiol* 1999; 28:567-572.
66. Hermann KG, Backhaus M, Schneider U, et al. Rheumatoid arthritis of the shoulder joint: comparison of conventional radiography, ultrasound, and dynamic contrast-enhanced magnetic resonance imaging. *Arthritis Rheum* 2003; 48:3338-3349.
67. Weishaupt D, Schweitzer ME. MR imaging of septic arthritis and rheumatoid arthritis of the shoulder. *Magn Reson Imaging Clin N Am* 2004; 12:111-124, vii.
68. Mason BJ, Kier R, Bindleglass DF. Occult fractures of the greater tuberosity of the humerus: radiographic and MR imaging findings. *AJR Am J Roentgenol* 1999; 172:469-473.
69. Saupé N, White LM, Bleakney R, et al. Acute traumatic posterior shoulder dislocation: MR findings. *Radiology* 2008; 248:185-193.
70. Giaroli EL, Major NM, Higgins LD. MRI of internal impingement of the shoulder. *AJR Am J Roentgenol* 2005; 185:925-929.
71. Giaroli EL, Major NM, Lemley DE, Lee J. Coracohumeral interval imaging in subcoracoid impingement syndrome on MRI. *AJR Am J Roentgenol* 2006; 186:242-246.
72. Tan V, Moore RS, Jr., Omarini L, Kneeland JB, Williams GR, Jr., Iannotti JP. Magnetic resonance imaging analysis of coracoid morphology and its relation to rotator cuff tears. *Am J Orthop* 2002; 31:329-333.
73. Tirman PF, Bost FW, Garvin GJ, et al. Posterosuperior glenoid impingement of the shoulder: findings at MR imaging and MR arthrography with arthroscopic correlation. *Radiology* 1994; 193:431-436.
74. Palmer WE, Caslowitz PL. Anterior shoulder instability: diagnostic criteria determined from prospective analysis of 121 MR arthrograms. *Radiology* 1995; 197:819-825.
75. Tung GA, Hou DD. MR arthrography of the posterior labrocapsular complex: relationship with glenohumeral joint alignment and clinical posterior instability. *AJR Am J Roentgenol* 2003; 180:369-375.
76. Willemsen UF, Wiedemann E, Brunner U, et al. Prospective evaluation of MR arthrography performed with high-volume intraarticular saline enhancement in patients with recurrent anterior dislocations of the shoulder. *AJR Am J Roentgenol* 1998; 170:79-84.
77. Beltran J, Kim DH. MR imaging of shoulder instability injuries in the athlete. *Magn Reson Imaging Clin N Am* 2003; 11:221-238.
78. Ho CP. MR imaging of rotator interval, long biceps, and associated injuries in the overhead-throwing athlete. *Magn Reson Imaging Clin N Am* 1999; 7:23-37.
79. Roger B, Skaf A, Hooper AW, Lektrakul N, Yeh L, Resnick D. Imaging findings in the dominant shoulder of throwing athletes: comparison of radiography, arthrography, CT arthrography, and MR arthrography with arthroscopic correlation. *AJR Am J Roentgenol* 1999; 172:1371-1380.
80. Schickendantz MS, Ho CP, Keppler L, Shaw BD. MR imaging of the thrower's shoulder. Internal impingement, latissimus dorsi/subscapularis strains, and related injuries. *Magn Reson Imaging Clin N Am* 1999; 7:39-49.
81. Major NM, Banks MC. MR imaging of complications of loose surgical tacks in the shoulder. *AJR Am J Roentgenol* 2003; 180:377-380.
82. Sugimoto H, Suzuki K, Mihara K, Kubota H, Tsutsui H. MR arthrography of shoulders after suture-anchor Bankart repair. *Radiology* 2002; 224:105-111.
83. Kanal E, Barkovich AJ, Bell C, et al. ACR guidance document for safe MR practices: 2007. *AJR Am J Roentgenol* 2007; 188:1447-1474.

84. American College of Radiology. Manual on Contrast Media. http://www.acr.org/SecondaryMainMenuCategories/quality_safety/contrast_manual.aspx. Accessed September 11, 2009.
85. Shellock FG. *Reference Manual for Magnetic Resonance Safety, Implants and Devices*. 2009 ed. Los Angeles, Calif: Biomedical Research Publishing Group; 2009.
86. Shellock FG, Crues JV. MR procedures: biologic effects, safety, and patient care. *Radiology* 2004; 232:635-652.
87. Kreitner KF, Loew R, Runkel M, Zollner J, Thelen M. Low-field MR arthrography of the shoulder joint: technique, indications, and clinical results. *Eur Radiol* 2003; 13:320-329.
88. Merl T, Scholz M, Gerhardt P, et al. Results of a prospective multicenter study for evaluation of the diagnostic quality of an open whole-body low-field MRI unit. A comparison with high-field MRI measured by the applicable gold standard. *Eur J Radiol* 1999; 30:43-53.
89. Sasaki M, Ehara S, Nakasato T, et al. MR of the shoulder with a 0.2-T permanent-magnet unit. *AJR Am J Roentgenol* 1990; 154:777-778.
90. Shellock FG, Bert JM, Fritts HM, Gundry CR, Easton R, Crues JV, 3rd. Evaluation of the rotator cuff and glenoid labrum using a 0.2-Tesla extremity magnetic resonance (MR) system: MR results compared to surgical findings. *J Magn Reson Imaging* 2001; 14:763-770.
91. Zlatkin MB, Hoffman C, Shellock FG. Assessment of the rotator cuff and glenoid labrum using an extremity MR system: MR results compared to surgical findings from a multi-center study. *J Magn Reson Imaging* 2004; 19:623-631.
92. Erickson SJ. High-resolution imaging of the musculoskeletal system. *Radiology* 1997; 205:593-618.
93. Loew R, Kreitner KF, Runkel M, Zoellner J, Thelen M. MR arthrography of the shoulder: comparison of low-field (0.2 T) vs high-field (1.5 T) imaging. *Eur Radiol* 2000; 10:989-996.
94. Tung GA, Entzian D, Green A, Brody JM. High-field and low-field MR imaging of superior glenoid labral tears and associated tendon injuries. *AJR Am J Roentgenol* 2000; 174:1107-1114.
95. Allmann KH, Walter O, Laubenberger J, et al. Magnetic resonance diagnosis of the anterior labrum and capsule. Effect of field strength on efficacy. *Invest Radiol* 1998; 33:415-420.
96. Magee T, Shapiro M, Williams D. Comparison of high-field-strength versus low-field-strength MRI of the shoulder. *AJR Am J Roentgenol* 2003; 181:1211-1215.
97. Glickstein MF. MR imaging of the shoulder: optimizing surface-coil positioning. *AJR Am J Roentgenol* 1989; 153:431-432.
98. Tuite MJ, De Smet AA, Norris MA, Orwin JF. MR diagnosis of labral tears of the shoulder: value of T2*-weighted gradient-recalled echo images made in external rotation. *AJR Am J Roentgenol* 1995; 164:941-944.
99. Farley TE, Neumann CH, Steinbach LS, Jahnke AJ, Petersen SS. Full-thickness tears of the rotator cuff of the shoulder: diagnosis with MR imaging. *AJR Am J Roentgenol* 1992; 158:347-351.
100. Davis SJ, Teresi LM, Bradley WG, Ressler JA, Eto RT. Effect of arm rotation on MR imaging of the rotator cuff. *Radiology* 1991; 181:265-268.
101. Flannigan B, Kursunoglu-Brahme S, Snyder S, Karzel R, Del Pizzo W, Resnick D. MR arthrography of the shoulder: comparison with conventional MR imaging. *AJR Am J Roentgenol* 1990; 155:829-832.
102. Saleem AM, Lee JK, Novak LM. Usefulness of the abduction and external rotation views in shoulder MR arthrography. *AJR Am J Roentgenol* 2008; 191:1024-1030.
103. Herold T, Bachthaler M, Hamer OW, et al. Indirect MR arthrography of the shoulder: use of abduction and external rotation to detect full- and partial-thickness tears of the supraspinatus tendon. *Radiology* 2006; 240:152-160.
104. Lee SY, Lee JK. Horizontal component of partial-thickness tears of rotator cuff: imaging characteristics and comparison of ABER view with oblique coronal view at MR arthrography initial results. *Radiology* 2002; 224:470-476.
105. Tirman PF, Bost FW, Steinbach LS, et al. MR arthrographic depiction of tears of the rotator cuff: benefit of abduction and external rotation of the arm. *Radiology* 1994; 192:851-856.
106. Patten RM, Spear RP, Richardson ML. Diagnostic performance of magnetic resonance imaging for the diagnosis of rotator cuff tears using supplemental images in the oblique sagittal plane. *Invest Radiol* 1994; 29:87-93.
107. Tuite MJ, Asinger D, Orwin JF. Angled oblique sagittal MR imaging of rotator cuff tears: comparison with standard oblique sagittal images. *Skeletal Radiol* 2001; 30:262-269.
108. Munk PL, Holt RG, Helms CA, Genant HK. Glenoid labrum: preliminary work with use of radial-sequence MR imaging. *Radiology* 1989; 173:751-753.
109. Kwak SM, Brown RR, Trudell D, Resnick D. Glenohumeral joint: comparison of shoulder positions at MR arthrography. *Radiology* 1998; 208:375-380.
110. Kneeland JB, Shimakawa A, Wehrli FW. Effect of intersection spacing on MR image contrast and study time. *Radiology* 1986; 158:819-822.
111. Rubin DA, Kneeland JB. MR imaging of the musculoskeletal system: technical considerations

- for enhancing image quality and diagnostic yield. *AJR Am J Roentgenol* 1994; 163:1155-1163.
112. Carrino JA, McCauley TR, Katz LD, Smith RC, Lange RC. Rotator cuff: evaluation with fast spin-echo versus conventional spin-echo MR imaging. *Radiology* 1997; 202:533-539.
 113. Quinn SF, Sheley RC, Demlow TA, Szumowski J. Rotator cuff tendon tears: evaluation with fat-suppressed MR imaging with arthroscopic correlation in 100 patients. *Radiology* 1995; 195:497-500.
 114. Sonin AH, Peduto AJ, Fitzgerald SW, Callahan CM, Bresler ME. MR imaging of the rotator cuff mechanism: comparison of spin-echo and turbo spin-echo sequences. *AJR Am J Roentgenol* 1996; 167:333-338.
 115. Parsa M, Tuite M, Norris M, Orwin J. MR imaging of rotator cuff tendon tears: comparison of T2*-weighted gradient-echo and conventional dual-echo sequences. *AJR Am J Roentgenol* 1997; 168:1519-1524.
 116. Sahin-Akyar G, Miller TT, Staron RB, McCarthy DM, Feldman F. Gradient-echo versus fat-suppressed fast spin-echo MR imaging of rotator cuff tears. *AJR Am J Roentgenol* 1998; 171:223-227.
 117. McCauley TR, Pope CF, Jokl P. Normal and abnormal glenoid labrum: assessment with multiplanar gradient-echo MR imaging. *Radiology* 1992; 183:35-37.
 118. Bush CH. The magnetic resonance imaging of musculoskeletal hemorrhage. *Skeletal Radiol* 2000; 29:1-9.
 119. De Smet AA, Fisher DR, Heiner JP, Keene JS. Magnetic resonance imaging of muscle tears. *Skeletal Radiol* 1990; 19:283-286.
 120. Hodler J, Kursunoglu-Brahme S, Snyder SJ, et al. Rotator cuff disease: assessment with MR arthrography versus standard MR imaging in 36 patients with arthroscopic confirmation. *Radiology* 1992; 182:431-436.
 121. Palmer WE, Brown JH, Rosenthal DI. Rotator cuff: evaluation with fat-suppressed MR arthrography. *Radiology* 1993; 188:683-687.
 122. Dinauer PA, Flemming DJ, Murphy KP, Doukas WC. Diagnosis of superior labral lesions: comparison of noncontrast MRI with indirect MR arthrography in unexercised shoulders. *Skeletal Radiol* 2007; 36:195-202.
 123. Kijowski R, Farber JM, Medina J, Morrison W, Ying J, Buckwalter K. Comparison of fat-suppressed T2-weighted fast spin-echo sequence and modified STIR sequence in the evaluation of the rotator cuff tendon. *AJR Am J Roentgenol* 2005; 185:371-378.
 124. Rybicki FJ, Chung T, Reid J, Jaramillo D, Mulkern RV, Ma J. Fast three-point dixon MR imaging using low-resolution images for phase correction: a comparison with chemical shift selective fat suppression for pediatric musculoskeletal imaging. *AJR Am J Roentgenol* 2001; 177:1019-1023.
 125. Chan KK, Muldoon KA, Yeh L, et al. Superior labral anteroposterior lesions: MR arthrography with arm traction. *AJR Am J Roentgenol* 1999; 173:1117-1122.
 126. Peh WC, Chan JH. Artifacts in musculoskeletal magnetic resonance imaging: identification and correction. *Skeletal Radiol* 2001; 30:179-191.
 127. McGee KP, Grimm RC, Felmler JP, Rydberg JR, Riederer SJ, Ehman RL. The shoulder: adaptive motion correction of MR images. *Radiology* 1997; 205:541-545.
 128. Haacke EM, Lenz GW. Improving MR image quality in the presence of motion by using rephasing gradients. *AJR Am J Roentgenol* 1987; 148:1251-1258.
 129. Patten RM. Vacuum phenomenon: a potential pitfall in the interpretation of gradient-recalled-echo MR images of the shoulder. *AJR Am J Roentgenol* 1994; 162:1383-1386.
 130. Timins ME, Erickson SJ, Estkowski LD, Carrera GF, Komorowski RA. Increased signal in the normal supraspinatus tendon on MR imaging: diagnostic pitfall caused by the magic-angle effect. *AJR Am J Roentgenol* 1995; 165:109-114.
 131. Schulte-Altendorneburg G, Gebhard M, Wohlgemuth WA, et al. MR arthrography: pharmacology, efficacy and safety in clinical trials. *Skeletal Radiol* 2003; 32:1-12.
 132. Shellock FG, Stoller D, Crues JV. MRI of the shoulder: a rational approach to the reporting of findings. *J Magn Reson Imaging* 1996; 6:268-270.
 133. Shellock FG. *Guide to MR Procedures and Metallic Objects: Update 2001*. 7th ed. Philadelphia, Pa: Lippincott, Williams & Wilkins; 2001.
 134. Sawyer-Glover AM, Shellock FG. Pre-MRI procedure screening: recommendations and safety considerations for biomedical implants and devices. *J Magn Reson Imaging* 2000; 12:92-106.
-
- *Guidelines and standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For guidelines and standards published before 1999, the effective date was January 1 following the year in which the guideline or standard was amended, revised, or approved by the ACR Council.
- Development Chronology for this Guideline
 2005 (Resolution 10)
 Amended 2006 (Resolution 35)
 Revised 2010 (Resolution 17)