

The American College of Radiology, with more than 30,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice guidelines and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice guidelines and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice guideline and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review, requiring the approval of the Commission on Quality and Safety as well as the ACR Board of Chancellors, the ACR Council Steering Committee, and the ACR Council. The practice guidelines and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice guideline and technical standard by those entities not providing these services is not authorized.

Revised 2007 (Resolution 24)*

ACR–AIUM PRACTICE GUIDELINE FOR THE PERFORMANCE OF AN ULTRASOUND EXAMINATION OF THE ABDOMEN AND/OR RETROPERITONEUM

PREAMBLE

These guidelines are an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. They are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care. For these reasons and those set forth below, the American College of Radiology cautions against the use of these guidelines in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the physician or medical physicist in light of all the circumstances presented. Thus, an approach that differs from the guidelines, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in the guidelines when, in the reasonable judgment of the practitioner, such course of action is indicated by the condition of the patient, limitations of available resources, or advances in knowledge or technology subsequent to publication of the guidelines. However, a practitioner who employs an approach substantially different from these guidelines is advised to document in the patient record information sufficient to explain the approach taken.

The practice of medicine involves not only the science, but also the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment.

Therefore, it should be recognized that adherence to these guidelines will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The sole purpose of these guidelines is to assist practitioners in achieving this objective.

I. INTRODUCTION

The clinical aspects contained in specific sections of this guideline (Introduction, Indications, Specifications of the Examination, and Equipment Specifications) were developed collaboratively by the American College of Radiology (ACR) and the American Institute of Ultrasound in Medicine (AIUM). Recommendations for physician requirements, written request for the examination, procedure documentation, and quality control vary between the two organizations and are addressed by each separately.

This guideline has been developed to assist practitioners performing ultrasound studies of the abdomen and/or retroperitoneum. Ultrasound examination is a proven and useful procedure for the evaluation of many structures within these anatomic areas. Depending on the clinical indications, an examination may include the entirety of the abdomen and/or retroperitoneum, a single organ, or several organs. A combination of structures may be imaged because of location (e.g., upper abdominal scan, right upper quadrant organs) or function (e.g., biliary system [liver, gallbladder, and bile ducts], both kidneys). For some patients, more focused examinations may be

appropriate for evaluating specific clinical indications or to follow up a known abnormality. In some cases, additional and/or specialized examinations may be necessary (e.g., spectral, color, and/or power Doppler). While it is not possible to detect every abnormality using ultrasound examination of the abdomen and/or retroperitoneum, adherence to the following guideline will maximize the probability of detecting abnormalities.

Throughout this guideline, references to Doppler evaluation may include spectral, color, or power Doppler individually or in any combination. Whenever a long axis view is indicated, it could be either a sagittal or coronal plane.

(For pediatric considerations, see sections V.A.2, V.A.5, V.B.3, and VII.)

II. QUALIFICATIONS AND RESPONSIBILITIES OF THE PHYSICIAN

Each organization addresses this requirement individually. ACR language is as follows:

See the [ACR–SPR–SRU Practice Guideline for Performing and Interpreting Diagnostic Ultrasound Examinations](#).

III. INDICATIONS/CONTRAINDICATIONS

Indications for ultrasound examination of the abdomen and/or retroperitoneum include, but are not limited to:

- A. Abdominal, flank, and/or back pain.
- B. Signs or symptoms that may be referred from the abdominal and/or retroperitoneal regions such as jaundice or hematuria.
- C. Palpable abnormalities such as an abdominal mass or organomegaly.
- D. Abnormal laboratory values or abnormal findings on other imaging examinations suggestive of abdominal and/or retroperitoneal pathology.
- E. Follow-up of known or suspected abnormalities in the abdomen and/or retroperitoneum.
- F. Search for metastatic disease or occult primary neoplasm.
- G. Evaluation of suspected congenital abnormalities.
- H. Abdominal trauma.
- I. Pretransplantation and post-transplantation evaluation.
- J. Planning and guidance for an invasive procedure.
- K. Search for the presence of free or loculated peritoneal and/or retroperitoneal fluid.

Abdominal and/or retroperitoneal ultrasound should be performed when there is a valid medical reason. There are no absolute contraindications.

IV. WRITTEN REQUEST FOR THE EXAMINATION

Each organization addresses this requirement individually. ACR language is as follows:

The written or electronic request for an abdomen and/or retroperitoneum ultrasound examination should provide sufficient information to demonstrate the medical necessity of the examination and allow for its proper performance and interpretation.

Documentation that satisfies medical necessity includes 1) signs and symptoms and/or 2) relevant history (including known diagnoses). Additional information regarding the specific reason for the examination or a provisional diagnosis would be helpful and may at times be needed to allow for the proper performance and interpretation of the examination.

The request for the examination must be originated by a physician or other appropriately licensed health care provider. The accompanying clinical information should be provided by a physician or other appropriately licensed health care provider familiar with the patient's clinical problem or question and consistent with the state's scope of practice requirements. (ACR Resolution 35, adopted in 2006)

V. SPECIFICATIONS FOR INDIVIDUAL EXAMINATIONS

Doppler may be useful to differentiate vascular from nonvascular structures in any location. Measurements should be considered for any abnormal area.

1. Liver

The examination of the liver should include long axis and transverse views. The liver parenchyma should be evaluated for focal and/or diffuse abnormalities. If possible, the echogenicity of the liver should be compared with that of the right kidney. In addition, the following should be imaged:

- a. The major vessels in the region of the liver, including the inferior vena cava (IVC), the hepatic veins, the main portal vein, and, if possible, the right and left branches of the portal vein.
- b. The hepatic lobes (right, left, and caudate) and, if possible, the right hemidiaphragm and the adjacent pleural space.
- c. For vascular examinations of the native or transplanted liver, Doppler evaluation should be used to document blood flow characteristics and blood flow direction. The structures that may be examined include the hepatic arteries, hepatic veins, portal veins, intrahepatic portion of the IVC, collateral

venous pathways, and transjugular intrahepatic portosystemic shunt (TIPS) stents.

2. Gallbladder and biliary tract

The gallbladder evaluation should include long-axis and transverse views obtained in the supine position; other positions, such as left lateral decubitus, erect, or prone positions, may be necessary to evaluate the gallbladder and its surrounding area completely, particularly when stones and/or sludge are observed. Measurements may aid in determining gallbladder wall thickening. If the patient presents with pain, tenderness to transducer compression should be assessed.

The intrahepatic ducts can be evaluated by obtaining views of the liver demonstrating the right and left branches of the portal vein. Doppler may be used to differentiate hepatic arteries and portal veins from bile ducts. The intrahepatic and extrahepatic bile ducts should be evaluated for dilatation, wall thickening, intraluminal findings, and other abnormalities. The size of the bile duct in the porta hepatis should be documented. When visualized, the distal common bile duct in the pancreatic head should be evaluated.

Routine gallbladder examination should be conducted on an adequately distended gallbladder. In most cases, fasting for 8 hours prior to examination will permit adequate distension of a normally functioning gallbladder in adults and children. In infants and some adults, adequate distention may be achieved in less time.

3. Pancreas

Whenever possible, all portions of the pancreas – head, uncinate process, body, and tail – should be identified. Orally administered water or contrast agent may afford better visualization of the pancreas. The following should be assessed in the examination of the pancreas:

- a. Parenchymal abnormalities.
- b. The distal common bile duct in the region of the pancreatic head.
- c. The pancreatic duct for dilatation and any other abnormalities, with dilatation confirmed by measurement.
- d. The peripancreatic region for adenopathy and/or fluid.

4. Spleen

Representative views of the spleen in long-axis and transverse projections should be obtained. Doppler may be used to determine the presence and direction of flow in the splenic vein and artery. Splenic measurement may be helpful in assessing enlargement. Echogenicity of the left kidney should be compared to splenic echogenicity when possible. An attempt should be made to demonstrate the left hemidiaphragm and the adjacent pleural space.

5. Bowel

The bowel may be evaluated for wall thickening, dilatation, muscular hypertrophy, masses, and other abnormalities. Sonography of the pylorus and surrounding structures may be indicated in the evaluation of the vomiting infant. Compression sonography may be necessary to visualize the appendix or other bowel loops. Measurements may aid in determining bowel wall thickening. Color Doppler imaging may be done to assess blood flow.

6. Peritoneal fluid

Evaluation for free or loculated peritoneal fluid should include documentation of the extent and location of any fluid identified.

For evaluating peritoneal spaces for bleeding after traumatic injury, particularly blunt trauma, the examination known as focused abdominal sonographic examination for trauma (FAST) assessment (or focused assessment with sonography for trauma) may be performed.

The objective of the abdominal portion of the examination is to analyze the abdomen for free fluid. Longitudinal and transverse plane images should be obtained in the right upper quadrant through the area of the liver with attention to fluid collections peripheral to the liver and in the subhepatic space. Longitudinal and transverse plane images should be obtained in the left upper quadrant through the area of the spleen, with attention to fluid collections peripheral to the spleen. Longitudinal and transverse images should be obtained at the periphery of the left and right abdomen in the areas of the left and right paracolic gutters for evidence of free fluid. Longitudinal and transverse midline images of the pelvis are obtained to evaluate for free pelvic fluid. Analysis through a fluid filled bladder (which if necessary can be filled through a Foley catheter, when possible) may help in the evaluation of the pelvis.

7. **Abdominal wall**
The examination should include images of the abdominal wall in the location of symptoms or signs. The relationship of any identified mass to the peritoneum should be demonstrated. Any defect in the peritoneum and abdominal wall musculature should be documented. The presence or absence of bowel, fluid, or other tissue contained within any abdominal wall defect should be noted. Images obtained in upright position and/or with use of the Valsalva maneuver may be helpful. Doppler examination may be useful to define the relationship of blood vessels to a detected mass.

8. **Kidneys**
The examination should include long-axis and transverse views of the upper poles, mid portions, and lower poles of the kidneys. The cortices and renal pelves should be assessed. A maximum measurement of renal length should be recorded for both kidneys. Decubitus, prone, or upright positioning may provide better images of the kidneys. When possible, renal echogenicity should be compared to the adjacent liver or spleen. The kidneys and perirenal regions should be assessed for abnormalities.

For vascular examination of the kidneys, Doppler can be used

- a. To assess renal arterial and venous patency.
- b. To evaluate adults suspected of having renal artery stenosis. For this application, angle-adjusted measurements of the peak systolic velocity should be made proximally, centrally, and distally in the extrarenal portion of the main renal arteries when possible. Peak systolic velocity of the adjacent aorta (or iliac artery in transplanted kidneys) should also be documented for calculating the ratio of renal to aortic peak systolic velocity. Spectral Doppler evaluation of the intrarenal arteries from the upper and lower portions of the kidneys, obtained to evaluate the early systolic peak, may be of value as indirect evidence of proximal stenosis in the main renal artery.

9. **Urinary bladder and adjacent structures**
When performing a complete ultrasound evaluation of the urinary tract, transverse and longitudinal images of the distended urinary bladder and its wall should be included, if possible. Bladder lumen or wall abnormalities should be noted. Dilatation or other distal ureteral abnormalities should be documented. Transverse and longitudinal scans may be used

to demonstrate any postvoid residual, which may be quantitated and reported.

10. **Adrenal glands**
When possible, usually in the newborn or young infant, long-axis and transverse images of the adrenal glands may be obtained. The adrenal glands are infrequently seen in adults. When visualized, the size of the gland should be documented as well as the presence of hemorrhage, masses, or other abnormalities.

11. **Aorta**
Representative images of the aorta in longitudinal and transverse planes should be obtained as appropriate. When evaluation of the aorta is specifically requested, see the [ACR–AIUM–SRU Practice Guideline for the Performance of Diagnostic and Screening Ultrasound of the Abdominal Aorta](#).

12. **Inferior vena cava**
Transverse and longitudinal images of the IVC should be obtained. Patency and abnormalities may be evaluated with Doppler. Vena cava filters, interruption devices, or catheters may need to be localized with respect to the hepatic and/or renal veins.

VI. DOCUMENTATION

Each organization addresses this requirement individually. ACR language is as follows:

Adequate documentation is essential for high-quality patient care. There should be a permanent record of the ultrasound examination and its interpretation. Comparison with prior relevant imaging studies may prove helpful. Images of all appropriate areas, both normal and abnormal, should be recorded. Variations from normal size should generally be accompanied by measurements. Images should be labeled with the patient identification, facility identification, examination date, and image orientation. An official interpretation (final report) of the ultrasound examination should be included in the patient's medical record. Retention of the ultrasound examination images should be consistent both with clinical need and with relevant legal and local health care facility requirements.

Reporting should be in accordance with the [ACR Practice Guideline for Communication of Diagnostic Imaging Findings](#).

VII. EQUIPMENT SPECIFICATIONS

Abdomen and/or retroperitoneum sonographic studies should be conducted with real-time scanners, preferably

using sector or linear (straight or curved) transducers. The equipment should be adjusted to operate at the highest clinically appropriate frequency, realizing that there is a trade-off between resolution and beam penetration. For most preadolescent pediatric patients, mean frequencies of 5 MHz or greater are preferred, and in newborns and small infants a higher frequency transducer is often necessary. For adults, mean frequencies between 2 and 5 MHz are most commonly used. When Doppler studies are performed, the Doppler frequency may differ from the imaging frequency. Diagnostic information should be optimized, while keeping total ultrasound exposure as low as reasonably achievable.

VIII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

Each organization addresses this requirement individually. ACR language is as follows:

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *Position Statement on QC & Improvement, Safety, Infection Control, and Patient Education* on the ACR web page (<http://www.acr.org/guidelines>).

Equipment performance monitoring should be in accordance with the [ACR Technical Standard for Diagnostic Medical Physics Performance Monitoring of Real Time Ultrasound Equipment](#).

ACKNOWLEDGEMENTS

This guideline was revised according to the process described under the heading *The Process for Developing ACR Practice Guidelines and Technical Standards* on the ACR web page (<http://www.acr.org/guidelines>) by the ACR Guidelines and Standards Committee of the Commission on Ultrasound in collaboration with the AIUM.

Principal Revisers

Harris L. Cohen, MD
John P. McGahan, MD

Collaborative Subcommittee

ACR

Beverly E. Hashimoto, MD, Chair
W. Dennis Foley, MD
Ronald R. Townsend, MD

AIUM

Barbara S. Hertzberg, MD
Jon W. Meilstrup, MD
Laurence Needleman, MD

ACR Guidelines and Standards Committee

Gretchen A. Gooding, MD, Chair
Raymond E. Bertino, MD
Mary C. Frates, MD
Ruth B. Goldstein, MD
Beatrice L. Madrazo, MD
Jon W. Meilstrup, MD
Michelle L. Melany, MD
Miriam N. Mikhail, MD
Sara M. O'Hara, MD
Suhass G. Parulekar, MD
John S. Pellerito, MD
Philip W. Ralls, MD
Michelle L. Robbin, MD
Carol M. Rumack, MD, Chair, Commission

Comments Reconciliation Committee

Julie K. Timins, MD, Co-Chair
Bill H. Warren, MD, Co-Chair
Albert L. Blumberg, MD
Harris L. Cohen, MD
Mary C. Frates, MD
Gretchen A. Gooding, MD
Beverly E. Hashimoto, MD
David C. Kushner, MD
Paul A. Larson, MD
Lawrence A. Liebscher, MD
Beatrice L. Madrazo, MD
John P. McGahan, MD
Jon W. Meilstrup, MD
Matthew S. Pollack, MD
Carol M. Rumack, MD

Suggested Reading (Additional articles that are not cited in the document but that the committee recommends for further reading on this topic)

1. Babcock DS, Patriquin HB. The pediatric kidney and adrenal glands. In: Rumack CM, Wilson SR, Charboneau JW, et al, eds. *Diagnostic Ultrasound*. 3rd edition. Philadelphia Pa: Elsevier Mosby; 2005:1905-1940.
2. Baxter GM. Imaging in renal transplantation. *Ultrasound Q* 2003;19:123-138.
3. Desser TS, Sze DY, Jeffrey RB. Imaging and intervention in the hepatic veins. *AJR* 2003;180:1583-1591.
4. Gandolfi L, Torresan F, Solmi L, Pucetti A. The role of ultrasound in biliary and pancreatic diseases. *Eur J Ultrasound* 2003;16:141-159.
5. Guimaraes CM, Correia MM, Baldisserott M, de Queiroz Aires EP, Coelho JF. Intraoperative ultrasonography of the liver in patients with

- abdominal tumors: a new approach. *J Ultrasound Med* 2004;23:1549-1555.
6. Hagen-Ansert SL. Introduction to abdominal scanning techniques and protocols. In: Hagen-Ansert SL, ed. *Textbook of Diagnostic Ultrasonography*. 5th edition. St. Louis, Mo: Mosby; 2001:42-69.
 7. Hagen-Ansert SL. The spleen. In: Hagen-Ansert SL, ed. *Textbook of Diagnostic Ultrasonography*. 5th edition. St. Louis, Mo: Mosby; 2001:308-326.
 8. Ho LM, Thomas J, Fine SA, Paulson EK. Usefulness of sonographic guidance during percutaneous biopsy of mesenteric masses. *AJR* 2003;180:1563-1566.
 9. Jeffrey RB. Right lower-quadrant pain: rule out appendicitis. In: Bluth EI, Arger PH, Benson CB, et al, eds. *Ultrasound: A Practical Approach to Clinical Problems*. New York, NY: Thieme; 2000:125-132.
 10. Kim NR, Kim SH, Lee JM, et al. Sonographic features of an intraductal polypoid mass: differentiation between hepatocellular carcinoma and intraductal cholangiocarcinoma. *J Ultrasound Med* 2004;23:1283-1291.
 11. Kitamura H, Kobayashi C. Impairment of change in diameter of the hepatic portion of the inferior vena cava: a sonographic sign of liver fibrosis or cirrhosis. *J Ultrasound Med* 2005;24:355-359.
 12. Kimura A, Otsuka T. Emergency center ultrasonography in the evaluation of hemoperitoneum: a prospective study. *J Trauma* 1991;31:20-23.
 13. Koito K, Namieno T, Nagakawa T, et al. Pancreas: imaging diagnosis with color/power Doppler ultrasonography, endoscopic ultrasonography, and intraductal ultrasonography. *Eur J Radiol* 2001;38:94-104.
 14. Laing FC. Jaundice. In: Bluth EI, Arger PH, Benson CB, et al, eds. *Ultrasound: A Practical Approach to Clinical Problems*. New York, NY: Thieme; 2000:17-29.
 15. Lingawi SS, Buckley AR. Focused abdominal US in patients with trauma. *Radiology* 2000;217:426-429.
 16. McGahan J, Richard J, Fogata M. Emergency ultrasound in trauma patients. *Radiol Clin N Am* 2004;42:417-425.
 17. McGahan JP, Rose J, Coates TL, Wisner DH, Newberry P. Use of ultrasonography in the patient with acute abdominal trauma. *J Ultrasound Med* 1997;16:653-662.
 18. Middleton WD. Right upper quadrant pain. In: Bluth EI, Arger PH, Benson CB, et al, eds. *Ultrasound: A Practical Approach to Clinical Problems*. New York, NY: Thieme; 2000:3-16.
 19. Middleton WD, Kurtz AB, Hertzberg BS. Kidney. In: *Ultrasound: The Requisites*. 2nd edition. St. Louis, Mo: Mosby; 2004.
 20. Muradali D, Wilson SR. Organ transplantation. In: Rumack CM, Wilson SR, Charboneau JW, et al, eds. *Diagnostic Ultrasound*. 3rd edition. Philadelphia, Pa: Elsevier Mosby; 2005:675-704.
 21. Radermacher J, Chavan J, Bleck J, et al. Use of Doppler ultrasonography to predict the outcome of therapy for renal-artery stenosis. *N Engl J Med* 2001;344:410-417.
 22. Ralls PW. Hyperamylasemia. In: Bluth EI, Arger PH, Benson CB, et al, eds. *Ultrasound: A Practical Approach to Clinical Problems*. New York, NY: Thieme; 2000:67-75.
 23. Ralls PW. Ultrasound to assess respectability of periampullary neoplasms. In: Bluth EI, Arger PH, Benson CB, et al, eds. *Ultrasound: A Practical Approach to Clinical Problems*. New York, NY: Thieme; 2000:58-66.
 24. Robbin ML, van Leeuwen DJ. Sonography in the evaluation of abnormal liver tests. In: Bluth EI, Arger PH, Benson CB, et al, eds. *Ultrasound: A Practical Approach to Clinical Problems*. New York, NY: Thieme; 2000:30-45.
 25. Scott T. Sonographic detection of the normal appendix: does the combination of real-time spatial compounding and tissue harmonic imaging make a difference? *J of Diagnostic Med Sonography* 2004;5:326-334.
 26. Siegel MJ. Liver. In: *Pediatric Sonography*. 3rd edition. Philadelphia, Pa: Lippincott, Williams and Wilkins; 2001.
 27. Siegel MJ. Gallbladder and biliary tract. In: *Pediatric Sonography*. 3rd edition. Philadelphia, Pa: Lippincott, Williams and Wilkins; 2001.
 28. Siegel MJ. Urinary tract. In: *Pediatric Sonography*. 3rd edition. Philadelphia, Pa: Lippincott, Williams and Wilkins; 2001.
 29. Sivit C, Siegel MJ. Gastrointestinal tract. In: Siegel MJ, ed. *Pediatric Sonography*. 3rd edition. Philadelphia, Pa: Lippincott, Williams and Wilkins; 2001.
 30. Sivit C, Siegel MJ. Spleen and peritoneal cavity. In: Siegel MJ, ed. *Pediatric Sonography*. 3rd edition. Philadelphia, Pa: Lippincott, Williams and Wilkins; 2001.
 31. Sofka CM. Multicentre aneurysm screening study (MASS): cost effectiveness analysis of screening for abdominal aortic aneurysms based on four-year results from randomized controlled trial. *Ultrasound Quarterly* 2003;19:106-107.
 32. Soudack M, Epelman M, Maor R, et al. Experience with focused abdominal sonography for trauma (FAST) in 313 pediatric patients. *J Clin Ultrasound* 2004;32:53-61.
 33. Thurston W, Wilson SR. The urinary tract. In: Rumack CM, Wilson SR, Charboneau JW, et al, eds. *Diagnostic Ultrasound*. 3rd edition. Philadelphia, Pa: Elsevier Mosby; 2005:321-394.

*Guidelines and standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For guidelines and standards published before 1999, the effective date

was January 1 following the year in which the guideline or standard was amended, revised, or approved by the ACR Council.

Development Chronology for this Guideline

1990 (Resolution 7)

Revised 1993 (Resolution 4)

Revised 2001 (Resolution 36)

Revised 2006 (Resolution 39, 35)

Revised 2007 (Resolution 24)