

# ACR Lung Cancer Screening Economics & Billing Quick Reference Guide



This guide is intended to answer commonly asked questions about lung cancer screening logistics, program requirements, economics, and billing issues.

| Definitions                        |   |
|------------------------------------|---|
| Chest CT                           | Generic term encompassing low-dose and non-low-dose CT, with or without IV contrast.  |
| LDCT                               | Chest CT using low-dose technique—an imaging technique used to evaluate the chest primarily for lung nodule detection, follow up, and lung disease. The LDCT (low-dose chest CT) technique is required for lung cancer screening.   |
| Interval CT                        | Short interval chest CT or CTA performed “off-cycle” between annual screening exams for diagnostic purposes. In most cases, performed with LDCT technique.  |
| Surveillance CT                    | Diagnostic chest CT performed for surveillance in patients with known lung cancer, generally evaluating for recurrent cancer.   |
| Program specifics                  |   |
| Site accreditation requirements    | <p><b>For CT:</b> CT accreditation is required for non-hospital-based outpatient facilities that bill for CT under part B of the Medicare physician fee schedule.</p> <p><b>For ACR® Lung Cancer Screening Center Designation:</b> Facility must have ACR CT Accreditation and participate in the ACR Lung Cancer Screening (LCS) Registry.</p> <p><b>For ACR LCS Registry participation:</b> ACR CT Accreditation is not required.</p> |
| Ordering provider                  | An order for LCS LDCT must be provided by a licensed independent practitioner. Order may be electronic or paper-based.  |
| Shared decision making             | <p>Face-to-face discussion by a licensed independent practitioner (physician, NP, and PA) (required by Medicare for first-time screen only); however, this discussion can be performed and reimbursed multiple times per year.</p> <p>Can be performed as part of the Evaluation and Management (E&amp;M) visit and billed with a 25 modifier. Optional but reimbursable (by Medicare) for subsequent annual LCS CT.</p>                |
| Smoking cessation intervention     | Smoking cessation interventions and services must be offered to current smokers. If smoking cessation counseling is provided, it must be documented separately. Smoking cessation counseling can be reported in addition to an E&M visit, performed on the same day by the same licensed independent provider, by appending 25 modifier to the appropriate level of E&M service.  |
| Lung-RADS®                         | <p>Used for interpreting low-dose LCS CT exams and interval follow-up CTs for screen-detected abnormalities that may be lung cancer.</p> <p>If an interval CT is performed for another reason, Lung-RADS® use is recommended when possible for the lung nodule-specific findings.</p> <p>Use is required for participation in the ACR Lung Cancer Screening Registry™.</p>  |
| ACR Lung Cancer Screening Registry | <p>Submission of data to a CMS-approved registry is required for CMS coverage of LCS CT. The ACR Lung Cancer Screening Registry is currently the only CMS-approved registry.</p> <p>Submission of data on all screening exams is required regardless of insurance coverage, i.e., not just Medicare patients. The ACR submits a subset of data to CMS.</p>  |

| <b>LCS coverage and eligibility criteria</b> |   |
|--|---|
| Medicare                                     | LCS is covered as a preventive service in patients aged 55–77 years; ≥30 pack-year smoking history; current smokers or quit within last 15 years; no signs or symptoms of lung cancer.  |
| Medicaid                                     | Varies, state-dependent. If covered, typically Medicare-like criteria. Verify specifics for your state.   |
| Commercial                                   | LCS is covered as a preventive service for patients who meet U.S. Preventive Services Task Force criteria—ages 55–80; ≥30 pack-years smoking history; current smoker or quit within last 15 years; no signs or symptoms of lung cancer. |
| Self-pay                                     | Permitted for patients who do not meet standard criteria but fulfill National Comprehensive Cancer Network group 2 or other high-risk criteria.<br><br>Coded as LCS LDCT (G0297); submission to the ACR LCS Registry is required.       |
| IDTF setting                                 | LCS LDCT is covered in an Independent Diagnostic Testing Facility (IDTF) setting for patients with commercial insurance. LCS LDCT coverage is not allowed in IDTF setting for patients with Medicare.                                   |
| <b>Billing</b>                               |   |
| Pre-authorization requirement                | Typically required for Medicare Advantage and commercial insurance.<br><br>Not required for traditional Medicare.<br><br>May be required for Medicaid (varies by state).  |
| Shared decision-making visit                 | Code: G0296; no co-pay (ICD 10 code — recommend Z87.891 or F17.210).  |
| LCS LDCT billing code                        | Code: G0297; no co-pay (ICD-10 code — recommend Z87.891 or F17.210).  |
| Tobacco cessation counseling                 | 3–10 minutes: CPT Code 99406 (ICD 10 code — recommend F17.210).<br>Over 10 minutes: CPT Code 99407 (ICD 10 code — recommend F17.210).   |
| Interval CT (diagnostic CT)                  | Diagnostic CT Code: 71250; co-pay typically required.<br>Diagnostic CTA Code: 71275; co-pay typically required.   |
| <b>Follow up</b>                             |   |
| Lung-RADS 3                                  | 6-month follow-up CT; if unchanged or smaller ➡ LCS LDCT 1 year after the follow-up CT.   |
| Lung-RADS 4A                                 | 3-month follow-up CT; if unchanged or smaller ➡ 6–9 month follow-up LDCT; Then, if unchanged or smaller ➡ LCS LDCT 1 year after last follow-up CT.  |
| Lung cancer diagnosis confirmed              | Discharge from LCS; patient may return to screening after appropriate post-treatment surveillance.  |

#### Commonly Used Acronyms:

|        |  |
|--------|--|
| ACR:   | American College of Radiology  |
| CMS:   | Centers for Medicare & Medicaid Services   |
| E&M:   | Evaluation and management  |
| IDTF:  | Independent Diagnostic Testing Facility  |
| LCS:   | Lung cancer screening  |
| LDCT:  | Low-dose chest CT  |
| NCCN:  | National Comprehensive Cancer Network (NCCN group 2 criteria = ages ≥50, at least 20 pack-years and another risk factor) |
| NLCRT: | National Lung Cancer Roundtable  |