

The American College of Radiology, with more than 30,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

2022 (Resolution 20)*

ACR–ASNR–SPR PRACTICE PARAMETER FOR THE PERFORMANCE OF COMPUTED TOMOGRAPHY (CT) IN THE EVALUATION AND CLASSIFICATION OF TRAUMATIC BRAIN INJURY

PREAMBLE

This document is an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care¹. For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner considering all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by variables such as the condition of the patient, limitations of available resources, or advances in knowledge or technology after publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document may consider documenting in the patient record information sufficient to explain the approach taken.

The practice of medicine involves the science, and the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The purpose of this document is to assist practitioners in achieving this objective.

¹ *Iowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing* 831 N.W.2d 826 (Iowa 2013) Iowa Supreme Court refuses to find that the *ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures* (Revised 2008) sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, *Stanley v. McCarver*, 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do not establish the standard of care.

I. INTRODUCTION

This practice parameter was developed collaboratively by the American College of Radiology (ACR), the American Society of Neuroradiology (ASNR), and the Society for Pediatric Radiology (SPR).

Traumatic brain injury (TBI) is one of the most common neurologic disorders, currently affecting 1.7 million Americans each year [1,2]. The incidence of TBI, especially mild TBI, is underestimated [3], as patients frequently dismiss their symptoms and never present to the emergency department, or they believe that the admission of symptoms may compromise their work situation (eg, athletes, military [4]). Although the majority of patients (nearly 80%) with diagnosed TBI are treated and released from the emergency department [5], the remaining 20% have more significant injuries resulting in approximately 275,000 hospitalizations and 52,000 deaths each year. Furthermore, TBI contributes to a third of all injury-related deaths in the United States. The economic cost of TBI was estimated at \$76.5 billion in 2010 (\$11.5 billion in direct medical costs and \$64.8 billion in indirect costs such as lost wages, lost productivity, and nonmedical expenditures) [6]. Moreover, affected military veterans generate an annual cost of \$11,700 of medical treatment per patient compared with \$2,400 in TBI-free veterans [7]. Leading causes of TBI in the general population include falls, motor vehicle accidents, assaults, and sports-related injuries.

Imaging plays an essential role in identifying TBI patients with intracranial injury. The goals of imaging include (1) detecting injuries that may require immediate surgical or procedural intervention; (2) detecting injuries that may benefit from early medical therapy and/or vigilant neurologic supervision; and (3) determining the prognosis of patients to tailor rehabilitative therapy or help with family counseling and discharge planning. A wide variety of imaging techniques have become available to assess patients presenting with TBI. This, coupled with the inconsistent use of clinical decision rules [8], has led to increased utilization and numerous variation in imaging practices. Among hospitals reporting to the National Hospital Ambulatory Medical Care Survey, CT utilization for head trauma in the pediatric population increased from 12.8% in 1995 to 28.6% in 2000 despite stable hospitalization rates from head trauma [9]. The practical challenge for physicians is to understand which imaging techniques to implement and how to use them optimally for specific patients. There are early reports at individual institutions that adoption of machine learning algorithms into the clinical workflow might assist physicians in triage and detection of intracranial hemorrhage in patients with TBI [10,11].

This practice parameter focuses on computed tomography (CT) and should assist referring physicians faced with the task of appropriately ordering CT scans in the particular TBI patient for whom they are providing care. It should also help radiologists advise their clinical colleagues on appropriate CT imaging utilization for TBI patients. For practical purposes, recommendations are presented separately for TBI severity and for acute, subacute, and chronic TBI, as defined by the Defense and Veterans Brain Injury Center (DVBIC) recommendations. Acute injuries refer to those from time of injury to 7 days; subacute injuries refer to those between 8 days to 89 days post-injury; and chronic injuries refer to those injuries 90 days or greater post-injury. The severity of TBI is usually classified by the Glasgow Coma Scale (GCS) score as mild, moderate, or severe. Mild TBI is generally defined as demonstrating GCS scores of 13 to 15, moderate TBI demonstrates GCS scores of 9 to 12, and severe TBI demonstrates GCS scores of 3 to 8 [12-14]. These recommendations are concordant with the ACR Appropriateness Criteria for head trauma [15].

II. INDICATIONS

Indications for CT TBI include, but are not limited to, the following:

1. Acute moderate or severe TBI [12,16-30]
2. Acute mild TBI [20,31-49]
3. Follow-up imaging in TBI [50-53]
4. Subacute to chronic TBI [54-56]
5. Pediatric TBI [15,24,57-83]
6. Neurovascular trauma [84-96]

For information on contrast, see the [ACR–SPR Practice Parameter for the Use of Intravascular Contrast Media](#) [97] and the [ACR Manual on Contrast Media](#) [98].

For the pregnant or potentially pregnant patient, see the [ACR–SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Patients with Ionizing Radiation](#) [99].

III. QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL

See the [ACR Practice Parameter for Performing and Interpreting Diagnostic Computed Tomography \(CT\)](#) [100] and the [ACR–ASNR–SPR Practice Parameter for the Performance and Interpretation of Cervicocerebral Computed Tomography Angiography \(CTA\)](#) [101].

A. Physician

Examinations must be performed under the supervision of and interpreted by a physician who has the following qualifications:

The physician should meet the criteria listed in the [ACR Practice Parameter for Performing and Interpreting Diagnostic Computed Tomography \(CT\)](#) [100] and in the [ACR–SPR Practice Parameter for the Use of Intravascular Contrast Media](#) [97] and should be trained in radiation safety.

1. The supervising physician must have adequate understanding of the indications, risks, and benefits of the examination, as well as alternative imaging procedures. The supervising physician is responsible for specifying the parameters of image acquisition; the route, volume, timing, type, and rate of contrast injection; and the method of image reconstruction and archival. The physician should monitor the quality of the images, be aware of potential artifacts [102], and interpret the study. Interpreting physicians must have knowledge of the benefits and risks of the procedures. Knowledge of the head and neck anatomy, including the vascular anatomy, and diseases of the intracranial and extracranial cerebrovascular system and their treatment is required.
2. Physicians meeting the aforementioned criteria additionally must have knowledge of the anatomy and pathophysiology relevant to the examination and of the spectrum of nonvascular abnormalities presenting on CT scans. They should be capable of identifying and characterizing important nonvascular abnormalities that may be visualized on CT scan and CT angiography (CTA), such as neoplasia, sequelae of infection, noninfectious inflammatory diseases, congenital anomalies, and normal anatomic variants, and any other abnormalities that may affect patient care and might necessitate treatment or further characterization through additional diagnostic testing.
3. The physician should be familiar with the use of 3-D processing workstations and be capable of performing or directing creation of 3-D renderings, multiplanar reformations, and measurements of vessel dimensions.
4. The physician should work with a Qualified Medical Physicist to optimize site-specific CT scan and CTA scan protocols, when possible.

B. Technologist

The technologist should have the responsibility of patient comfort, preparing and positioning the patients for the CT examination, monitoring the patient during the examination, and obtaining the CT data in a manner prescribed by the supervising physician. For the intravenous (IV) administration of contrast material for CTA, qualifications for technologists performing IV injections should be in compliance with current ACR policy and existing operating procedures or manuals at the imaging facility. The technologist should perform the regular quality control testing of the CT system under the supervision of a medical physicist ([ACR–SPR Practice Parameter for the Use of Intravascular Contrast Media](#) [97]).

The technologist performing CT examinations should be certified by the American Registry of Radiologic Technologists or have an unrestricted state license with documented training and experience in CT.

IV. SPECIFICATIONS OF THE EXAMINATION

The written or electronic request for CT of the head should provide sufficient information to demonstrate the medical necessity of the examination and allow for the proper performance and interpretation of the examination.

Documentation that satisfies medical necessity includes 1) signs and symptoms and/or 2) relevant history (including known diagnoses). The provision of additional information regarding the specific reason for the examination or a provisional diagnosis would be helpful and may at times be needed to allow for the proper performance and interpretation of the examination.

The request for the examination must be originated by a physician or other appropriately licensed health care provider. The accompanying clinical information should be provided by a physician or other appropriately licensed health care provider familiar with the patient's clinical problem or question and consistent with the state scope of practice requirements. (ACR Resolution 35 adopted in 2006 – revised in 2016, Resolution 12-b)

A. General Considerations

CT protocols for brain imaging should be designed to answer the specific clinical question. The supervising physician should be familiar with the indications for each examination, relevant patient history, and potential adverse reactions to contrast media. The supervising physician should be familiar with how individual CT settings affect radiation dose and image quality, including field of view (FOV), collimation, pitch, automated exposure control, and image reconstruction algorithms such as iterative reconstruction [103]. The goal of CT scanning is to obtain diagnostic information from images of sufficient quality. Protocols should be optimized to deliver the lowest dose required to achieve appropriate image quality, especially in pediatric patients, and should be reviewed and updated as needed in light of new clinically applicable developments [104-112].

B. Head CT

Performance Standards

To achieve acceptable clinical CT scans of the brain, the CT scanner should meet or exceed the following specifications:

1. Scan times: per slice or image not more than 2 seconds.
2. Slice thickness: acquired slice thickness should be 2 mm or less, whereas reconstructed slice thickness should be 5 mm or less. Axial reconstructions might be performed to correct acquisition plane if necessary.
3. Limiting spatial resolution: must be measured to verify that it meets the unit manufacturer's specifications. Limiting spatial resolution should be >10 lp/cm for a display FOV <24 cm.
4. Table pitch: no greater than 2 for most CT scanners, but pitch may be increased for dual-source scanners for sole evaluation of bone anatomy (craniofacial)

For further information, see the American Association of Physicists in Medicine (AAPM) [Routine Adult Head \(Brain\) Protocols](#) [113].

CT brain imaging is performed for the evaluation of a variety of pathologies that require appropriate techniques for acquisition and viewing. CT brain imaging may be performed with a sequential single-slice technique, multislice helical (spiral) protocol, or multidetector multislice algorithm [114,115]. Use of these techniques is dependent on clinical indication, scanner capability, and image quality requirements. For CT of the brain, contiguous or overlapping axial slices should be acquired with a slice thickness of no greater than 5 mm. In addition to directly acquired axial images, reformatted images in coronal, sagittal, true axial, or other more complex planes may be

constructed from the axial data set to answer specific clinical questions. Additionally, axial reconstructed images should be presented with at least 2 different kernels, using both a brain/soft-tissue and bone kernel. Brain images should be reviewed at dedicated workstations and with window settings appropriate for demonstrating brain, bone, and soft-tissue abnormalities as well as hemorrhage.

C. Cervicocerebral CTA

1. Patient Selection and Preparation

Patients without absolute contraindication to the administration of iodinated contrast media are candidates for cervicocerebral CTA. In cases of relative contraindication to the administration of iodinated contrast medium, measures to reduce the possibility of contrast medium reactions or nephrotoxicity should be followed to the extent that the patient's condition allows, as defined in the [ACR–SPR Practice Parameter for the Use of Intravascular Contrast Media](#) [97,116].

When possible, patients should be well hydrated, and IV access should be established. A 20-gauge or larger antecubital IV catheter should be placed ideally on the right side to accommodate an optimal rate of 4 or 5 mL/s of iodinated contrast media. Smaller catheters that can withstand the prescribed injection rates can be used, and lower injection rates may be used for pediatric patients. All catheters used for the CTA examination should first be tested with a rapidly injected bolus of sterile saline to ensure that the venous access is secure and can accommodate the rapid bolus, minimizing the risk of contrast medium extravasations. The injection site should be monitored by medical personnel trained in the rapid recognition of IV extravasations. Department procedures for care of IV extravasations should be documented and applied if necessary.

2. Examination Technique

The CTA acquisition should be performed with a section thickness of 1.5 mm or less, depending on the vascular territory to be assessed. The scan should be reconstructed with overlapping sections. In the setting of trauma, CTA imaging of the neck should be obtained and the acquisition should at least cover the aortic arch, the origin and cervical course of the subclavian and carotid arteries, and proximal subclavian arteries, through the Circle of Willis. Automated tube voltage selection can also be employed in conjunction with tube current modulation when available. Finally, the display FOV must be sufficient to allow an assessment of the vasculature of interest, the end-organ, and adjacent tissues.

Because of substantial variations in the time required for an IV injection of nonionic contrast medium (iodine, 300-370 mg/mL) to reach the target vascular anatomy, an assessment of patient-specific circulation time is frequently required, especially for arterial imaging, although not mandatory. Circulation timing can be performed using one of the following techniques [117].

- a. IV injection of a small test bolus (eg, 10-15 mL) of contrast medium at the same rate and through the same access that will be used for the CTA followed by acquisition of sequential cine CT images at the level of the artery or vein of interest. The rate and intensity of enhancement of the lumen of interest are then used to create a time density curve. The peak of the curve is used to calculate the scanning delay postinjection. A perfusion CT series performed before the CTA can be used similarly to a test bolus for determining the timing of the CTA acquisition.
- b. The use of automated or semiautomated triggering software based on monitoring of the attenuation within the vessel of interest (or a great vessel such as the aorta) by the CT scanner following initiation of the full dose of contrast media injection. The CTA is automatically started when the enhancement in the vessel reaches a predetermined operator-selected level.

Ideally the administration of iodinated contrast media for the CTA should be performed with a minimum flow rate of 4 mL/s in any patient weighing 50 kg or more. Higher flow rates up to 6 mL/s are frequently required for larger patients, and in general, higher flow rates are required for shorter acquisitions. In

children, contrast medium dosing should be scaled to body weight. Injection rate should be scaled similarly and preferably delivered via powered injection. For young children and infants, a 22- or 24-gauge IV catheter may be the only option, and a 2 mL/s injection rate may be reasonable for these patients. For patients under 50 kg, a dose of 2 mL/kg should be considered. In summary, contrast injection parameters should be modified on an individual patient basis, and the volume of contrast medium should be selected with consideration of the patient's weight and comorbidities that might increase the risk of nephrotoxicity. When performing a cervicocerebral CTA, a right-arm injection is preferable to a left-arm injection to avoid artifacts from undiluted contrast medium in the left brachiocephalic vein. When possible, a bolus of saline should follow the iodinated contrast medium injection as this may reduce the volume of contrast medium required to achieve adequate vascular opacification.

V. DOCUMENTATION

Reporting should be in accordance with the [ACR Practice Parameter for Communication of Diagnostic Imaging Findings](#) [118].

Since 2009, multiple healthcare agencies involving experts from the international TBI community have worked on developing and refining Common Data Elements (CDEs) in TBI to promote the use of consistent terminology and definitions in characterizing intracranial injuries across all imaging studies, as well as all clinical aspects of TBI [119,120]. These CDEs can be used in a consistent manner for clinical practice, research, and treatment trials across multiple institutions and research studies. The CDEs include a list of the injuries that can be identified, with definitions of terms used to describe these injuries on the images, and recommended protocols and descriptors for image acquisition methods. The goal of the CDEs is to promote consistency across the field in future investigations aimed at evaluating TBI imaging. Clinical implementation of CDEs is time-consuming and challenging. Different groups have been working on streamlining the clinical implementation and adoption of CDEs for the reporting of TBI imaging studies. One such example is the NIRIS system [17,18,29,30] mentioned above. The supervising and interpreting physician should be aware of efforts to standardize reporting and consider adopting these when appropriate, to enhance communication with clinical care team members.

Cervicocerebral CTAs are preferentially interpreted on equipment that allows stacked dynamic paging of the primary axial and the reformatted CTA sections. A complete interpretation includes review of all images, including the scout and the axial CT sections (source images) and, as indicated, multiplanar/curved reformations, volume renderings, maximum-intensity projections, and other reconstructions produced during postprocessing. On occasion, the interpreting physician will personally create postprocessed images documenting important findings that are essential to the interpretation of the study [121]. These images should be archived with the patient's original study or other postprocessed images. Interpretation of the cervicocerebral CTA includes an assessment of the patency and caliber of the carotid and vertebral arteries, their origins, the carotid bifurcations, the intracranial arteries, possible occlusion, dissection, stenosis, and aneurysmal dilatation.

The visible regional anatomy and pathology should be commented on when appropriate. In the setting of suspected traumatic injury, the soft issues surrounding the vasculature and adjacent bony structures in the cervical region should be assessed. Comparison with prior studies should be performed when appropriate.

VI. EQUIPMENT SPECIFICATIONS

For specific issues regarding CT quality control, see the [ACR Practice Parameter for Performing and Interpreting Diagnostic Computed Tomography \(CT\)](#) [100].

Equipment monitoring should be in accordance with the [ACR–AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Computed Tomography \(CT\) Equipment](#) [122].

- A. For diagnostic quality CTA, the CT scanner should meet or exceed the following specifications:
 1. Cervicocerebral CTA should be performed on a multidetector CT (MDCT) scanner, preferably with greater than or equal to 4 active detector rows.

2. Gantry rotation: 1 second or less for cervicocerebral CTA.
3. Tube heat capacity that allows for a single ≥ 10 -second acquisition.
4. Section thickness: no greater than 1.5 mm.
5. A contrast medium power injector that allows programming of both the volume and flow rate must be used for head and neck CTA examinations.

To maximize information available from the CT scan and thus derive the full diagnostic benefit for the patient following X-ray irradiation, any CT scanner used for CTA must allow display and interpretation of the full 12 bits (from $-1,000$ to $3,095$ Hounsfield units) of attenuation information. Dual-energy CTA can be obtained when available to decrease total patient radiation dose, lower contrast administration, distinguish contrast from hemorrhage and calcium, and reduce hardware artifacts [123-126].

- B. Patient monitoring equipment and facilities for cardiopulmonary resuscitation, including vital signs monitoring equipment and support equipment, should be immediately available.

Appropriate emergency equipment and medications must be immediately available to treat adverse reactions associated with administered contrast. The equipment and medications should be monitored for inventory and drug expiration dates on a regular basis. The equipment, medications, and other emergency support must also be appropriate for the range of ages or sizes in the patient populations.

Radiologists, technologists, and staff members should be able to assist with procedures, patient monitoring, and patient support. A written policy should be in place for dealing with emergencies, such as cardiopulmonary arrest.

VII. RADIATION SAFETY IN IMAGING

Radiologists, medical physicists, non-physician radiology providers, radiologic technologists, and all supervising physicians have a responsibility for safety in the workplace by keeping radiation exposure to staff, and to society as a whole, "as low as reasonably achievable" (ALARA) and to assure that radiation doses to individual patients are appropriate, taking into account the possible risk from radiation exposure and the diagnostic image quality necessary to achieve the clinical objective. All personnel who work with ionizing radiation must understand the key principles of occupational and public radiation protection (justification, optimization of protection, application of dose constraints and limits) and the principles of proper management of radiation dose to patients (justification, optimization including the use of dose reference levels). https://www-pub.iaea.org/MTCDD/Publications/PDF/PUB1775_web.pdf

Nationally developed guidelines, such as the [ACR's Appropriateness Criteria](#)®, should be used to help choose the most appropriate imaging procedures to prevent unnecessary radiation exposure.

Facilities should have and adhere to policies and procedures that require ionizing radiation examination protocols (radiography, fluoroscopy, interventional radiology, CT) to vary according to diagnostic requirements and patient body habitus to optimize the relationship between appropriate radiation dose and adequate image quality. Automated dose reduction technologies available on imaging equipment should be used, except when inappropriate for a specific exam. If such technology is not available, appropriate manual techniques should be used.

Additional information regarding patient radiation safety in imaging is available from the following websites – Image Gently® for children (www.imagegently.org) and Image Wisely® for adults (www.imagewisely.org). These advocacy and awareness campaigns provide free educational materials for all stakeholders involved in imaging (patients, technologists, referring providers, medical physicists, and radiologists).

Radiation exposures or other dose indices should be periodically measured by a Qualified Medical Physicist in accordance with the applicable ACR Technical Standards. Monitoring or regular review of dose indices from patient imaging should be performed by comparing the facility's dose information with national benchmarks, such

as the ACR Dose Index Registry and relevant publications relying on its data, applicable ACR Practice Parameters, NCRP Report No. 172, Reference Levels and Achievable Doses in Medical and Dental Imaging: Recommendations for the United States or the Conference of Radiation Control Program Director’s National Evaluation of X-ray Trends; 2006, 2009, amended 2013, revised 2023 (Res. 2d).

When possible, CT imaging of the head should consider the following to minimize radiation dose and maintain image quality:

1. Center the patient in the gantry [127].
2. Angling of the gantry to exclude the orbits during brain imaging [128,129].
3. Remove unnecessary objects from the patient.

Dose-minimization CT techniques should be used, especially in the pediatric population [130].

Diagnostic Reference Levels (DRL) and Achievable Doses (AD) are national benchmarks for radiation protection and optimization that provide a comparison for facilities in order to review techniques and determine whether acceptable image quality can be achieved at lower doses. Published levels are available [131]. For further information, see the [ACR–AAPM–SPR Practice Parameter for Diagnostic Reference Levels and Achievable Doses in Medical X-Ray Imaging](#) [132]. Attention to dose is particularly important but also particularly challenging in the pediatric population, when age and size specific protocols should be considered [133].

VIII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *Position Statement on Quality Control & Improvement, Safety, Infection Control, and Patient Education* on the ACR website (<https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Quality-Control-and-Improvement>).

In addition to CT radiation safety and quality control, appropriateness studies, and utilization review, a facilitation of best practices for CT brain imaging should be considered and encouraged as part of a comprehensive continuous quality improvement program [23,134-141]. Moreover, best practices that are evidence based and instituted as part of care team and/or multidisciplinary policies to assure equitable access and utilization and the reduction of health disparities should also be considered and encouraged [142-144].

ACKNOWLEDGEMENTS

This practice parameter was revised according to the process described under the heading *The Process for Developing ACR Practice Parameters and Technical Standards* on the ACR website (<https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards>) by the Committee on Practice Parameters – Neuroradiology of the ACR Commission on Neuroradiology and the Committee on Practice Parameters – Pediatric Radiology of the ACR Commission on Pediatric Radiology in collaboration with the ASNR and the SPR.

Writing Committee – members represent their societies in the initial and final revision of this practice parameter

ACR

Max Wintermark, MD, Chair
Einat Blumfield, MD
John E. Jordan, MD, MPP, FACR
Sumit N. Niogi, MD, PhD

ASNR

Jason W. Allen, MD, PhD
Kavita K. Erickson, MD
Nandini D. Patel, MD
Eric J. Russell, MD, FACR

SPR

Aaron M. Betts, MD

Mai-Lan Ho, MD
Gaurav Saigal, MD
Nicholas V. Stence, MD

Committee on Practice Parameters – Neuroradiology

(ACR Committee responsible for sponsoring the draft through the process)

Steven W. Hetts, MD, Chair	Gerald Drocton, MD
Lubdha M. Shah, MD, Vice Chair	Kavita K. Erickson, MD
Ashley H. Aiken, MD	Adam E. Flanders, MD
Sameer A. Ansari, MD, PhD	Masis Isikbay, MD, BS
Kristine A. Blackham, MD	Raymond K. Tu, MD, FACR
Gloria C Chiang, MD	Max Wintermark, MD

Committee on Practice Parameters – Pediatric Radiology

(ACR Committee responsible for sponsoring the draft through the process)

Terry L. Levin, MD, FACR, Chair	Jane Sun Kim, MD
John B. Amodio, MD, FACR	Jennifer A Knight, MD
Jesse Berman, MD	Jessica Kurian, MD
Tara M. Catanzano, MB, BCh	Helen R. Nadel, MD
Harris L. Cohen, MD, FACR	Erica Poletto, MD
Kassa Darge, MD, PhD	Richard B. Towbin, MD, FACR
Dorothy L. Gilbertson-Dahdal, MD	Andrew T. Trout, MD
Lauren P. Golding, MD	Esben S. Vogelius, MD
Adam Goldman-Yassen, MD	Jason Wright, MD
Safwan S. Halabi, MD	

John E. Jordan, MD, MPP, FACR, Chair, Commission on Neuroradiology
Richard A. Barth, MD, FACR, Chair, Commission on Pediatric Radiology
David B. Larson, MD, MBA, Chair, Commission on Quality and Safety
Mary S. Newell, MD, FACR, Chair, Committee on Practice Parameters and Technical Standards

Comments Reconciliation Committee

Gaurang V. Shah, MD, FACR, Chair	David B. Larson, MD, MBA
Daniel G. Gridley, MD, Co-Chair	Paul A. Larson, MD, FACR
Jason W. Allen, MD	Terry L. Levin, MD, FACR
Joseph M. Aulino, MD	Mary S. Newell, MD, FACR
Richard A. Barth, MD, FACR	Sumit N. Niogi, MD, PhD
Aaron M. Betts, MD	Thomas J. O'Neill, MD
Einat Blumfield, MD	Nandini D. Patel, MD
Julie Bykowski, MD	Eric J. Russell, MD, FACR
Timothy A. Crummy, MD, FACR	Gaurav Saigal, MD
Kavita K. Erickson, MD	William F. Sensakovic, PhD
Steven W. Hetts, MD	Lubdha M. Shah, MD
Mai-Lan Ho, MD	Nicholas V. Stence, MD
John E. Jordan, MD, MPP, FACR	Max Wintermark, MD
Amy L. Kotsenas, MD, FACR	

REFERENCES

1. Faul M, Xu L, Wald M, Coronado V. Traumatic brain injury in the United States: emergency department visits, hospitalizations and deaths 2002-2006. Center for Disease Control and Prevention, National Center for Injury Prevention and Control 2010.

2. Marin JR, Weaver MD, Yealy DM, Mannix RC. Trends in visits for traumatic brain injury to emergency departments in the United States. *JAMA* 2014;311:1917-9.
3. Ilie G, Boak A, Adlaf EM, Asbridge M, Cusimano MD. Prevalence and correlates of traumatic brain injuries among adolescents. *JAMA* 2013;309:2550-2.
4. AD G. *Brain Injury: Applications from War and Terrorism*: Lippincott Williams & Wilkins; 2014.
5. Centers for Disease, Control and Prevention NCFIPaC. *Report to congress on mild traumatic brain injury in the United States: steps to prevent a serious public health problem*: National Center for Injury Prevention and Control; 2003.
6. Center for Disease Control. Injury and violence prevention and control: traumatic brain injury. Available at: <https://www.cdc.gov/traumaticbraininjury/index.html>. Accessed February 25, 2021.
7. Congressional Budget Office. The veterans health administration's treatment of PTSD and traumatic brain injury among recent combat veterans. Available at: <https://www.cbo.gov/publication/42969>. Accessed February 25, 2021.
8. Graham ID, Stiell IG, Laupacis A, O'Connor AM, Wells GA. Emergency physicians' attitudes toward and use of clinical decision rules for radiography. *Acad Emerg Med* 1998;5:134-40.
9. Blackwell CD, Gorelick M, Holmes JF, Bandyopadhyay S, Kuppermann N. Pediatric head trauma: changes in use of computed tomography in emergency departments in the United States over time. *Ann Emerg Med* 2007;49:320-4.
10. Arbabshirani MR, Fornwalt BK, Mongelluzzo GJ, et al. Advanced machine learning in action: identification of intracranial hemorrhage on computed tomography scans of the head with clinical workflow integration. *NPJ Digit Med* 2018;1:9.
11. O'Neill TJ, Xi Y, Stehel E, et al. Active Reprioritization of the Reading Worklist Using Artificial Intelligence Has a Beneficial Effect on the Turnaround Time for Interpretation of Head CT with Intracranial Hemorrhage. *Radiol Artif Intell* 2021;3:e200024.
12. Cushman JG, Agarwal N, Fabian TC, et al. Practice management guidelines for the management of mild traumatic brain injury: the EAST practice management guidelines work group. *J Trauma* 2001;51:1016-26.
13. Iverson GL, Lovell MR, Smith S, Franzen MD. Prevalence of abnormal CT-scans following mild head injury. *Brain Inj* 2000;14:1057-61.
14. Servadei F, Teasdale G, Merry G, Neurotraumatology Committee of the World Federation of Neurosurgical S. Defining acute mild head injury in adults: a proposal based on prognostic factors, diagnosis, and management. *J Neurotrauma* 2001;18:657-64.
15. Expert Panel on Neurological I, Shih RY, Burns J, et al. ACR Appropriateness Criteria(R) Head Trauma: 2021 Update. *J Am Coll Radiol* 2021;18:S13-S36.
16. Care NCCfA. *Head injury: triage, assessment, investigation and early management of head injury in infants, children and adults*. London: National Collaborating Centre for Acute Care; 2007.
17. Chen H, Li Y, Jiang B, et al. Demographics and clinical characteristics of acute traumatic brain injury patients in the different Neuroimaging Radiological Interpretation System (NIRIS) categories. *J Neuroradiol* 2019.
18. Creeden S, Ding VY, Parker JJ, et al. Interobserver Agreement for the Computed Tomography Severity Grading Scales for Acute Traumatic Brain Injury. *J Neurotrauma* 2020;37:1445-51.
19. Jacobs B, Beems T, van der Vliet TM, Diaz-Arrastia RR, Borm GF, Vos PE. Computed tomography and outcome in moderate and severe traumatic brain injury: hematoma volume and midline shift revisited. *J Neurotrauma* 2011;28:203-15.
20. Jagoda AS, Bazarian JJ, Bruns JJ, Jr., et al. Clinical policy: neuroimaging and decisionmaking in adult mild traumatic brain injury in the acute setting. *Ann Emerg Med* 2008;52:714-48.
21. Maas AI, Hukkelhoven CW, Marshall LF, Steyerberg EW. Prediction of outcome in traumatic brain injury with computed tomographic characteristics: a comparison between the computed tomographic classification and combinations of computed tomographic predictors. *Neurosurgery* 2005;57:1173-82; discussion 73-82.
22. Marshall LF, Marshall SB, Klauber MR, et al. The diagnosis of head injury requires a classification based on computed axial tomography. *J Neurotrauma* 1992;9 Suppl 1:S287-92.
23. Nelson DW, Nystrom H, MacCallum RM, et al. Extended analysis of early computed tomography scans of traumatic brain injured patients and relations to outcome. *J Neurotrauma* 2010;27:51-64.
24. Nigrovic LE, Lee LK, Hoyle J, et al. Prevalence of clinically important traumatic brain injuries in children with minor blunt head trauma and isolated severe injury mechanisms. *Arch Pediatr Adolesc Med* 2012;166:356-61.
25. Panczykowski DM, Puccio AM, Scruggs BJ, et al. Prospective independent validation of IMPACT modeling as a prognostic tool in severe traumatic brain injury. *J Neurotrauma* 2012;29:47-52.
26. Raj R, Siironen J, Skrifvars MB, Hernesniemi J, Kivisaari R. Predicting outcome in traumatic brain injury: development of a novel computerized tomography classification system (Helsinki computerized tomography score). *Neurosurgery* 2014;75:632-46; discussion 46-7.
27. Tavender EJ, Bosch M, Green S, et al. Quality and consistency of guidelines for the management of mild traumatic brain injury in the emergency department. *Acad Emerg Med* 2011;18:880-9.
28. Thelin EP, Nelson DW, Vehvilainen J, et al. Evaluation of novel computerized tomography scoring systems in human traumatic brain injury: An observational, multicenter study. *PLoS Med* 2017;14:e1002368.

29. Wintermark M, Li Y, Ding VY, et al. Neuroimaging Radiological Interpretation System for Acute Traumatic Brain Injury. *J Neurotrauma* 2018;35:2665-72.
30. Zhou B, Ding VY, Li Y, et al. Validation of the NeuroImaging Radiological Interpretation System for Acute Traumatic Brain Injury. *J Comput Assist Tomogr* 2019;43:690-96.
31. af Geijerstam JL, Britton M. Mild head injury - mortality and complication rate: meta-analysis of findings in a systematic literature review. *Acta Neurochir (Wien)* 2003;145:843-50; discussion 50.
32. af Geijerstam JL, Oredsson S, Britton M, Investigators OS. Medical outcome after immediate computed tomography or admission for observation in patients with mild head injury: randomised controlled trial. *BMJ* 2006;333:465.
33. Barbosa RR, Jawa R, Watters JM, et al. Evaluation and management of mild traumatic brain injury: an Eastern Association for the Surgery of Trauma practice management guideline. *J Trauma Acute Care Surg* 2012;73:S307-14.
34. Bazarian JJ, Biberthaler P, Welch RD, et al. Serum GFAP and UCH-L1 for prediction of absence of intracranial injuries on head CT (ALERT-TBI): a multicentre observational study. *Lancet Neurol* 2018;17:782-89.
35. Borg J, Holm L, Cassidy JD, et al. Diagnostic procedures in mild traumatic brain injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med* 2004:61-75.
36. de Andrade AF, de Almeida AN, Bor-Seng-Shu E, Lourenco L, Mandel M, Marino R, Jr. The value of cranial computed tomography in high-risk, mildly head-injured patients. *Surg Neurol* 2006;65 Suppl 1:S1:10-1:13.
37. Fabbri A, Servadei F, Marchesini G, et al. Prospective validation of a proposal for diagnosis and management of patients attending the emergency department for mild head injury. *J Neurol Neurosurg Psychiatry* 2004;75:410-6.
38. Gill J, Latour L, Diaz-Arrastia R, et al. Glial fibrillary acidic protein elevations relate to neuroimaging abnormalities after mild TBI. *Neurology* 2018;91:e1385-e89.
39. Haydel MJ, Preston CA, Mills TJ, Luber S, Blaudeau E, DeBlieux PM. Indications for computed tomography in patients with minor head injury. *N Engl J Med* 2000;343:100-5.
40. Hsiang JN, Yeung T, Yu AL, Poon WS. High-risk mild head injury. *J Neurosurg* 1997;87:234-8.
41. Levin HS, Hanten G, Roberson G, et al. Prediction of cognitive sequelae based on abnormal computed tomography findings in children following mild traumatic brain injury. *J Neurosurg Pediatr* 2008;1:461-70.
42. Livingston DH, Lavery RF, Passannante MR, et al. Emergency department discharge of patients with a negative cranial computed tomography scan after minimal head injury. *Ann Surg* 2000;232:126-32.
43. Manolakaki D, Velmahos GC, Spaniolas K, de Moya M, Alam HB. Early magnetic resonance imaging is unnecessary in patients with traumatic brain injury. *J Trauma* 2009;66:1008-12; discussion 12-4.
44. Miller EC, Holmes JF, Derlet RW. Utilizing clinical factors to reduce head CT scan ordering for minor head trauma patients. *J Emerg Med* 1997;15:453-7.
45. Mower WR, Hoffman JR, Herbert M, et al. Developing a clinical decision instrument to rule out intracranial injuries in patients with minor head trauma: methodology of the NEXUS II investigation. *Ann Emerg Med* 2002;40:505-14.
46. Ro YS, Shin SD, Holmes JF, et al. Comparison of clinical performance of cranial computed tomography rules in patients with minor head injury: a multicenter prospective study. *Acad Emerg Med* 2011;18:597-604.
47. Sherer M, Stouter J, Hart T, et al. Computed tomography findings and early cognitive outcome after traumatic brain injury. *Brain Inj* 2006;20:997-1005.
48. Stiell IG, Wells GA, Vandemheen K, et al. The Canadian CT Head Rule for patients with minor head injury. *Lancet* 2001;357:1391-6.
49. Welch RD, Ayaz SI, Lewis LM, et al. Ability of Serum Glial Fibrillary Acidic Protein, Ubiquitin C-Terminal Hydrolase-L1, and S100B To Differentiate Normal and Abnormal Head Computed Tomography Findings in Patients with Suspected Mild or Moderate Traumatic Brain Injury. *J Neurotrauma* 2016;33:203-14.
50. Brown CV, Zada G, Salim A, et al. Indications for routine repeat head computed tomography (CT) stratified by severity of traumatic brain injury. *J Trauma* 2007;62:1339-44; discussion 44-5.
51. Cohen DB, Rinker C, Wilberger JE. Traumatic brain injury in anticoagulated patients. *J Trauma* 2006;60:553-7.
52. Reljic T, Mahony H, Djulbegovic B, et al. Value of repeat head computed tomography after traumatic brain injury: systematic review and meta-analysis. *J Neurotrauma* 2014;31:78-98.
53. Washington CW, Grubb RL, Jr. Are routine repeat imaging and intensive care unit admission necessary in mild traumatic brain injury? *J Neurosurg* 2012;116:549-57.
54. Mataro M, Poca MA, Sahuquillo J, et al. Neuropsychological outcome in relation to the traumatic coma data bank classification of computed tomography imaging. *J Neurotrauma* 2001;18:869-79.
55. McKee AC, Stern RA, Nowinski CJ, et al. The spectrum of disease in chronic traumatic encephalopathy. *Brain* 2013;136:43-64.
56. Stern RA, Daneshvar DH, Baugh CM, et al. Clinical presentation of chronic traumatic encephalopathy. *Neurology* 2013;81:1122-9.
57. Bigler ED. Neuroimaging biomarkers in mild traumatic brain injury (mTBI). *Neuropsychol Rev* 2013;23:169-209.
58. Bulas DI, Goske MJ, Applegate KE, Wood BP. Image Gently: why we should talk to parents about CT in children. *AJR Am J Roentgenol* 2009;192:1176-8.
59. Gonzalez AJ. Biological effects of low doses of ionizing radiation: a fuller picture. In: IAEA, ed. *IAEA Bulletin*; 1994.

60. Lindberg DM, Stence NV, Grubenhoff JA, et al. Feasibility and Accuracy of Fast MRI Versus CT for Traumatic Brain Injury in Young Children. *Pediatrics* 2019;144.
61. Lumba-Brown A, Lee MO, Brown I, et al. Emergency department implementation of abbreviated magnetic resonance imaging for pediatric traumatic brain injury. *J Am Coll Emerg Physicians Open* 2020;1:994-99.
62. Mathews JD, Forsythe AV, Brady Z, et al. Cancer risk in 680,000 people exposed to computed tomography scans in childhood or adolescence: data linkage study of 11 million Australians. *BMJ* 2013;346:f2360.
63. Maxfield MW, Schuster KM, McGillicuddy EA, et al. Impact of adaptive statistical iterative reconstruction on radiation dose in evaluation of trauma patients. *J Trauma Acute Care Surg* 2012;73:1406-11.
64. Nagayama Y, Nakaura T, Tsuji A, et al. Radiation dose reduction using 100-kVp and a sinogram-affirmed iterative reconstruction algorithm in adolescent head CT: Impact on grey-white matter contrast and image noise. *Eur Radiol* 2017;27:2717-25.
65. Nigrovic LE, Kuppermann N. Children With Minor Blunt Head Trauma Presenting to the Emergency Department. *Pediatrics* 2019;144.
66. Park JE, Choi YH, Cheon JE, et al. Image quality and radiation dose of brain computed tomography in children: effects of decreasing tube voltage from 120 kVp to 80 kVp. *Pediatr Radiol* 2017;47:710-17.
67. Pearce MS, Salotti JA, Little MP, et al. Radiation exposure from CT scans in childhood and subsequent risk of leukaemia and brain tumours: a retrospective cohort study. *Lancet* 2012;380:499-505.
68. Rozovsky K, Ventureyra EC, Miller E. Fast-brain MRI in children is quick, without sedation, and radiation-free, but beware of limitations. *J Clin Neurosci* 2013;20:400-5.
69. Suzuki S, Furui S, Ishitake T, et al. Lens exposure during brain scans using multidetector row CT scanners: methods for estimation of lens dose. *AJNR Am J Neuroradiol* 2010;31:822-6.
70. Dunning J, Daly JP, Lomas JP, et al. Derivation of the children's head injury algorithm for the prediction of important clinical events decision rule for head injury in children. *Arch Dis Child* 2006;91:885-91.
71. Durham SR, Liu KC, Selden NR. Utility of serial computed tomography imaging in pediatric patients with head trauma. *J Neurosurg* 2006;105:365-9.
72. Expert Panel on Pediatric I, Wootton-Gorges SL, Soares BP, et al. ACR Appropriateness Criteria((R)) Suspected Physical Abuse-Child. *J Am Coll Radiol* 2017;14:S338-S49.
73. Gerdung C, Dowling S, Lang E. Review of the CATCH study: a clinical decision rule for the use of computed tomography in children with minor head injury. *CJEM* 2012;14:243-7.
74. Kuppermann N, Holmes JF, Dayan PS, et al. Identification of children at very low risk of clinically-important brain injuries after head trauma: a prospective cohort study. *Lancet* 2009;374:1160-70.
75. Nigrovic LE, Schunk JE, Foerster A, et al. The effect of observation on cranial computed tomography utilization for children after blunt head trauma. *Pediatrics* 2011;127:1067-73.
76. Orman G, Wagner MW, Seeburg D, et al. Pediatric skull fracture diagnosis: should 3D CT reconstructions be added as routine imaging? *J Neurosurg Pediatr* 2015;16:426-31.
77. Osmond MH, Klassen TP, Wells GA, et al. CATCH: a clinical decision rule for the use of computed tomography in children with minor head injury. *CMAJ* 2010;182:341-8.
78. Pickering A, Harnan S, Fitzgerald P, Pandor A, Goodacre S. Clinical decision rules for children with minor head injury: a systematic review. *Arch Dis Child* 2011;96:414-21.
79. Ryan ME, Jaju A, Ciolino JD, Alden T. Rapid MRI evaluation of acute intracranial hemorrhage in pediatric head trauma. *Neuroradiology* 2016;58:793-9.
80. Schonfeld D, Fitz BM, Nigrovic LE. Effect of the duration of emergency department observation on computed tomography use in children with minor blunt head trauma. *Ann Emerg Med* 2013;62:597-603.
81. Sheridan DC, Newgard CD, Selden NR, Jafri MA, Hansen ML. QuickBrain MRI for the detection of acute pediatric traumatic brain injury. *J Neurosurg Pediatr* 2017;19:259-64.
82. Cothren CC, Moore EE, Ray CE, Jr., Johnson JL, Moore JB, Burch JM. Cervical spine fracture patterns mandating screening to rule out blunt cerebrovascular injury. *Surgery* 2007;141:76-82.
83. Kopelman TR, Leeds S, Berardoni NE, et al. Incidence of blunt cerebrovascular injury in low-risk cervical spine fractures. *Am J Surg* 2011;202:684-8; discussion 88-9.
84. Berne JD, Cook A, Rowe SA, Norwood SH. A multivariate logistic regression analysis of risk factors for blunt cerebrovascular injury. *J Vasc Surg* 2010;51:57-64.
85. Biff WL, Moore EE, Offner PJ, Brega KE, Franciose RJ, Burch JM. Blunt carotid arterial injuries: implications of a new grading scale. *J Trauma* 1999;47:845-53.
86. Burlew CC, Biff WL, Moore EE, Barnett CC, Johnson JL, Bensard DD. Blunt cerebrovascular injuries: redefining screening criteria in the era of noninvasive diagnosis. *J Trauma Acute Care Surg* 2012;72:330-5; discussion 36-7, quiz 539.
87. Chokshi FH, Munera F, Rivas LA, Henry RP, Quencer RM. 64-MDCT angiography of blunt vascular injuries of the neck. *AJR Am J Roentgenol* 2011;196:W309-15.
88. Desai NK, Kang J, Chokshi FH. Screening CT angiography for pediatric blunt cerebrovascular injury with emphasis on the cervical "seatbelt sign". *AJNR Am J Neuroradiol* 2014;35:1836-40.

89. Dhillon RS, Barrios C, Lau C, et al. Seatbelt sign as an indication for four-vessel computed tomography angiogram of the neck to diagnose blunt carotid artery and other cervical vascular injuries. *Am Surg* 2013;79:1001-4.
90. Fleck SK, Langner S, Baldauf J, Kirsch M, Rosenstengel C, Schroeder HW. Blunt craniocervical artery injury in cervical spine lesions: the value of CT angiography. *Acta Neurochir (Wien)* 2010;152:1679-86.
91. Franz RW, Willette PA, Wood MJ, Wright ML, Hartman JF. A systematic review and meta-analysis of diagnostic screening criteria for blunt cerebrovascular injuries. *J Am Coll Surg* 2012;214:313-27.
92. Liang T, McLaughlin PD, Louis L, Nicolaou S. Review of multidetector computed tomography angiography as a screening modality in the assessment of blunt vascular neck injuries. *Can Assoc Radiol J* 2013;64:130-9.
93. Lohrer L, Vieth V, Nassenstein I, et al. Blunt cerebrovascular injuries in acute trauma care: a screening protocol. *Eur Spine J* 2012;21:837-43.
94. McKinney A, Ott F, Short J, McKinney Z, Truwit C. Angiographic frequency of blunt cerebrovascular injury in patients with carotid canal or vertebral foramen fractures on multidetector CT. *Eur J Radiol* 2007;62:385-93.
95. Patterson BO, Holt PJ, Cleanthis M, et al. Imaging vascular trauma. *Br J Surg* 2012;99:494-505.
96. Sliker CW. Blunt cerebrovascular injuries: imaging with multidetector CT angiography. *Radiographics* 2008;28:1689-708; discussion 709-10.
97. American College of Radiology. ACR–SPR practice parameter for the use of intravascular contrast media Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/IVCM.pdf>. Accessed February 26, 2021.
98. American College of Radiology. ACR manual on contrast media. Available at: https://www.acr.org/-/media/ACR/Files/Clinical-Resources/Contrast_Media.pdf. Accessed February 26, 2021.
99. American College of Radiology. ACR–SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Patients with Ionizing Radiation. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Pregnant-Pts.pdf>. Accessed January 27, 2021.
100. American College of Radiology. ACR practice parameter for performing and interpreting diagnostic computed tomography (CT). Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CT-Perf-Interpret.pdf>. Accessed January 27, 2021.
101. American College of Radiology. ACR–ASNR–SPR practice parameter for the Performance and interpretation of cervicocerebral computed tomography angiography (CTA) Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CervicoCerebralCTA.pdf>. Accessed February 26, 2021.
102. Kim JJ, Dillon WP, Glastonbury CM, Provenzale JM, Wintermark M. Sixty-four-section multidetector CT angiography of carotid arteries: a systematic analysis of image quality and artifacts. *AJNR Am J Neuroradiol* 2010;31:91-9.
103. O'Hora L, Foley SJ. Iterative reconstruction and automatic tube voltage selection reduce clinical CT radiation doses and image noise. *Radiography (Lond)* 2018;24:28-32.
104. Brady Z, Ramanauskas F, Cain TM, Johnston PN. Assessment of paediatric CT dose indicators for the purpose of optimisation. *Br J Radiol* 2012;85:1488-98.
105. Diekmann S, Siebert E, Juran R, et al. Dose exposure of patients undergoing comprehensive stroke imaging by multidetector-row CT: comparison of 320-detector row and 64-detector row CT scanners. *AJNR Am J Neuroradiol* 2010;31:1003-9.
106. Hoang JK, Wang C, Frush DP, et al. Estimation of radiation exposure for brain perfusion CT: standard protocol compared with deviations in protocol. *AJR Am J Roentgenol* 2013;201:W730-4.
107. Kilic K, Erbas G, Guryildirim M, et al. Quantitative and qualitative comparison of standard-dose and low-dose pediatric head computed tomography: a retrospective study assessing the effect of adaptive statistical iterative reconstruction. *J Comput Assist Tomogr* 2013;37:377-81.
108. Mahesh M, Scatarige JC, Cooper J, Fishman EK. Dose and pitch relationship for a particular multislice CT scanner. *AJR Am J Roentgenol* 2001;177:1273-5.
109. Perisinakis K, Seimenis I, Tzedakis A, Papadakis AE, Damilakis J. The effect of head shape, miscentering, and bowtie filter on peak patient tissue doses from modern brain perfusion 256-slice CT: how can we minimize the risk for deterministic effects? *Med Phys* 2013;40:011911.
110. Reimann AJ, Davison C, Bjarnason T, et al. Organ-based computed tomographic (CT) radiation dose reduction to the lenses: impact on image quality for CT of the head. *J Comput Assist Tomogr* 2012;36:334-8.
111. Vorona GA, Zuccoli G, Sutcliffe T, Clayton BL, Ceschin RC, Panigrahy A. The use of adaptive statistical iterative reconstruction in pediatric head CT: a feasibility study. *AJNR Am J Neuroradiol* 2013;34:205-11.
112. Yamauchi-Kawara C, Fujii K, Aoyama T, Yamauchi M, Koyama S. Radiation dose evaluation in multidetector-row CT imaging for acute stroke with an anthropomorphic phantom. *Br J Radiol* 2010;83:1029-41.
113. American Association of Physicists in Medicine. Adult routine head CT protocols. Available at: <https://www.aapm.org/pubs/ctprotocols/documents/adultroutineheadct.pdf>. Accessed March 2, 2021.
114. Ertl-Wagner B, Eftimov L, Blume J, et al. Cranial CT with 64-, 16-, 4- and single-slice CT systems-comparison of image quality and posterior fossa artifacts in routine brain imaging with standard protocols. *Eur Radiol* 2008;18:1720-6.

115. Jones TR, Kaplan RT, Lane B, Atlas SW, Rubin GD. Single- versus multi-detector row CT of the brain: quality assessment. *Radiology* 2001;219:750-5.
116. Oleinik A, Romero JM, Schwab K, et al. CT angiography for intracerebral hemorrhage does not increase risk of acute nephropathy. *Stroke* 2009;40:2393-7.
117. Lell M, Tomandl BF, Anders K, Baum U, Nkenke E. Computed tomography angiography versus digital subtraction angiography in vascular mapping for planning of microsurgical reconstruction of the mandible. *Eur Radiol* 2005;15:1514-20.
118. American College of Radiology. ACR practice parameter for communication of diagnostic imaging findings. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CommunicationDiag.pdf>. Accessed January 27, 2021.
119. Duhaime AC, Gean AD, Haacke EM, et al. Common data elements in radiologic imaging of traumatic brain injury. *Arch Phys Med Rehabil* 2010;91:1661-6.
120. Haacke EM, Duhaime AC, Gean AD, et al. Common data elements in radiologic imaging of traumatic brain injury. *J Magn Reson Imaging* 2010;32:516-43.
121. Leong JL, Batra PS, Citardi MJ. Three-dimensional computed tomography angiography of the internal carotid artery for preoperative evaluation of sinonasal lesions and intraoperative surgical navigation. *Laryngoscope* 2005;115:1618-23.
122. American College of Radiology. ACR–AAPM technical standard for diagnostic medical physics performance monitoring of computed tomography (CT) equipment. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CT-Equip.pdf>. Accessed January 27, 2021.
123. Ferda J, Novak M, Mirka H, et al. The assessment of intracranial bleeding with virtual unenhanced imaging by means of dual-energy CT angiography. *Eur Radiol* 2009;19:2518-22.
124. Jiang XY, Zhang SH, Xie QZ, et al. Evaluation of Virtual Noncontrast Images Obtained from Dual-Energy CTA for Diagnosing Subarachnoid Hemorrhage. *AJNR Am J Neuroradiol* 2015;36:855-60.
125. Kamalian S, Lev MH, Pomerantz SR. Dual-Energy Computed Tomography Angiography of the Head and Neck and Related Applications. *Neuroimaging Clin N Am* 2017;27:429-43.
126. Zhao L, Li F, Zhang Z, et al. Assessment of an advanced virtual monoenergetic reconstruction technique in cerebral and cervical angiography with third-generation dual-source CT: Feasibility of using low-concentration contrast medium. *Eur Radiol* 2018;28:4379-88.
127. Habibzadeh MA, Ay MR, Asl AR, Ghadiri H, Zaidi H. Impact of miscentering on patient dose and image noise in x-ray CT imaging: phantom and clinical studies. *Phys Med* 2012;28:191-9.
128. Nikupaavo U, Kaasalainen T, Reijonen V, Ahonen SM, Kortensniemi M. Lens dose in routine head CT: comparison of different optimization methods with anthropomorphic phantoms. *AJR Am J Roentgenol* 2015;204:117-23.
129. Wang J, Duan X, Christner JA, Leng S, Grant KL, McCollough CH. Bismuth shielding, organ-based tube current modulation, and global reduction of tube current for dose reduction to the eye at head CT. *Radiology* 2012;262:191-8.
130. Trattner S, Pearson GDN, Chin C, et al. Standardization and optimization of CT protocols to achieve low dose. *J Am Coll Radiol* 2014;11:271-78.
131. Kanal KM, Butler PF, Sengupta D, Bhargavan-Chatfield M, Coombs LP, Morin RL. U.S. Diagnostic Reference Levels and Achievable Doses for 10 Adult CT Examinations. *Radiology* 2017;284:120-33.
132. American College of Radiology. ACR–AAPM–SPR practice parameter for diagnostic reference levels and achievable doses in medical x-ray imaging. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Diag-Ref-Levels.pdf>. Accessed March 3, 2021.
133. Al Mahrooqi KMS, Ng CKC, Sun Z. Pediatric Computed Tomography Dose Optimization Strategies: A Literature Review. *J Med Imaging Radiat Sci* 2015;46:241-49.
134. Jhaveri KS, Saini S, Levine LA, et al. Effect of multislice CT technology on scanner productivity. *AJR Am J Roentgenol* 2001;177:769-72.
135. Jordan MJ, Lightfoote JB, Jordan JE. Quality outcomes of reinterpretation of brain CT imaging studies by subspecialty experts in neuroradiology. *J Natl Med Assoc* 2006;98:1326-8.
136. Jordan YJ, Jordan JE, Lightfoote JB, Ragland KD. Quality outcomes of reinterpretation of brain CT studies by subspecialty experts in stroke imaging. *AJR Am J Roentgenol* 2012;199:1365-70.
137. Korn A, Fenchel M, Bender B, et al. Iterative reconstruction in head CT: image quality of routine and low-dose protocols in comparison with standard filtered back-projection. *AJNR Am J Neuroradiol* 2012;33:218-24.
138. Rapalino O, Kamalian S, Kamalian S, et al. Cranial CT with adaptive statistical iterative reconstruction: improved image quality with concomitant radiation dose reduction. *AJNR Am J Neuroradiol* 2012;33:609-15.
139. Ren Q, Dewan SK, Li M, et al. Comparison of adaptive statistical iterative and filtered back projection reconstruction techniques in brain CT. *Eur J Radiol* 2012;81:2597-601.
140. Tan JS, Tan KL, Lee JC, Wan CM, Leong JL, Chan LL. Comparison of eye lens dose on neuroimaging protocols between 16- and 64-section multidetector CT: achieving the lowest possible dose. *AJNR Am J Neuroradiol* 2009;30:373-7.

141. You JJ, Gladstone J, Symons S, Rotstein D, Laupacis A, Bell CM. Patterns of care and outcomes after computed tomography scans for headache. *Am J Med* 2011;124:58-63 e1.
142. Marin JR, Rodean J, Hall M, et al. Racial and Ethnic Differences in Emergency Department Diagnostic Imaging at US Children's Hospitals, 2016-2019. *JAMA Netw Open* 2021;4:e2033710.
143. Ross AB, Kalia V, Chan BY, Li G. The influence of patient race on the use of diagnostic imaging in United States emergency departments: data from the National Hospital Ambulatory Medical Care survey. *BMC Health Serv Res* 2020;20:840.
144. Schragger JD, Patzer RE, Kim JJ, et al. Racial and Ethnic Differences in Diagnostic Imaging Utilization During Adult Emergency Department Visits in the United States, 2005 to 2014. *J Am Coll Radiol* 2019;16:1036-45.

*Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised, or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

Development Chronology for this Practice Parameter

2022 (Resolution 20)

Amended 2023 (Resolution 2c, 2d)