

The American College of Radiology, with more than 30,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

Revised 2017 (CSC/BOC)*

ACR–ASTRO PRACTICE PARAMETER FOR THE PERFORMANCE OF TOTAL BODY IRRADIATION

PREAMBLE

This document is an educational tool designed to assist practitioners in providing appropriate radiation oncology care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care¹. For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner in light of all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by the condition of the patient, limitations of available resources, or advances in knowledge or technology subsequent to publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document is advised to document in the patient record information sufficient to explain the approach taken.

The practice of medicine involves not only the science, but also the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The sole purpose of this document is to assist practitioners in achieving this objective.

¹ [Iowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing](#), ___ N.W.2d ___ (Iowa 2013) Iowa Supreme Court refuses to find that the *ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures* (Revised 2008) sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, [Stanley v. McCarver](#), 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do not establish the standard of care.

I. INTRODUCTION

This practice parameter was revised collaboratively by the American College of Radiology (ACR) and the American Society for Radiation Oncology (ASTRO).

Total body irradiation (TBI) is a radiotherapy technique that may be used as a component of preparative regimens for hematopoietic stem cell transplant (HSCT). TBI, usually in conjunction with chemotherapeutic agents, has proven useful for eradicating residual malignant or genetically disordered cells, for ablating hematopoietic stem cells, and for immunosuppression to reduce the risk of graft rejection.

According to data summarized by the Center for International Blood and Marrow Transplant Research, in 2013 the diseases most commonly treated with HSCT were (in decreasing order of frequency) multiple myeloma, non-Hodgkin lymphoma, acute myelogenous leukemia, myelodysplastic syndrome/myeloproliferative disease, Hodgkin disease, acute lymphoid leukemia, and additional malignant and nonmalignant diseases [1]. TBI has been used for many of these diseases but is not routine for all HSCT (eg, TBI is not commonly used for multiple myeloma transplants), and ongoing studies are evaluating the effectiveness of TBI-containing conditioning regimens as compared to chemotherapy alone for individual diseases [2-8]. HSCT is considered autologous if native stem cells are reinfused and allogeneic if the hematopoietic graft is derived from someone other than the recipient. Allogeneic grafts can be from related or unrelated individuals, but “matching” of donor and recipient is performed. The graft may be in the form of bone marrow, peripheral stem cells, and umbilical cord blood [9].

Unique features of TBI that make it a valuable component of transplant preparative regimens include:

1. No sparing of “sanctuary” sites such as testes and the central nervous system
2. Dose homogeneity to the whole body regardless of blood supply
3. Less chance of cross-resistance with other antineoplastic agents (chemotherapy)
4. No problems with excretion or detoxification
5. Ability to tailor the dose distribution by shielding specific organs or by “boosting” sites

A wide variety of TBI dose and fractionation schedules have been studied. The optimal regimen depends on a range of clinical variables, including patient age, disease, and type of HSCT. With competing goals of disease eradication and avoidance of toxicity, the most commonly accepted total dose of TBI for myeloablative HSCT is 12 to 15 Gy delivered in 6 to 12 fractions over 3 to 5 days [10-14]. Numerous investigators have shown that efficacy is improved and a variety of important late toxicities are significantly decreased when TBI is fractionated in 2 to 3 treatments per day [15,16]. Relatively low dose rates are also important for optimal outcome [17] and for reducing the risk of interstitial pneumonitis [18,19]. Many protocols require a dose rate of less than 0.2 Gy per minute, some as low as 0.1 Gy per minute.

Low-dose TBI, often in conjunction with chemotherapy, has recently emerged as an effective form of conditioning in reduced intensity HSCT for patients who may not be able to tolerate myeloablation because of poor performance status or comorbidity. Notable studies have included TBI doses of 2 to 6 Gy in 2 to 4 fractions [20-24]. Low-dose TBI is also being used as part of the conditioning regimen in salvage allogeneic hematopoietic cell transplantation in patients with graft rejection [25,26].

It is essential that the complicated treatment and care of the patient receiving TBI be well coordinated among the various subspecialties (medical oncology, radiation oncology, etc) and caregivers (physicians, nurses, physicists, psychologists, dieticians, transplant coordinators, etc). TBI presents a unique challenge because it results in potentially lethal myeloablation without intensive medical support and stem cell backup. Incorrectly delivered TBI may result in fatal toxicity. Anticipated immediate toxicity includes the following signs and symptoms: nausea, emesis, parotitis, xerostomia, headache, fatigue, mucositis, diarrhea, and loss of appetite [27]. Prophylactic interventions to manage these toxicities include intravenous hydration, antiemetics, and antimucositis agents [28]. Patients must be counseled regarding the risks of long-term sequelae of TBI, which vary in incidence depending on the clinical scenario, age at transplant, and TBI regimen, with unique side-effect

profile inherent to the age at the time of transplant. Some late risks may include pneumonopathy [29], veno-occlusive disease of the liver [30], kidney dysfunction [31], cataracts [17], hypothyroidism [32], infertility [33], secondary malignancies [14,34-36], growth and developmental delay, decreased bone density, insulin resistance, and neurocognitive effects [37,38]. Because of the significant risk associated with this treatment, the entire team must take great care to assure the best possible multidisciplinary care with attention to all facets of TBI.

Although the techniques of TBI vary widely from institution to institution, certain basic principles apply, such as the achievement of relative-dose homogeneity throughout the body, with the exception of intentionally shielded or boosted areas. Some centers use opposing anterior and posterior fields with the patient standing upright several meters from the source and the beam pointed horizontally. A beam spoiler may be used to prevent skin sparing [39]. An alternative approach irradiates patients with lateral fields in a sitting or partly reclining position [40]. This approach is usually better tolerated by patients but can present additional dosimetric challenges that must be considered and addressed to improve dose uniformity. Very young children who require anesthesia may be treated lying on the floor with the gantry pointing downward and with the spoiler and blocks placed above the patient. Successful planning and delivery of TBI require close interaction and coordination among the radiation oncologists, medical physicists, dosimetrists, nurses, and radiation therapists.

This practice parameter describes a quality assurance program for both adult and pediatric TBI and is supplementary to the [ACR–ASTRO Practice Parameter for Radiation Oncology](#) [41] and the [ACR–AAPM Technical Standard for the Performance of Radiation Oncology Physics for External Beam Therapy](#) [42].

II. PROCESS OF TOTAL BODY IRRADIATION

The use of TBI is a complex process involving many trained personnel who carry out highly coordinated activities.

A. Clinical Evaluation

The initial evaluation should include a detailed history, including a review of issues that may have an impact on treatment tolerance (previous radiotherapy to sensitive organs, including the spinal cord and whole brain [pediatric patients]; factors affecting pulmonary, renal, or hepatic function; presence of implanted battery-operated medical devices; cancer predisposition syndromes; and exposure to infectious agents); physical examination; review of all pertinent diagnostic and laboratory tests, including pulmonary function studies; and communication with the referring physician and other physicians involved in the patient’s care in accordance with the [ACR–ASTRO Practice Parameter for Communication: Radiation Oncology](#) [43]. Careful review of the applicable treatment plan or clinical trial protocol for the particular disease being treated is essential since standardized institutional or cooperative group protocols are typically used for transplantation.

As with delivery of any chemotherapy or radiotherapy, policies and procedures should be in place to determine whether a female patient is pregnant before initiating any component of a transplant program, including TBI. If the patient is determined to be pregnant, other therapies should be considered in an effort to preserve the pregnancy. Alternatively, if the patient wishes to proceed with transplant, the pregnancy may be electively terminated.

B. Informed Consent

Prior to simulation and treatment, informed consent must be obtained, documented, and in compliance with applicable laws, regulations, or policies, in accordance with the [ACR Practice Parameter on Informed Consent – Radiation Oncology](#) [44]. This should include a detailed discussion of the benefits and potential tissue-specific acute and late toxicities of TBI, as well as the details of, rationale for, and alternatives to TBI.

C. Treatment Planning

Treatment planning for TBI requires detailed knowledge of the specific transplant program to be followed (either on or off of a clinical trial). Specific treatment parameters to be determined in advance of treatment include field size, collimator rotation, treatment distance, dose per fraction, dose rate, total dose, number of fractions per day, interval between fractions, beam energy, geometry to achieve dose homogeneity, bolus or beam spoilers to increase skin dose, shielding and dose compensation requirements (eg, lungs, kidneys), and boost specifications (eg, testes, chest wall, brain, craniospinal axis, etc). Patient thickness measurements should be obtained at the prescription point (which is typically at the point of maximum separation, often at the level of the umbilicus) and at other points of interest for possible dose calculations and homogeneity determinations, such as head, neck, mid-mediastinum, mid-lung, pelvis, knee, ankle, etc. Patient height is recorded to determine the appropriate source-to-patient distance to fit the patient within the beam with sufficient margin around the patient (usually greater than 5 cm). Special attention should be paid to the dramatic decrease in dose that can be seen in the field corners for many treatment units when the collimator is in the full open position.

Organ shielding may be considered when necessary [45]. For high dose TBI regimens, total lung dose is limited to about 8 to 10 Gy [46,47]. Lung shielding can be performed by treatment in lateral position with the arms down and/or by use of partial (50% to 80%) transmission blocks. In some cases, shielding of the kidneys, thyroid, lens, liver, or parotid glands is performed. Testes shielding may be considered for non-malignant disease, eg, myelodysplastic syndromes.

D. Simulation of Treatment

For lung or other organ blocking, simulation or other treatment planning is generally done in the treatment position, ie, if the patient is standing for TBI, the simulation should be done in the standing position if possible. As an alternative to CT-simulation in the supine position, lung blocks may be designed on megavoltage radiographs generated by a linear accelerator with the patient in an upright position. If the planning session is performed in another position, positional differences in organ location should be considered and the medical physicist should be consulted. Reference points for block placement at the time of treatment should be marked on the patient's body for reproducibility. If the patient is treated in the lateral position, additional lung shielding may not be necessary.

E. Calculations

Calculations are performed by the medical physicist or his or her designee to determine the beam-on time necessary to achieve the prescribed dose, dose homogeneity, and any other relevant dose points. Consideration should always be given to differences in the patient's separation in different body regions with the resulting dose heterogeneities. For example, adjustments should be considered for overweight patients who can experience severe head and neck mucositis, as well as prescription doses in excess of 20% over the cervical spinal cord when only umbilical separation is used for prescribing dose [48]. These considerations are especially important in patients with a history of prior radiation therapy. A medical physicist or a dosimetrist who did not perform the initial computation should independently check the calculation before the first fraction is delivered. It is recommended here that in-vivo dosimetry be used to assess dose homogeneity. Every effort should be made to maintain dose inhomogeneity to within $\pm 10\%$.

F. Treatment Aids

Special TBI stands, treatment couches, or treatment tables are often used to aid in immobilization, placement of organ shields, and patient support and comfort.

G. Treatment Delivery

TBI containing myeloablative transplant programs typically use fractionated or hyperfractionated regimens (twice a day or 3 times a day) over several days in order to minimize both acute and chronic toxicities and to minimize overall treatment time. Consideration should be given to the time interval between fractions delivered on the same day (typically treatments are separated by a 4- to 6-hour interval). Prior to treatment, any shielding of normal organs should be checked clinically or with portal images. In the setting of low-dose TBI, where total doses are typically only 2 to 4 Gy, organ shielding is usually not used. Dosimetry should be checked against department protocols to verify dose delivery at the extended distances that are used for treatment. A medical physicist should be available during all treatments in case of questions regarding dosimetric details, equipment function, patient setup, etc. Treatments are carried out by the radiation therapist per the [ACR–ASTRO Practice Parameter for Radiation Oncology](#) [41].

A physician should be in close proximity to manage any problems related to treatment. Avoidance of medications that may cause orthostatic hypotension and the administration of IV fluids for hydration or transfusions for anemia may help to prevent syncope or near-syncope episodes if the patient is treated in the standing position.

III. QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL

Application of this practice parameter should be in accordance with the [ACR–ASTRO Practice Parameter for Radiation Oncology](#) [41].

A. Radiation Oncologist

The radiation oncologist should be currently proficient in TBI procedures prior to embarking on any of these regimens. It is encouraged that TBI be performed in high-volume transplant centers.

The responsibilities of the radiation oncologist include:

1. Consultation and decision-making regarding the appropriate course of treatment.
2. Coordination of the patient's care with the transplantation service and other physicians.
3. Oversight and participation in the treatment planning process (immobilization techniques, simulation, block design, prescription, dosimetric and physics review, etc).
4. Review and approval of treatment verification images.
5. Clinical assessment of the patient's tolerance during the treatment course.

B. Qualified Medical Physicist

The responsibilities of the Qualified Medical Physicist include: (see references [21-22] for helpful details relating to this section)

1. Establish and manage the system of dosimetric measurements, calculating and shielding.
2. Establish the system for beam-spoiling designed to adjust the dose at the beam entry surface.
3. Initiate and maintain a quality assurance program for TBI performance.
4. Act as a technical resource for planning of immobilization devices, dosimetry techniques, shielding, dose compensation devices, and bolus methods.
5. Calibrate the external beam delivery system and the in-vivo measurement system.
6. Direct supervision of dosimetry measurements and calculations for TBI delivery.
7. Verify the calculations performed by the dosimetrist.

C. Dosimetrist

The responsibilities of the dosimetrist include:

1. Generation of the dose calculations for treatment.
2. Dosimetry measurements.

D. Radiation Therapist

The responsibilities of the radiation therapist include:

1. Setting up the patient in the treatment position, including using appropriate treatment devices.
2. Verifying that the prescribed and calculated treatment distances match the utilized treatment distances.
3. Performing and reviewing of imaging procedures to verify the setup and blocking, if any.
4. Treating the patient according to the prescription and plan provided.
5. Monitoring and evaluating the patient during the treatments.

E. Nurse

The responsibilities of the nurse may include:

1. Educating the patient and family about the procedures, acute/late side effects, and procedures taken to promote safe/comfortable treatment.
2. Monitoring the patient's tolerance of the procedure to promote adequate supportive care.
3. Communicating any special precautions to the rest of the team regarding the care of immunosuppressed patients.

IV. EQUIPMENT

A treatment room large enough to accommodate extended SSD may be required for treatment of adults using conventional TBI techniques. A backup beam delivery system must be available in case of unanticipated machine failure. High-energy photon beams in the range of 4 to 18 MV are preferred for TBI. Early investigations in the use of helical tomotherapy or volumetric arc therapy for total body or selective total marrow irradiation show promise and may be used, but enrollment in clinical trial(s) evaluating this modality is highly encouraged [49-51]. Additional equipment may include a fluoroscopy or CT simulator, immobilization devices, equipment for the manufacture of shielding, computers for dose calculations, a beam spoiler, custom bolus, custom compensators, and dosimetry and calibration devices.

V. PATIENT AND PERSONNEL SAFETY

A. Safety measures

Safety measures should be in accordance with the [ACR–ASTRO Practice Parameter for Radiation Oncology](#) [41].

B. Special Patient Protection Measures

1. Timing of TBI delivery must be precisely coordinated with chemotherapy regimens, procurement of stem cells, and subsequent stem cell infusion. Confirmation with the transplant team immediately before initiating TBI is important to identify any unanticipated delays or changes to the treatment plan.
2. Charting systems for prescription; delineation of treatment parameters of the setup, including any position settings of the TBI stand; and treatment delivery record, including time of delivery for multiple treatments in a day

3. Physics program for calibrating the treatment machine, independent checking of dose calculations, and monitoring of dose delivery to the patient
4. Visual and audio contact with the patient during treatment

VI. DOCUMENTATION

Reporting should be in accordance with the [ACR–ASTRO Practice Parameter for Communication: Radiation Oncology](#) [43].

VII. EDUCATIONAL PROGRAM

Continuing medical education programs should include radiation oncologists, physicists, dosimetrists, nurses, and radiation therapists. The program should be in accordance with the [ACR Practice Parameter for Continuing Medical Education \(CME\)](#) [52].

VIII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *Position Statement on QC & Improvement, Safety, Infection Control, and Patient Education* on the ACR website (<http://www.acr.org/parameters>).

The Medical Director of Radiation Oncology is responsible for the institution and ongoing supervision of the Continuing Quality Improvement (CQI) program as described in the [ACR–ASTRO Practice Parameter for Radiation Oncology](#) [41]. It is the responsibility of the director to identify problems, see that actions are taken, and evaluate the effectiveness of the actions.

SUMMARY

Total body irradiation is a specialized radiation technique often used prior to HSCT. Delivery of TBI requires knowledge of the clinical indications and specialized treatment setup as well as the presence of dosimetric and physics staff with training in the procedures. Safe and accurate delivery of TBI can be performed with attention to the special indications, specific morbidities, and specialized treatment delivery measurements and techniques required for this procedure.

ACKNOWLEDGEMENTS

This practice parameter was revised according to the process described under the heading *The Process for Developing ACR Practice Parameters and Technical Standards* on the ACR website (<http://www.acr.org/parameters>) by the Committee on Practice Parameters – Radiation Oncology of the ACR Commission on Radiation Oncology in collaboration with the ASTRO.

Collaborative Committee – members represent their societies in the initial and final revision of this practice parameter

ACR

Lesley A. Jarvis, MD, PhD, Chair
Mohammad K. Khan, MD, PhD
Suzanne L. Wolden, MD

ASTRO

Andrea Ng, MD, MPH
Chelsea Pinnix, MD, PhD

Committee on Practice Parameters – Radiation Oncology
(ACR Committee responsible for sponsoring the draft through the process)

Alan C. Hartford, MD, PhD, FACR, Chair
Nathan H. J. Bittner, MD
Chee-Wai Cheng, PhD, FAAPM
Nancy A. Ellerbroek, MD, FACR
Beth A. Erickson, MD, FACR
Lesley A. Jarvis, MD, PhD
Bill W. Loo, MD, PhD
Jeff M. Michalski, MD, MBA, FACR

Christopher H. Pope, MD
Naomi R. Schechter, MD
Nikhil Thaker, MD
Suzanne L. Wolden, MD
Ying Xiao, PhD
Sue S. Yom, MD, PhD
Bassem I. Zaki, MD

Seth A. Rosenthal, MD, FACR, Chair, Commission on Radiation Oncology

Comments Reconciliation Committee

William Small, Jr., MD, FACR, Chair
Debra S. Dyer, MD, FACR, Co-Chair
Neil E. Dunlap, MD
Ralph P. Ermoian, MD
Laura M. Freedman, MD
Lesley A. Jarvis, MD, PhD
Alan C. Hartford, MD, PhD, FACR
William T. Herrington, MD, FACR
Christopher Kelsey, MD
Mohammad K. Khan, MD, PhD
Chi Lin, MD

Andrea Ng, MD, MPH
Rahul R. Parikh, MD
Chelsea Pinnix, MD, PhD
Seth A. Rosenthal, MD, FACR
Peter J. Rossi, MD
Wenyin Shi, MD
Timothy L. Swan, MD, FACR, FSIR
Yolanda Tseng, MD
Paul E. Wallner, DO, FACR
Suzanne L. Wolden, MD

REFERENCES

1. Pasquini MC, Zhu M.C. Current use and outcome of hematopoietic stem cell transplantation: CIBMTR Summary Slides. 2015.
2. Bredeson C, LeRademacher J, Kato K, et al. Prospective cohort study comparing intravenous busulfan to total body irradiation in hematopoietic cell transplantation. *Blood*. 2013;122(24):3871-3878.
3. Copelan EA, Avalos BR, Ahn KW, et al. Comparison of outcomes of allogeneic transplantation for chronic myeloid leukemia with cyclophosphamide in combination with intravenous busulfan, oral busulfan, or total body irradiation. *Biol Blood Marrow Transplant*. 2015;21(3):552-558.
4. Copelan EA, Hamilton BK, Avalos B, et al. Better leukemia-free and overall survival in AML in first remission following cyclophosphamide in combination with busulfan compared with TBI. *Blood*. 2013;122(24):3863-3870.
5. Eroglu C, Pala C, Kaynar L, et al. Comparison of total body irradiation plus cyclophosphamide with busulfan plus cyclophosphamide as conditioning regimens in patients with acute lymphoblastic leukemia undergoing allogeneic hematopoietic stem cell transplant. *Leukemia & lymphoma*. 2013;54(11):2474-2479.
6. Nagler A, Rocha V, Labopin M, et al. Allogeneic hematopoietic stem-cell transplantation for acute myeloid leukemia in remission: comparison of intravenous busulfan plus cyclophosphamide (Cy) versus total-body irradiation plus Cy as conditioning regimen--a report from the acute leukemia working party of the European group for blood and marrow transplantation. *J Clin Oncol*. 2013;31(28):3549-3556.
7. Uberti JP, Agovi MA, Tarima S, et al. Comparative analysis of BU and CY versus CY and TBI in full intensity unrelated marrow donor transplantation for AML, CML and myelodysplasia. *Bone Marrow Transplant*. 2011;46(1):34-43.
8. Chen YB, Lane AA, Logan BR, et al. Impact of conditioning regimen on outcomes for patients with lymphoma undergoing high-dose therapy with autologous hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2015;21(6):1046-1053.
9. Barker C, Wolden SL. Total Body Irradiation. In: Leibel SA, Phillips TL, ed. *Textbook of Radiation Oncology*. 2nd ed. Philadelphia, Pa: WB Saunders; 2010.

10. Alyea E, Neuberger D, Mauch P, et al. Effect of total body irradiation dose escalation on outcome following T-cell-depleted allogeneic bone marrow transplantation. *Biol Blood Marrow Transplant.* 2002;8(3):139-144.
11. Marks DI, Forman SJ, Blume KG, et al. A comparison of cyclophosphamide and total body irradiation with etoposide and total body irradiation as conditioning regimens for patients undergoing sibling allografting for acute lymphoblastic leukemia in first or second complete remission. *Biol Blood Marrow Transplant.* 2006;12(4):438-453.
12. Kelsey CR, Horwitz ME, Chino JP, et al. Severe pulmonary toxicity after myeloablative conditioning using total body irradiation: an assessment of risk factors. *Int J Radiat Oncol Biol Phys.* 2011;81(3):812-818.
13. Kelsey CR, Scott JM, Lane A, et al. Cardiopulmonary exercise testing prior to myeloablative allo-SCT: a feasibility study. *Bone Marrow Transplant.* 2014;49(10):1330-1336.
14. Marnitz S, Zich A, Martus P, et al. Long-term results of total body irradiation in adults with acute lymphoblastic leukemia. *Strahlentherapie und Onkologie : Organ der Deutschen Rontgengesellschaft ... [et al].* 2014;190(5):453-458.
15. Deeg HJ, Sullivan KM, Buckner CD, et al. Marrow transplantation for acute nonlymphoblastic leukemia in first remission: toxicity and long-term follow-up of patients conditioned with single dose or fractionated total body irradiation. *Bone Marrow Transplant.* 1986;1(2):151-157.
16. Thomas ED, Clift RA, Hersman J, et al. Marrow transplantation for acute nonlymphoblastic leukemia in first remission using fractionated or single-dose irradiation. *Int J Radiat Oncol Biol Phys.* 1982;8(5):817-821.
17. Ozsahin M, Belkacemi Y, Pene F, et al. Total-body irradiation and cataract incidence: a randomized comparison of two instantaneous dose rates. *Int J Radiat Oncol Biol Phys.* 1994;28(2):343-347.
18. Abugideiri M, Nanda RH, Butker C, et al. Factors Influencing Pulmonary Toxicity in Children Undergoing Allogeneic Hematopoietic Stem Cell Transplantation in the Setting of Total Body Irradiation-Based Myeloablative Conditioning. *Int J Radiat Oncol Biol Phys.* 2016;94(2):349-359.
19. Carruthers SA, Wallington MM. Total body irradiation and pneumonitis risk: a review of outcomes. *British journal of cancer.* 2004;90(11):2080-2084.
20. Baron F, Maris MB, Sandmaier BM, et al. Graft-versus-tumor effects after allogeneic hematopoietic cell transplantation with nonmyeloablative conditioning. *J Clin Oncol.* 2005;23(9):1993-2003.
21. Hegenbart U, Niederwieser D, Sandmaier BM, et al. Treatment for acute myelogenous leukemia by low-dose, total-body, irradiation-based conditioning and hematopoietic cell transplantation from related and unrelated donors. *J Clin Oncol.* 2006;24(3):444-453.
22. Laport GG, Sandmaier BM, Storer BE, et al. Reduced-intensity conditioning followed by allogeneic hematopoietic cell transplantation for adult patients with myelodysplastic syndrome and myeloproliferative disorders. *Biol Blood Marrow Transplant.* 2008;14(2):246-255.
23. Stelljes M, Bornhauser M, Kroger M, et al. Conditioning with 8-Gy total body irradiation and fludarabine for allogeneic hematopoietic stem cell transplantation in acute myeloid leukemia. *Blood.* 2005;106(9):3314-3321.
24. Tomblyn M, Brunstein C, Burns LJ, et al. Similar and promising outcomes in lymphoma patients treated with myeloablative or nonmyeloablative conditioning and allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant.* 2008;14(5):538-545.
25. Baron F, Storb R, Storer BE, et al. Factors associated with outcomes in allogeneic hematopoietic cell transplantation with nonmyeloablative conditioning after failed myeloablative hematopoietic cell transplantation. *J Clin Oncol.* 2006;24(25):4150-4157.
26. Gyurkocza B, Cao TM, Storb RF, et al. Salvage allogeneic hematopoietic cell transplantation with fludarabine and low-dose total body irradiation after rejection of first allografts. *Biol Blood Marrow Transplant.* 2009;15(10):1314-1322.
27. Buchali A, Feyer P, Groll J, Massenkeil G, Arnold R, Budach V. Immediate toxicity during fractionated total body irradiation as conditioning for bone marrow transplantation. *Radiother Oncol.* 2000;54(2):157-162.
28. Matsuoka S, Okamoto S, Watanabe R, et al. Granisetron plus dexamethasone versus granisetron alone in the prevention of vomiting induced by conditioning for stem cell transplantation: a prospective randomized study. *Int J Hematol.* 2003;77(1):86-90.
29. Peters SG, Afessa B. Acute lung injury after hematopoietic stem cell transplantation. *Clin Chest Med.* 2005;26(4):561-569, vi.
30. Wadleigh M, Ho V, Momtaz P, Richardson P. Hepatic veno-occlusive disease: pathogenesis, diagnosis and treatment. *Curr Opin Hematol.* 2003;10(6):451-462.

31. Hingorani S. Chronic kidney disease in long-term survivors of hematopoietic cell transplantation: epidemiology, pathogenesis, and treatment. *J Am Soc Nephrol*. 2006;17(7):1995-2005.
32. Chemaitilly W, Boulad F, Heller G, et al. Final height in pediatric patients after hyperfractionated total body irradiation and stem cell transplantation. *Bone Marrow Transplant*. 2007;40(1):29-35.
33. Mertens AC, Ramsay NK, Kouris S, Neglia JP. Patterns of gonadal dysfunction following bone marrow transplantation. *Bone Marrow Transplant*. 1998;22(4):345-350.
34. Rizzo JD, Curtis RE, Socie G, et al. Solid cancers after allogeneic hematopoietic cell transplantation. *Blood*. 2009;113(5):1175-1183.
35. Armenian SH, Sun CL, Vase T, et al. Cardiovascular risk factors in hematopoietic cell transplantation survivors: role in development of subsequent cardiovascular disease. *Blood*. 2012;120(23):4505-4512.
36. Kunkele A, Engelhard M, Hauffa BP, et al. Long-term follow-up of pediatric patients receiving total body irradiation before hematopoietic stem cell transplantation and post-transplant survival of >2 years. *Pediatric blood & cancer*. 2013;60(11):1792-1797.
37. Bizzarri C, Pinto RM, Ciccone S, Brescia LP, Locatelli F, Cappa M. Early and progressive insulin resistance in young, non-obese cancer survivors treated with hematopoietic stem cell transplantation. *Pediatric blood & cancer*. 2015;62(9):1650-1655.
38. Oudin C, Auquier P, Bertrand Y, et al. Metabolic syndrome in adults who received hematopoietic stem cell transplantation for acute childhood leukemia: an LEA study. *Bone Marrow Transplant*. 2015;50(11):1438-1444.
39. Van Dyk J, Galvin JM, Glasgow GP, Podgorsak EB. AAPM Report No. 17: The physical aspects of total and half body photon irradiation. 1986. http://www.aapm.org/pubs/reports/RPT_17.pdf. Accessed July 1, 2010.
40. Khan FM, Williamson JF, Sewchand W, Kim TH. Basic data for dosage calculation and compensation. *Int J Radiat Oncol Biol Phys*. 1980;6(6):745-751.
41. American College of Radiology. ACR-ASTRO practice parameter for radiation oncology. 2014; Available at: <http://www.acr.org/~media/7B19A9CEF68F4D6D8F0CF25F21155D73.pdf>. Accessed June 6, 2016.
42. American College of Radiology. ACR-AAPM technical standard for the performance of radiation oncology physics for external beam therapy. 2015; Available at: <http://www.acr.org/~media/EC3F4C2566F24865B1317E2A5B622B8C.pdf>. Accessed June 6, 2016.
43. American College of Radiology. ACR-ASTRO practice parameter for communication: radiation oncology 2015; Available at: <http://www.acr.org/~media/735B26D150674A2291E3E8E69B0C4EF3.pdf>. Accessed June 6, 2016.
44. American College of Radiology. ACR practice parameter on informed consent - radiation oncology. 2012; Available at: <http://www.acr.org/~media/DD525D52FBFD458FBA07B5C8BFCE8F1D.pdf>. Accessed June 6, 2016.
45. Anderson JE, Appelbaum FR, Schoch G, et al. Relapse after allogeneic bone marrow transplantation for refractory anemia is increased by shielding lungs and liver during total body irradiation. *Biol Blood Marrow Transplant*. 2001;7(3):163-170.
46. Della Volpe A, Ferreri AJ, Annaloro C, et al. Lethal pulmonary complications significantly correlate with individually assessed mean lung dose in patients with hematologic malignancies treated with total body irradiation. *Int J Radiat Oncol Biol Phys*. 2002;52(2):483-488.
47. Soule BP, Simone NL, Savani BN, et al. Pulmonary function following total body irradiation (with or without lung shielding) and allogeneic peripheral blood stem cell transplant. *Bone Marrow Transplant*. 2007;40(6):573-578.
48. Chi P, Pinnix C, Dabaja B, Wang C, Aristophanous M, Tung S. Simple field-in-field technique for total body irradiation in large patients. *Med Phys*. 2014;41(6):318.
49. Hui SK, Kapatoes J, Fowler J, et al. Feasibility study of helical tomotherapy for total body or total marrow irradiation. *Med Phys*. 2005;32(10):3214-3224.
50. Wong JY, Rosenthal J, Liu A, Schultheiss T, Forman S, Somlo G. Image-guided total-marrow irradiation using helical tomotherapy in patients with multiple myeloma and acute leukemia undergoing hematopoietic cell transplantation. *Int J Radiat Oncol Biol Phys*. 2009;73(1):273-279.
51. Springer A, Hammer J, Winkler E, et al. Total body irradiation with volumetric modulated arc therapy: Dosimetric data and first clinical experience. *Radiat Oncol*. 2016;11:46.

52. American College of Radiology. ACR practice parameter for continuing medical education (CME). 2016; Available at: <http://www.acr.org/~media/FBCDC94E0E25448DAD5EE9147370A8D1.pdf>. Accessed June 6, 2016.

Development Chronology for this Practice Parameter

*Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

As of May 2015, all practice parameters and technical standards that are collaborative with the American Society for Radiation Oncology (ASTRO) are approved by the ACR Council Steering Committee and the ACR Board of Chancellors and do not go through the ACR Council (ACR Resolution 54, 2015). The effective date is the first of the first month after 60 days of the date the approval was finalized. The effective date is displayed below:

Effective date October 1 of the year noted

- 2001 (Resolution 19)
- Revised 2006 (Resolution 23)
- Revised 2011 (CSC/BOC) – Effective August 1, 2011
- Amended 2014 (Resolution 39)
- Revised 2017 (BOC/CSC); approved May 23, 2017, 60 days following July 22, 2017, effective August 1, 2017